

ENVIRONMENTAL MONITORING PLAN FOR DELAWARE'S INLAND BAYS

REVISED: JANUARY 2018

Compiled by:

Marianne Walch and Andrew McGowan
Delaware Center for the Inland Bays
39375 Inlet Road, Rehoboth Beach, DE 19971
www.inlandbays.org

James M. Eisenhardt
Senior Manager, RK&K
110 S. Poplar Street, Suite 102, Wilmington, DE 19801
www.rkk.com



DELAWARE CENTER FOR THE
INLAND BAYS
Research. Educate. Restore.

Get on board with the bays.
inlandbays.org

Rehoboth | Indian River | Little Assawoman

This page intentionally left blank.

TABLE OF CONTENTS

Table of Contents	i
List of Tables	ii
List of Figures	ii
Acknowledgements	iii
Executive Summary	1
Introduction	
The Inland Bays and Their Watershed	2
Inland Bays Environmental Monitoring Plan	4
Background	4
Objectives	5
Summary of Revision Process	5
Monitoring Plan Updates.....	5
Assessment and Reporting	
Environmental Indicators and State of the Estuary Reports.....	6
TMDL Reporting and Benchmark Goals.....	6
Inland Bays Monitoring Programs	6
Questions Addressed by Monitoring	6
Existing Monitoring Programs.....	8
Data Management and Quality Control	8
Recommendations and Supporting Programs	17
Recommendations Table.....	18
Hydrodynamic/Watershed Model.....	23
Upgrade of Citizen Monitoring Program Database.....	24
Continuous Water Quality Monitoring	25
Monitoring of Submerged Aquatic Vegetation	26
Local Indicators of Sea Level Rise	26
Indian River Inlet Tidal Flushing	27
Monitoring of Oyster Recruitment and Growth	29
Monitoring of Shoreline Condition and Modification	30
Continuation of Tidal Marsh Acreage/Condition as Indicators	31
Estuary Acidification.....	31
Recreational Blue Crab and Clam Harvests	32
Emerging Contaminants.....	32
References.....	33

Appendices:

- APPENDIX A - 2012 CCMP Addendum Actions and Performance Measures
- APPENDIX B - Environmental Monitoring Plan Revision Process and Meeting Notes
- APPENDIX C - Descriptions of Existing Long-Term Monitoring Programs in The Inland Bays Study Area
- APPENDIX D - Assessment, Listing, And Reporting Methodologies for 305(B) Report and 303(D) List Development, 2016

List of Tables:

- TABLE 1 Monitoring and Indicators Matrix
- TABLE 2 Recommendations for Monitoring in the Inland Bays

List of Figures:

- FIGURE 1 Map of the Delaware Inland Bays Watershed
- FIGURE 2 Map: Watershed Land Use, 2012
- FIGURE 3 Hydraulic measurements made at the Indian River Inlet in 2004, showing current measurement transects
- FIGURE 4 Hydraulic measurements made at the Indian River Inlet in 2004

ACKNOWLEDGEMENTS

The development of this update to the Inland Bays Monitoring Plan was funded wholly or in part by the United States Environmental Protection Agency under assistance agreements CE09939912 and CE09939913 to the Delaware Center for the Inland Bays. The contents of this document do not necessarily reflect the views and policies of the Environmental Protection Agency, nor does the EPA endorse trade names or recommend the use of commercial products mentioned in the document.

Input was provided by members of the Inland Bays Monitoring Plan Subcommittee and the Center for the Inland Bays Scientific and Technical Advisory Committee (STAC). The STAC also provided review of the final plan. Leslie Jamka of RK&K provided invaluable assistance in collating questionnaire results and assembling early drafts of the plan.

Special thanks go to those who participated in the Water Quality Monitoring Workgroup meeting held in Dover, DE on July 30, 2015: Robin Tyler, David Wolanski, Michael Bott, Debbie Rouse, Hassan Mirsajadi, John Schneider, Larry Trout, Joanna York, Kevin Brinson, Tina Callahan, Ed Whereat, Bill Ullman, Joe Farrell, Judy Denver, and Scott Andres.

The Delaware Center for the Inland Bays is a nonprofit organization and a National Estuary Program. It was created to promote the wise use and enhancement of the Inland Bays watershed by conducting public outreach and education, developing and implementing restoration projects, encouraging scientific inquiry and sponsoring needed research, and establishing a long-term process for the protection and preservation of the Inland Bays watershed.

EXECUTIVE SUMMARY

The purpose of the Inland Bays Environmental Monitoring Plan (IBEMP) is to track the status and trends of key environmental indicators used to assess the chemical, physical, and biological integrity of the estuary and surrounding study area, and to evaluate whether the goals of the Inland Bays Comprehensive Conservation and Management Plan (CCMP) are being met. It is a comprehensive inventory of existing, new, and proposed monitoring activities to meet these objectives and is intended to guide future research and monitoring efforts. The plan is also intended to lead to increased integration of work and consolidation of resources.

The original Monitoring Plan for the Delaware Inland Bays was written in 1995 and last updated in 1996. Since then, collection of additional relevant parameters has been initiated, responsibilities for collection have changed, and both monitoring technology and the scientific understanding of the Bays have evolved significantly. The Center for the Inland Bays (CIB) manages or supports some of these programs, but most are led and supported financially by academic, county, state, and federal partners. It is the CIB's role to facilitate these partnerships and regularly synthesize, analyze, and report indicator data for the Inland Bays. This updated document provides an opportunity to re-engage stakeholders around its cooperative implementation.

This IBEMP presents an inventory of existing projects and programs that conduct ongoing, long-term environmental monitoring in Delaware's Inland Bays. Many of these programs contribute data that are used by the CIB to develop *State of the Delaware Inland Bays* reports every five years. Other entities produce data that may illuminate progress toward achieving goals of the CCMP, provide data for new environmental models, or may be useful for development of new indicators in the future. For each program we identify CCMP objectives addressed, responsible entities, data collected, data gaps, frequency of data collection and reporting, and how the data are shared, reported and used. The programs are organized into seven sections that cover monitoring of: (1) surface water; (2) groundwater; (3) wastewater; (4) atmospheric deposition; (5) wetlands; and (6) climate.

Of primary importance in this IBEMP are recommendations made for new monitoring programs, or enhancement of existing programs. These recommendations are made based upon critical data gaps (including emerging issues), the availability of new methods or technologies, and/or changes needed to make programs sustainable over the long term.

Highest priority recommendations are:

- Development of a new hydrodynamic/watershed model for the Inland Bays;
- Upgrade of the University of Delaware's Citizen Monitoring Program database to a format that is sustainable long-term and can serve data to the public through STORET and/or the state's Water Quality Portal;
- Long-term, continuous monitoring of dissolved oxygen and chlorophyll at key stations;
- Monitoring of submerged aquatic vegetation in tidal regions of the Inland Bays; and
- Monitoring of local indicators of sea level rise.

Other recommendations, judged to be important but of slightly lower priority, include:

- Continued monitoring of the tidal prism at the Indian River Inlet;
- Long-term monitoring of oyster recruitment and growth in the Bays, particularly as aquaculture begins, and shellfish restoration and enhancement projects are undertaken;
- Shoreline condition and modification monitoring to evaluate the effectiveness of living shoreline initiatives;
- Continued analyses of tidal marsh acreage and condition using GIS methodology established in a 2014 study conducted by the University of Delaware;
- Monitoring of estuary acidification; and
- Monitoring of recreational Blue Crab and Hard Clam harvests from the Inland Bays.

- Build and maintain a list of research and monitoring activities focused on emerging contaminants in the Inland Bays.

For each recommended program, partner organizations or agencies are identified to be responsible for, or participate in, its implementation. Where possible, estimated costs and potential funding sources are provided. Coordination among those organizations involved in data collection, processing and analysis, storage and provision, and presentation is key to the success of monitoring in the Inland Bays. The Center for the Inland Bays Scientific and Technical Advisory Committee (STAC) will be responsible for ensuring the implementation of this plan. This includes engagement of all partners collecting data and ensuring that data are submitted to the CIB for State of the Bays reporting.

The IBEMP is a living document intended to evolve to meet future needs for tracking the status and trends of conditions within the Inland Bays and progress toward meeting the goals of each of the CCMP focus areas. As the CCMP is revised, or new monitoring opportunities or technologies appear, revisions to the plan may need to be made. In order to ensure that monitoring programs are implemented and coordinated, and that the IBEMP is kept up to date, a CCMP/Monitoring subcommittee of the STAC will be responsible for biannual review/update, on a schedule corresponding with the state's development of its 305(b) report to the U.S. EPA.

INTRODUCTION

THE INLAND BAYS AND THEIR WATERSHED

Delaware's Inland Bays (the Bays) refers to all tidal waters and tidal wetlands encompassing the Indian River Bay, Indian River, Rehoboth Bay, and Little Assawoman Bay (Figure 1). The 292 square mile Inland Bays watershed is located in southeastern Sussex County, and drains to 35 square miles of bays and tidal tributaries. Rehoboth Bay and Indian River Bay are tidally connected to the Atlantic Ocean by the Indian River Inlet. Little Assawoman Bay is connected by the Ocean City Inlet, 10 miles to the south in Maryland. As of 2012, agriculture represented the largest land use in the watershed (31%), followed by developed/developing lands (24%), forested lands (17%), wetlands (16%), and open water (12%) (Figure 2).

Historically, the Bays have been extremely dynamic. Prior to the 1930's the Indian River system consisted entirely of freshwater with the only connection to the Atlantic occurring during storm surges when the barrier island was breached, at various locations. The Indian River Inlet, as it exists today, was stabilized in the late 1930's and has deepened over time, passing greater volumes of water and increasing the tidal range of the Bays. This has led to a long-term increase in the salinity of the Bays. The greatest impacts of the salinity shift are evident in the upper




FIGURE 1 – Map of the Delaware Inland Bays Watershed

reaches of the tributaries where tidal freshwater segments have been virtually eliminated. The dynamic flux of the Inland Bays poses an exceptional challenge to those responsible for monitoring the health of the system and establishing scientifically defensible status and trends data and analyses.

The degradation of the Bays has been a gradual process occurring over many decades, and it is anticipated that the recovery process will proceed over a similar period of time. Nutrient contaminated groundwater in the Inland Bays watershed, for example, moves very slowly, and the contamination reaches depths of just over 100 feet. If all contamination of the aquifer were to stop immediately, it is predicted that it would take 75 to over 100 years for replacement water to purge the system and reach the Bays.

While there are other issues raised in the CCMP, two areas of priority problems have been identified in the Inland Bays: eutrophication caused by excessive nutrient loading, and habitat loss and modification.




FIGURE 2 – Map: Watershed Land Use, 2012

THE INLAND BAYS ENVIRONMENTAL MONITORING PLAN

Background

A key requirement of National Estuary Programs is to monitor the effectiveness of actions taken to implement their Comprehensive Conservation and Management Plans (CCMPs). This type of research must include the understanding of the natural variability of the ecosystems and populations that make up the estuary and its watershed.

The original Inland Bays CCMP was developed in 1995 to guide the work of the partners and cooperators charged with its implementation. Subsequent to this, Total Maximum Daily Load (TMDL) regulations for nitrogen and phosphorus were established for Indian River, Indian River Bay, and Rehoboth Bay in 1998, and for Little Assawoman Bay and the major tributaries of the Inland Bays in 2005. In 2008 the Inland Bays Pollution Control Strategy (PCS) was promulgated with the intention to implement the TMDLs. Furthermore, since 1995 population growth and development have brought significant changes to the watershed.

A comprehensive update to the Inland Bays CCMP was published in 2012 as an Addendum to the original plan (Delaware Center for the Inland Bays, 2012). The Addendum includes 10 goals and 81 objectives, organized under eight focus areas:

- Nutrient Management
- Wastewater Management
- Stormwater Management
- Water Quality Management
- Managing Living Resources and Their Habitat
- Planning for Climate Change
- Coordinating Land and Water Use Decisions
- Outreach and Education

Actions that would be required to accomplish the goals and objectives were written, as were Performance Measures that could be used to track progress (Appendix A).

During the Center for the Inland Bays (CIB's) original CCMP development process that culminated in 1995, a *Plan for Inland Bays Environmental Monitoring* was produced and included as Appendix G of that report. The plan assembled the metadata of relevant environmental parameters collected at the time and put forth hypotheses for their change based on CCMP implementation. An updated *Monitoring and Assessment Plan for Delaware's Inland Bays (1996-2000)* was published in December 1996.

Since then, collection of additional relevant parameters has been initiated, responsibilities for collection have changed, and both monitoring technology and the scientific understanding of the Bays have evolved significantly. This requires that the Plan be revised and provides an opportunity to re-engage stakeholders around its cooperative implementation.

The current update to the IBEMP was developed with input from the CIB's Scientific and Technical Advisory Committee (STAC), and the participants of a facilitated Ambient Water Quality Monitoring Plan Workshop held in August 2015. These participants included representatives from the Delaware Department of Natural Resources and Environmental Control (DNREC), the University of Delaware, and the U.S. Geological Survey.

Objectives

The purpose of the Inland Bays Environmental Monitoring Plan (IBEMP) is to monitor the conditions of the Inland Bays Estuary and Watershed used to evaluate the overall effectiveness of the CCMP. It serves as a comprehensive blueprint for monitoring activities that relate to the mission of the CIB.

The IBEMP is a living document intended to evolve to meet future needs for tracking the status and trends of eutrophication and habitat loss/modification within the Inland Bays. It is a comprehensive inventory of projects that are ongoing, or are needed to fill data gaps, to monitor progress toward meeting the CCMP goals. The plan is intended to guide research, monitoring, and assessment efforts, and may lead to increased integration of work and consolidation of resources. If the CCMP is revised or updated, the IBEMP will be reviewed and revised if necessary to reflect that new document starting in 2019.

The primary objectives of the IBEMP are to:

- Identify monitoring needed to track progress toward meeting the goals for improving water quality and living resources within the Bays;
- Measure the effectiveness of CCMP actions in bringing about environmental change;
- Identify projects/programs/agencies that are conducting monitoring that meets these goals;
- Identify gaps where data and information are needed (including emerging issues), and suggest alternatives for filling those gaps where possible through integration of work being carried out under active projects;
- Make recommendations for data synthesis, and for coordination among those organizations involved in data collection, processing and analysis, storage and provision, and presentation;
- Identify funding needs and strategies to address data gaps and implement the IBEMP.

Summary of Plan Revision Process

Stakeholders and partners assisted the CIB in the development of this IBEMP update. The CIB hosted a facilitated workshop to discuss ambient water quality monitoring in the Inland Bays. A meeting of the STAC was also devoted to discussion of the plan. Notes from the workshop and STAC meeting are included in Appendix B.

MONITORING PLAN UPDATES

Periodic reevaluation of the IBEMP must be conducted to ensure that data gaps are addressed. During this reevaluation, monitoring programs will be checked for current relevance, applicability to emerging needs, and improvements in technology. If necessary, the CIB will revise the IBEMP to reflect any updates.

A standing CCMP Monitoring Subcommittee is to be appointed by the Chair of the STAC in 2018. This subcommittee will be responsible for biannual review of, and updates to, the Inland Bays Environmental Monitoring Plan starting 2019, as well as development of strategies for implementation. The biannual period for review was selected to correspond with the State's Combined 305(b) and 303(d) reports to EPA. At a minimum this subcommittee shall include representatives from the Center for the Inland Bays, DNREC Watershed Assessment Section, DNREC Environmental Laboratory, University of Delaware Citizen Monitoring Program, Center for Environmental Monitoring and Analysis (CEMA), and the Delaware Geological Survey.

Schedule for Next Review of the Plan: Spring 2019, Spring 2021.

ASSESSMENT AND REPORTING

ENVIRONMENTAL INDICATORS AND STATE OF THE ESTUARY REPORTS

Conditions in the Inland Bays are dynamic and it is essential to monitor temporal and spatial changes in water quality across the watershed and understand how changes relate to the health of the estuary.

Comprehensive assessments of the condition of the Inland Bays were published in 1995, 2004, 2011, and 2016. The *State of the Delaware Inland Bays* report currently is updated and published every five years. The 2016 report included assessment of 35 individual environmental indicators, which are included in the IBEMP and identified in Table 1. The indicators are used to assess the status and trends of water quality and the health of Inland Bays habitats and living resources. Status and trends are assigned using best professional judgment and reviewed by scientists knowledgeable in these areas. For each indicator, long-term trends are addressed, as well as short-term changes that have occurred since the previous report was published.

The IBEMP is intended, in part, to ensure that long-term collection of data needed to develop these environmental indicators for the Bays is continued.

TMDL REPORTING AND BENCHMARK GOALS

Section 305(b) of the Federal Clean Water Act requires that states prepare and submit a Watershed Assessment Report to EPA on April 1st of every even numbered year. The 305(b) reports and monitoring data are used to compile a list of impaired waters, commonly referred to as the 303(d) list. When waters are identified as impaired on 303(d) lists, a Total Maximum Daily Load (TMDL) must be developed.

All of the 305(b) Reports and 303(d) lists that Delaware has submitted to EPA are available on the DNREC website

(<http://www.dnrec.delaware.gov/swc/wa/Pages/WatershedAssessment305band303dReports.aspx>). Draft Core Documents for the 2016 305(b)/303(d) Integrated Report were posted at the time of completion of this monitoring plan (March 2017). A copy of the assessment, listing, and reporting methodologies, including benchmark goals and hypotheses, is included here as Appendix D.

INLAND BAYS MONITORING PROGRAMS

QUESTIONS ADDRESSED BY MONITORING

Two types of monitoring are used to track progress toward meeting goals of the CCMP – programmatic and environmental. The IBEMP focuses on environmental outcomes, i.e., changes in environmental conditions, ecological functions, and biological populations.

The objectives of the 2012 Addendum to the CCMP were expressed as monitoring questions, which drive much of the environmental monitoring conducted by the CIB and its partners. Understanding of both responses to stressors and natural variability is critical for determining relationships between actions taken and responses within the estuary system. These questions, listed below, are organized to correspond with the CCMP Addendum focus areas. Question numbers are referenced in the Monitoring and Indicator Matrix (Table 1).

MONITORING QUESTIONS

Focus Area: Nutrient Management

1. Do nutrient loadings from nonpoint and atmospheric sources meet established TMDL targets?

Focus Area: Wastewater Management

2. Do nutrient loadings from point sources meet established TMDL targets?
3. Are the frequency and spatial distribution of emerging contaminants of concern increasing, decreasing, or remaining stable?

Focus Area: Stormwater Management

4. Is the aerial coverage of effective impervious surface in the watershed increasing, decreasing, or remaining stable?

Focus Area: Water Quality Management

5. Is tributary water quality improving, declining, or remaining stable?
6. Are phytoplankton biomass levels (as indicated by chlorophyll-a concentrations) above, below, or consistent with established targets?
7. Are water column nutrient concentrations above, below, or consistent with established targets?
8. Is water clarity above, below, or consistent with established targets?
9. Is the areal extent of low dissolved oxygen concentrations increasing, decreasing, or remaining stable?
10. Are the frequency and spatial distribution of macroalgal blooms increasing, decreasing, or remaining stable?
11. Are the concentrations and spatial distribution of fecal bacteria increasing, decreasing, or remaining stable?
12. Do concentrations of toxics in water and sediment meet water quality standards?

Focus Area: Managing Living Resources and their Habitat

13. Is the acreage of bay grasses increasing, decreasing or remaining stable?
14. Is the acreage and condition of freshwater wetlands, including isolated wetlands, increasing, decreasing or remaining stable?
15. Is the acreage and condition of tidal wetlands increasing, decreasing or remaining stable?
16. Is the average width of vegetated buffers on waterways increasing, decreasing, or remaining stable?
17. Is the percentage of hardened shorelines in the Inland Bays increasing, decreasing, or remaining stable?
18. Are populations of migratory fish increasing, decreasing or remaining stable?
19. Is the acreage of approved shellfishing waters increasing, decreasing or remaining stable?
20. Are the density and distribution of economically important shellfish (oysters, hard clams, blue crabs) increasing, decreasing or remaining stable?
21. Are the density and distribution of invasive plant and animal species increasing, decreasing or remaining stable?

Focus Area: Planning for Climate Change

25. Are water levels in the estuary increasing, decreasing, or remaining stable?
26. Are water temperatures increasing, decreasing, or remaining stable?
27. Are pH conditions in the estuary increasing, decreasing, or remaining stable?
28. Are shifts of dominant aquatic species changing over time in response to long-term temperature, pH, or salinity changes?

Focus Area: Coordinating Land and Water Use Decisions

29. Is the acreage of natural habitat protected or restored increasing, decreasing, or remaining stable?
30. Are populations of Species of Greatest Conservation Need (as defined by the Delaware Wildlife Action Plan) in the watershed increasing, decreasing or remaining stable?

EXISTING MONITORING PROGRAMS

The IBEMP provides a framework that builds upon existing programs that are conducting ongoing, long-term environmental monitoring within the Delaware's Inland Bays study area. Many of these programs contribute data that are used to develop *State of the Delaware Inland Bays* reports every five years. Others produce data that may illuminate progress toward achieving goals of the CCMP and/or Pollution Control Strategy, provide data for new environmental models, or may be useful for development of new indicators in the future.

CCMP objectives and monitoring questions addressed by these programs, responsible entities, data collected, data gaps, frequency of data collection and reporting, and how the data are shared, reported and used are summarized in the Monitoring and Indicators Matrix (Table 1).

The monitoring programs listed in the Matrix are organized into the following sections:

1. Surface Water
2. Groundwater
3. Wastewater
4. Atmospheric Deposition
5. Wetlands
6. Living Resources
7. Climate

Many of the monitoring programs in Table 1 are described in greater detail in Appendix C. Cross-references to sections of the Appendix are provided in column two of the Table.

The expectation and recommendation of this Plan is that these programs will continue to be funded and conducted long-term. Coordination, collaboration, and long-term support for these monitoring programs are critical for the success of the IBEMP.

The CIB funds and leads the following monitoring programs:

- Long-term salt marsh monitoring
- Seaweed abundance
- Inshore fish and blue crab surveys
- Horseshoe crab surveys and tagging

Responsibility for implementation of the remaining programs lies primarily with other entities, as noted in Table 1. The CIB will work with these entities, as needed, to facilitate partnerships and funding that ensure the sustainability of these programs. Through the STAC and state/regional workgroups (such as the Delaware Environmental Sensing Workgroup), the CIB will ensure that data gaps are discussed and addressed in each biannual review of the IBEMP. The CIB also will continue to synthesize Inland Bays monitoring data and communicate findings to the public, stakeholders, and decision makers through *State of the Bays* reports and other media.

DATA MANAGEMENT AND QUALITY CONTROL

The data managers identified for each monitoring program (Table 1) are responsible for the following:

- Collection and analysis of data according to existing or updated monitoring plans. Changes in monitoring frequency or protocols must be communicated to the CIB, and included in any updates to the *Environmental Monitoring Plan for Delaware's Inland Bays*.

- Providing data to the CIB and/or its partners upon request for use in indicator reports or research projects.
- Quality Assurance Plans must be kept up to date and provided to data users (including the CIB) upon request.
- All monitoring programs funded partially or in whole by the U.S. EPA must have an up to date, approved Quality Assurance Project Plan (QAPP). QAPPs are to be updated every three years. A copy of the approved QAPP must be provided to the CIB for its records.

TABLE 1 - Monitoring and Indicators Matrix (◆ = used to develop indicator for State of the Bays reports)

Monitoring Program	Appendix E Cross-reference	CCMP Objectives/ Monitoring Questions	Indicators & Measures	Data Collected	Collection Frequency	Record of Collection	Responsible Entity/Entities	Frequency of Reporting	Sharing/Reporting	Gaps & Funding Needs
T. SURFACE WATER MONITORING PROGRAMS										
Ambient Water Quality, State of Delaware General Assessment Monitoring Network (GAMN) ◆	E.1.1	CCMP: Water Quality Management, Objectives 1, 3, and 6 Questions: 1, 3, 5, 6, 7, 8, 9, 11, 12, 26, 27, 28	Comparison with TMDL targets for nitrogen, phosphorus, and bacteria. Data used by CIB to develop a Water Quality Index (WQI) to indicate suitability for eelgrass reestablishment.	TP, ortho-P, ammonia N, NO ₃ -NO ₂ , TN, TOC, DOC, Chl a, BOD, DO, TSS, alkalinity, hardness, pH, conductivity, salinity, temperature, Secchi depth, light attenuation, turbidity, chloride, total Enterococcus, Cu, Pb, Zn, As	Most stations monitored 6 times/year for three years, then 12 times/year for two years.	1998 to present.	DNREC Division of Water, Environmental Laboratory DNREC Division of Watershed Stewardship, Watershed Assessment Section	Biannually	GAMN data entered into STORET and publicly available via the Delaware Water Quality Portal. Data published biannually in 305(b) and 303(d) combined reports. Indicator in State of the Delaware Inland Bays reports.	Continue long-term program funding. Continuous/high frequency monitoring for DO and other parameters that fluctuate significantly over short time scales. New hydrodynamic/watershed model for the Inland Bays. Monitoring of estuary acidification. Emerging contaminants of concern.
Ambient Water Quality, Citizen Monitoring Program (CMP) ◆	E.1.2	CCMP: Water Quality Management, Objectives 1, 3, and 6; Outreach and Education, Objective 4 Questions: 5, 6, 7, 8, 9, 11, 26, 27, 28	Data used by CIB to develop a Water Quality Index (WQI) to indicate suitability for eelgrass reestablishment. Data included in state's Combined 305(b) Report and 303(d) List, but not used in listing determinations.	Ortho-P, ammonia N, NO ₃ -NO ₂ , DO, Chl a, TSS, pH, conductivity, salinity, temperature, Secchi depth, rainfall	weekly to biweekly, dependent upon parameter	Varies by station. Longest records begin in 2000.	University of Delaware, Sea Grant Marine Advisory Service	Semi-monthly summary reports	Summary reports posted online. Indicator in State of the Delaware Inland Bays reports.	Additional funding to support continued nutrient analyses (<\$10K/yr). Funding needed to upgrade CMP database to make data publicly available via STORET/WQ Portal (\$50-100K).
Fecal contaminants, Citizen Monitoring Program (CMP) ◆	E.1.2	CCMP: Water Quality Management, Objectives 1, 3, and 6; Managing Living Resources, Objective 5; Outreach and Education, Objective 4 Questions: 11	Indicator derived from percentage of summer samples exceeding safe swimming standard. Harmful algal bloom and TE data used to support State's Shellfish Monitoring Program.	Total Enterococcus	Biweekly	Varies by station. Longest records begin in 2000.	University of Delaware, Sea Grant Marine Advisory Service	Semi-monthly summary reports by CMP. Every five years in CIB indicator reports.	Summary reports posted online by CMP. Indicator in State of the Delaware Inland Bays reports.	Along with ambient WQ, data should be made publicly available via STORET/WQ Portal.
Fecal contaminants, State of Delaware Coliform Monitoring Program	E.1.3	CCMP: Water Quality Management, Objectives 1, 3, and 6; Managing Living Resources, Objective 5; Outreach and Education, Objective 4. Questions: 11, 19	Assess the suitability of the Inland Bays waters for shellfish harvest using federal standards.	Total/fecal coliforms	9-10 times per year	2008 to present. Fecal coliform collection began in 2016, with plans to succeed total coliforms as the method to assess the suitability of waters.	DNREC Division of Watershed Stewardship Shellfish Program	Changes in shellfish advisories reported as made.	Shellfish advisory interactive map on DNREC website.	Refinement of source tracking. Continue current funding.
Stream/Tide Gaging	E.1.4	CCMP: Water Quality Management, Objectives 1 and 6; Planning for Climate Change, Objective 1. Questions: 1, 25	Tide elevation. Real-time and long-term, current and historical streamflow.	Discharge/flow, tide elevation.	Continuous	Varies by station. Longest record begins 1985.	MD-DE-DC Water Science Center	Daily summary data online	Stream/tide data shared online by USGS. Delaware data also available through the Delaware Environmental Observing System (DEOS) site.	Continue funding to maintain existing stream and tide gages. Need for monitoring of local indicators of sea level rise, including a flood monitoring network.
Tidal Flushing at Indian River Inlet ◆	—	CCMP: Water Quality Management, Objective 1. Questions: 5, 28	Tidal prism calculations. Biological Index, derived to categorize stream water quality according to condition (excellent to severely degraded)	Tidal range, surface area of basin calculated from transects and bathymetry data.	No set frequency. No plans for future monitoring.	1939-2004 (irregularly)	U.S. Army Corps of Engineers, Coastal Planning Section	No set frequency	Technical report submitted to CIB for each monitoring event. Indicator in State of the Delaware Inland Bays reports.	Funding to repeat tidal prism measurements.
Biological assessments of water quality in wadable streams	E.1.5	CCMP: Water Quality Management, Objectives 2 and 3. Questions: 5	Benthic macroinvertebrates, periphyton		Biannually	2000 to present	DNREC Division of Water, Environmental Laboratory	Biannual	Used in 303(d) listings.	Continue current funding

TABLE 1 - Monitoring and Indicators Matrix (◆ = used to develop indicator for State of the Bays reports)

Monitoring Program	Appendix E Cross-reference	CCMP Objectives/ Monitoring Questions	Indicators & Measures	Data Collected	Collection Frequency	Record of Collection	Responsible Entity/Entities	Frequency of Reporting	Sharing/Reporting	Gaps & Funding Needs
2. GROUNDWATER PROGRAMS										
Delaware Groundwater Monitoring Network	E.2.1	CCMP: Wastewater Management, Objective 3; Water Quality Management, Objectives 1, 3, and 6; Planning for Climate Change, Objective 1. Questions: 25, 28	Water levels in major aquifers, saltwater intrusion.	Long time-series of water levels in 15 wells in the Inland Bays watershed. Salinity sensors installed in three wells. Groundwater quality data collected when funded.	Continuous	Varies by well. Records for the oldest wells go to 1957.	Delaware Geological Survey	Continuous	Data and reports shared on DGS website.	Funding to support continued groundwater quality data collection.
Agricultural Shallow Groundwater Monitoring Network	E.2.2	CCMP: Nutrient Management, Objective 1; Wastewater Management, Objective 3; Water Quality Management, Objective 3. Questions: 1	Trends in nitrate concentrations.	Nutrients, major ions, pesticides, and groundwater age.	Continuous	2014 to present.	USGS and Delaware Department of Agriculture	Annually	Reports available from Delaware Dept. of Agriculture.	Continue current funding.
Ambient Groundwater Quality, Public Water Supply Wells	E.2.3	CCMP: Wastewater Management, Objectives 2 and 3; Water Quality Management, Objective 3. Questions: 12	Results evaluated with respect to Primary Maximum Contaminant Levels (PMLs), Secondary Maximum Contaminant Levels (SMCLs), and Health Advisories (HAs) for public water-supply systems.	Chemical data collected from public water-supply wells. Data drawn from DHSS's Safe Drinking Water Information System (SDWIS) database, DNREC's Source Water Assessment and Protection Program (SWAPP) database, and Tidewater Utilities.	Analyses conducted biannually	2008 to present	DNREC Division of Water, Water Supply Section	Biannually	306(b) Groundwater-Quality Assessment Based on Public-Well Data reports	Continue current funding.
3. WASTEWATER MONITORING PROGRAMS										
Point Source Wastewater Discharges ◆	E.3.1	CCMP: Wastewater Management, Objective 2; Water Quality Management, Objectives 1, 2, and 3. Questions: 2, 3	Flow, nutrient loads, eutrophication indicators. Data used to assess progress toward TMDL goals.	Flow, BOD, TSS, TP, TN, pH, DO. Some facilities also monitoring fractions of P and N.	Varies by permitted facility/outfall	Varies by permitted facility	DNREC Division of Water, Surface Water Discharges Section	Annually	Data must be requested from DNREC. Point source nutrient loads indicator in State of the Delaware Inland Bays reports.	Continue current funding.
Land Application of Wastewater	E.3.2	CCMP: Nutrient Management, Objective 1; Wastewater Management, Objectives 1 and 3; Water Quality Management, Objectives 1, 2, and 3. Questions: 1, 2, 3	Nutrient loads, eutrophication indicators. Maintenance of soil function. Compliance with drinking water standards.	Effluent: TSS, BOD, TN, TP (and fractions of nitrogen), pH, DO. Groundwater: water table depth, soluble constituents (under and adjacent to spray site).	Varies by permitted facility	Varies by permitted facility	DNREC Division of Water, Groundwater Discharges Section, Large Systems Branch	Annually	Compliance/monitoring reports prepared by permittees.	Continue current funding.
Nutrient Management Practices ◆	—	CCMP: Nutrient Management, Objective 1; Wastewater Management, Objectives 1 and 3; Stormwater Management, Objective 1; Water Quality Management, Objective 2. Questions: 1	Progress toward PCS goals	Agricultural nutrient management practices, EDU's converted from septic to sewer, acres of stormwater retrofits.	Annually	2005 to present	U.S. Department of Agriculture, DNREC Division of Watershed Stewardship, DNREC Division of Water, Sussex Conservation District, Sussex County	Every five years	Indicator in State of the Delaware Inland Bays report.	Incomplete data collection/reporting on agricultural practices in the watershed.

TABLE 1 - Monitoring and Indicators Matrix (used to develop indicator for State of the Bays reports)

Monitoring Program	Appendix E Cross-reference	CCMP Objectives/Monitoring Questions	Indicators & Measures	Data Collected	Collection Frequency	Record of Collection	Responsible Entity/Entities	Frequency of Reporting	Sharing/Reporting	Gaps & Funding Needs
4. Atmospheric Deposition of Nutrients	E.4.1	CCMP: Water Quality Management Objectives 1 and 2. Questions: 1	Wet and dry deposition rates of N and P species to open waters and contiguous marshes in the Inland Bays.	Precipitation, free acidity (H ⁺), pH, conductance, Ca ²⁺ , Mg ²⁺ , Na ⁺ , K ⁺ , SO ₄ ²⁻ , NO ₃ ⁻ , Cl ⁻ , NH ₄ ⁺ .	Daily within 24 hours of the start of precipitation.	1992 to present	University of Delaware, College of Earth, Ocean, and Environment	5-6 month time lag between sample collection and posting on website.	Data from the Lewes station DE07 accessed through the main NADP/ARMON website or the AIRMON Data Retrieval Site. Indicator in State of the Delaware Inland Bays report.	Continue current funding to maintain local station.
5. WETLANDS MONITORING PROGRAMS										
State of Delaware Wetlands Monitoring and Assessment Program	E.5.1	CCMP: Managing Living Resources and Their Habitat, Objectives 2 and 6; Planning for Climate Change, Objective 1; Coordinating Land and Water Use Decisions, Objective 2. Questions: 14, 15	Wetland acreage, condition, and major stressors. DECAP scores for nontidal wetland functions (habitat, hydrology, buffers, soil cycling). MidTRAM scores for condition of aquatic emergent tidal wetlands, validated with biological data based on bird community and biomass levels.	Delaware Comprehensive Assessment Procedure (DECAP) and Delaware Rapid Assessment Procedure (DERAP) for nontidal wetlands. MidAtlantic Tidal Rapid Assessment Method (MidTRAM) protocol for tidal wetlands.	Rotating basin approach planned following initial assessments.	Nontidal wetlands: 2005-2006 Tidal wetlands: 2008	DNREC Division of Watershed Stewardship, Wetlands Monitoring and Assessment Program	Final reports issued following assessment of each basin.	Data managed by DNREC Wetland Monitoring and Assessment Program.	DNREC should follow through on plans to make wetlands data available through STORET.
Long-term Salt Marsh elevation	E.5.2	CCMP: Managing Living Resources and Their Habitat, Objectives 2 and 6; Planning for Climate Change, Objective 1. Questions: 15, 25	Overall marsh height change, changes in the marsh height, as a result of surface accretion/erosion, changes in the root zone.	Sediment elevation, surface accretion, water level	Semi-annually	2008 to present	Delaware Center for the Inland Bays	Every five years	Project technical report published by the CIB.	Continue current support for three Inland Bays stations. Collection of vegetation data from these sites would be useful.
Salt Marsh Acreage and Condition		CCMP: Managing Living Resources and Their Habitat, Objectives 2 and 6; Planning for Climate Change, Objective 1; Coordinating Land and Water Use Decisions, Objective 2. Questions: 15	Historic trends in the extent and condition of the tidal marshes of the Inland Bays	Extent of vegetated marsh, fractured pooling, ditching, and hardening. Data derived from geospatial analysis of aerial imagery and land cover data.	N/A	Study completed in 2014. Documented data from a number of years between 1937 and 2007.	University of Delaware Water Resources Agency Delaware Center for the Inland Bays	Every five years for State of the Delaware Inland Bays reporting.	Technical report produced by the University of DE. Geospatial data available. Acreage and fractured pooling used as indicators in the State of the Delaware Inland Bays report.	Funding to continue future analyses of tidal marsh acreage and condition using updated imagery and land cover data.
Shoreline condition		CCMP: Managing Living Resources and Their Habitat, Objective 2; Planning for Climate Change, Objective 1. Questions: 17	Condition of tidal shorelines.	Riparian land use, bank condition, shoreline features (including hard structure)	N/A	Behoboth Bay: 2006, Indian River Bay: 2012	DNREC Division of Watershed Stewardship, Wetlands Monitoring and Assessment Program Delaware Center for the Inland Bays	N/A	Current data available on WIMS Center for Coastal Resources Management Shoreline Inventory website. Summary report published for each bay inventory.	Shoreline condition data collection for Little Assawoman Bay. Ongoing updates to shoreline data needed to assess progress of Living Shoreline Initiative.

TABLE 1 – Monitoring and Indicators Matrix (♦ = used to develop indicator for State of the Bays reports)

Monitoring Program	Appendix E Cross-reference	CCMP Objectives/ Monitoring Questions	Indicators & Measures	Data Collected	Collection Frequency	Record of Collection	Responsible Entity/Entries	Frequency of Reporting	Sharing/Reporting	Gaps & Funding Needs
♦ Land Use Land Cover ♦ 6. LIVING RESOURCES AND HABITATS	—	CCMP: Stormwater Management, Objective 1; Managing Living Resources and Their Habitat, Objective 2; Planning for Climate Change, Objective 1; Coordinating Land and Water Use Decisions, Objectives 1 and 2; Outreach and Education, Objective 5. Questions: 4, 14, 15, 16, 29	Land use change. Impervious surface coverage, arisan width of forested buffers on croplands	Aerial estimates	Every five years	1992 to present.	State of Delaware, Office of Management and Budget, Delaware Geographic Data Committee	Five-year updates	Delaware First Map. Indicator in State of the Delaware Inland Bays report.	More frequent updates of impervious cover layer.
Human Population ♦	—	CCMP: Wastewater Management, Objectives 1 and 3; Coordinating Land and Water Use Decisions, Objective 1. Questions: 4, 29	Resident and seasonal population. Sussex County and watershed	U.S. census, wastewater flows	Every ten years	County: 1790 to present Watershed: 1990 to present	U.S. Census Bureau, Sussex County	Every five years.	Indicator in the State of the Delaware Inland Bays report.	A better estimate of seasonal population is needed.
Natural Aquatic Resource Surveys (NARS)	E.6.1	CCMP: Water Quality Management, Objectives 1 and 3; Managing Living Resources and Their Habitat, Objective 2. Questions: 5, 6, 7, 8, 9, 11, 12, 14, 15, 18, 21, 26, 27, 28	Four individual surveys that are implemented on a rotating basis - National Coastal Condition Assessment, National Lakes Assessment, National Rivers & Streams Assessment, and National Wetland Condition Assessment. Each has Biological, chemical, physical, and human health indicators.	Biological (benthic macroinvertebrates, Chl a, zooplankton, fish diversity), chemical (DO, salinity, pH, nutrients, sediment contaminants/toxicity), physical (clarity, habitat complexity and disturbance), recreational/human health (legal toxins, cyanobacteria, enterococci)	Five-year cycles	2007 to present	U.S. Environmental Protection Agency, Office of Wetlands, Oceans, and Watersheds DNR/REC Division of Water, Environmental Laboratory Section	Five-year cycles for reporting status, trends for each survey.	Data, summary reports, and manuals shared on EPA website.	Continue long-term funding of program.
Natural Habitat Protection/ Restoration ♦	—	CCMP: Managing Living Resources and Their Habitat, Objective 2. Questions: 29	Cumulative acres	Data retrieved from U.S. EPA National Estuary Program Online Reporting Tool (NEPORT) Habitat Reporting	Annually	2003 to present	Delaware Center for the Inland Bays	Annually	Indicator in State of the Delaware Inland Bays reports.	NEPORT may not capture all projects in the watershed.
Vegetation Community and Land Cover Mapping	E.6.2	CCMP: Managing Living Resources and Their Habitat, Objectives 2 and 6. Questions: 14, 30	Rarity and status of vegetation communities.	Aerial imagery analyses of public lands, combined with field data.	Delaware Statewide Vegetation Community Map updated approximately monthly.	Analyses published for 2011-2013.	DNR/REC Wildlife Species Conservation and Research Program	Monthly updates	Data held by DNR/REC Wildlife Species Conservation and Research Program, and the University of Delaware Water Resources Agency.	Continue current program funding.
Seaweed Abundance ♦	E.6.3	CCMP: Water Quality Management, Objectives 3, 5 and 6; Managing Living Resources and Their Habitats, Objective 5. Questions: 5, 10	Macroalgae density	Volume of macroalgae collected at fixed stations by grapping hook.	Monthly, May to September	1999, 2009, 2011, Annually since 2017.	Delaware Center for the Inland Bays	Annually	Prior to 2017, reports for each survey published by DNR/REC. Currently published on CIB website. Status and trend data included in State of the Delaware Inland Bays reports.	Coverage could be increased by citizen scientists.

TABLE 1 - Monitoring and Indicators Matrix (◆ = used to develop indicator for State of the Bays reports)

Monitoring Program	Appendix E Cross-reference	CCMP Objectives/ Monitoring Questions	Indicators & Measures	Data Collected	Collection Frequency	Record of Collection	Responsible Entity/Entities	Frequency of Reporting	Sharing/Reporting	Gaps & Funding Needs
Coastal Finfish Assessment ◆	E.6.4	CCMP: Managing Living Resources and Their Habitat, Objectives 3, 4, 5, and 6; Questions: 18,21,28	Abundance and distribution of recreationally important finfish species. Annual young-of-the-year indices, catch per tow. Blue Crab abundance.	Other trawl surveys (Indian River and Rehoboth Bays). Count by species, fork length, surface temperature, salinity, and DO, tidal stage, weather conditions, water depth, engine speed.	Monthly, April to October	1986 to present.	DNREC Division of Fish and Wildlife, Fisheries Section	Annually	Annual summary reports published by DNREC DFW. Status and trends for indicator species reported in the State of the Delaware Inland Bays reports.	Continue current program funding.
Inshore Fish and Blue Crabs	E.6.5	CCMP: Managing Living Resources and Their Habitat, Objectives 3, 4, 5, and 6; Outreach and Education, Objective 4; Questions: 18,20,21,28	Abundance and distribution of juvenile fish and Blue Crabs.	Same counts of fish by species; fish lengths; Blue Crab counts by size class, salinity, DO, temperature, wave height, antecedent precipitation, wind speed, tide cycle.	Semi-monthly from April to October.	2011 to present.	Delaware Center for the Inland Bays	Annually	Annual data reports published on CIB website. Five-year trend analyses published by CIB.	Continue current funding and level of effort.
Recreational Fishing ◆	E.6.6	CCMP: Managing Living Resources and Their Habitat, Objective 3; Coordinating Land and Water Use Decisions, Objectives 1 and 3; Outreach and Education, Objective 5; Questions: 18,28	Marine recreational fishing catch and effort.	Catch rates and effort, by species. Landings per year. Harvester intercept surveys conducted at Indian River Inlet.	Annually	2004 to present	NOAA Fisheries Service, Marine Recreational Information Program (MRIP) DNREC Division of Fish and Wildlife	Annually	MRIP reports, DNREC DFW reports. Status and trends for indicator species reported in the State of the Delaware Inland Bays reports.	Monitoring of recreational blue crab and hard clam harvests.
Hard Clam Landings ◆	-	CCMP: Managing Living Resources and Their Habitat, Objective 5; Coordinating Land and Water Use Decisions, Objectives 1 and 3; Outreach and Education, Objective 5; Questions: 20	Number of clams	Commercial and recreational catch	Annually	1943 to present	DNREC Division of Fish and Wildlife, Fisheries Section	Annually	Data available annually from the DNREC Div. of Fish and Wildlife. Used as an indicator in State of the Delaware Inland Bays reports.	Continue current program funding.
Fish Tissue ◆	E.1.6	CCMP: Wastewater Management, Objective 2; Water Quality Management, Objective 4; Managing Living Resources and Their Habitat, Objective 5; Questions: 3,12	Fish consumption advisories.	Contaminants in fish and shellfish.	Varies	1992-present.	DNREC Division of Fish and Wildlife & Division of Watershed Stewardship Delaware Health and Social Services, Division of Public Health	Annual updates to advisories.	Fish consumption advisory reports. Number of advisories used as indicator in State of the Delaware Inland Bays report.	No data to date have led DNREC and DHSS to conclude that an advisory is needed in the Inland Bays as a result of contaminant sources within the Inland Bays.
Fish Kills ◆	-	CCMP: Water Quality management, Objectives 5 and 6; Questions: 5,9	Number of kills reported.	Reports to DNREC	As reported	1981 to present	DNREC Division of Fish and Wildlife, Fisheries Section	Annually	Data shared by DNREC upon request. Indicator in the State of the Delaware Inland Bays reports.	Continue collection of fish kill data.
Shellfish Abundance (Hard Clams, Scallops, Oysters) ◆	E.6.7	CCMP: Managing Living Resources and Their Habitat, Objective 5; Questions: 20	Hard Clam/Scallop abundance in Indian River and Rehoboth Bays, health of the Inland Bays clam shellfishery.	Density and distribution of hard clams and scallops in Indian River and Rehoboth Bays.	No set frequency	1987, 1976, 2011	DNREC Division of Watershed Stewardship	At conclusion of each survey.	Technical report published by DNREC at conclusion of survey. Data included in State of the Delaware Inland Bays reports.	Hard Clam data collection repeated at least every five years, for inclusion in State of the Delaware Inland Bays reports. Long-term monitoring of oyster recruitment and growth. No long-term surveys currently exist.

TABLE 1 - Monitoring and Indicators Matrix (◆ = used to develop indicator for State of the Bays reports)

Monitoring Program	Appendix E Cross-reference	CCMP Objectives/ Monitoring Questions/ Resources and Their Habitats, Objective 2; Outreach and Education, Objective 4. Questions: 17, 28, 30	Indicators & Measures	Data Collected	Collection Frequency	Record of Collection	Responsible Entity/Entities	Frequency of Reporting	Sharing/Reporting	Gaps & Funding Needs
Horseshoe Crabs	E.6.8	CCMP: Managing Living Resources and Their Habitats, Objective 2; Outreach and Education, Objective 4. Questions: 17, 28, 30	Spawning surveys conducted on selected beaches.	Counts of spawning crabs along beach transects, sex ratios, wave height, wind speed, salinity, water temperature.	Semi-monthly during spawning season, around full and new moons.	2012 to present.	Delaware Center for the Inland Bays	Annually	Annual data reports published on CIB website. Five-year trend analyses.	Protocol changed in 2015 to facilitate comparisons with Delaware Bay surveys. Inland Bays data should be incorporated into regional datasets.
Breeding Bird Atlas	E.6.9	CCMP: Managing Living Resources and Their Habitats, Objective 2; Coordinating Land and Water Use Decisions, Objective 2; Outreach and Education, Objective 4. Questions: 20, 30	Trends in breeding bird populations, distribution, and diversity that occur with changes in land use, habitats, and climate.	Verified evidence of breeding, all species, within 10-sq. mi. blocks.	No set frequency. Two atlas surveys conducted to date, 25 years apart.	1993-1997; 2008-2012.	DNREC Division of Fish and Wildlife, Wildlife Species Conservation & Research Program	Report after each 5-year survey period. No set frequency for surveys.	Published as a book at the end of each atlas survey.	As more data are collected, long-term trend analyses to correlate changes with environmental factors such as disappearance of interior forest and climate change.
◆ Mid-Winter Waterfowl	E.6.10	CCMP: Managing Living Resources and Their Habitats, Objective 2; Coordinating Land and Water Use Decisions, Objective 2. Questions: 13, 14, 15, 30	Wintering populations of ducks and geese, by species.	Aerial survey counts for each species, by zone.	Annually	1974 to present	DNREC Wildlife Species Conservation and Research Program	Annually	Data shared through Delaware Open Data Portal. USFWS Atlantic Flyway annual reports. Status and trends for selected indicator species reported in State of the Delaware Inland Bays reports.	Continue funding for January surveys each year, at a minimum.
Bald Eagle and Osprey Nesting	E.6.11	CCMP: Managing Living Resources and Their Habitats, Objective 2; Coordinating Land and Water Use Decisions, Objective 2; Outreach and Education, Objective 4. Questions: 30	Bald Eagle and Osprey populations, measured by number of active nests.	Aerial surveys of active nests.	Annually	Bald Eagles: 1997 to present. Ospreys: 1991 to present.	DNREC Division of Fish and Wildlife, Wildlife Species Conservation & Research Program	Annually	Osprey data shared through OspreyWatch program. Indicator in State of the Delaware Inland Bays reports.	Funding to continue at least annual surveys for both species. Some years have been missed.
◆ Submerged Aquatic Vegetation	—	CCMP: Water Quality Management, Object 6; Managing Living Resources and Their Habitat, Objective 1. Questions: 13	Acres, distribution, species.	No current data collection.	No current data collection	Last reported natural eelgrass bed in 1975. Acres reported since then are anecdotal.	None currently.	N/A	Reported as an indicator in State of the Delaware Inland Bays reports.	No current program exists to survey map SAV in the Inland Bays. Periodic surveys should be implemented.

TABLE 1 - Monitoring and Indicators Matrix (◆ = used to develop indicator for State of the Bays reports)

Monitoring Program	Appendix E Cross-reference	CCMP Objectives/ Monitoring Questions	Indicators & Measures	Data Collected	Collection Frequency	Record of Collection	Responsible Entity/Entities	Frequency of Reporting	Sharing/Reporting	Gaps & Funding Needs
Sea Level ◆	—	CCMP: Planning for Climate Change, Objective 1 Questions: 25	Change in mean sea level at Lewes, DE tide gage.	water level	Continuous	1919 to present	NOAA National Ocean Service	Real-time	Data shared on NOAA Tides and Currents website. Status and trends for this station included as an indicator in the State of the Delaware Inland Bays report.	Lewes gage was offline from 2011 to 2015. Went back online in 2016. Continue long term funding for this station. Need for monitoring of local indicators of sea level rise, including a flood monitoring network.
Ocean Acidification ◆	—	CCMP: Planning for Climate Change, Objective 1 Questions: 27, 28	Change in ocean pH, measured at Oahu, Hawaii.	pH	Monthly	1988 to present	Hawaii Ocean Time-Series Program	Monthly	Data shared on HOT website. Status and trends for this station included as an indicator in the State of the Delaware Inland Bays report.	Data on estuary acidification.
Atmospheric Carbon Dioxide Concentration ◆	—	CCMP: Planning for Climate Change, Objective 1 Questions: 25, 27, 28	Change in CO ₂ concentration, measured at NOAA Mauna Loa Observatory.	Mean CO ₂ concentration, reported as dry air mole fraction	Annually	1958 to present	NOAA Earth System Research Laboratory, Global Monitoring Division	Monthly	Data shared on NOAA website. Indicator in State of the Delaware Inland Bays report.	Continue status quo.
Climate Characteristics ◆	—	CCMP: Planning for Climate Change, Objective 1 Questions: 26, 28	Average annual air temperature, annual rainfall, growing season length	Weather characteristics (temperature, precipitation, frost dates)	Hourly	Air temperature, precipitation: 1995 to present Growing Season Length: 1946 to Present	Delaware Environmental Observing System (DEOS)	Daily	DEOS website. Indicator in State of the Delaware Inland Bays report.	Few stations in the watershed provide inadequate resolution for microclimate variations.

RECOMMENDATIONS AND SUPPORTING PROGRAMS

The previous section of this plan describes ongoing monitoring programs in the Inland Bays that exist currently. Many of these existing programs provide data that are used to develop the long-term indicators for the *State of the Delaware Inland Bays* reports that are published every five years by the CIB.

This section of the plan provides recommendations for new monitoring programs, or enhancement of existing programs. These recommendations are made based upon critical data gaps (identified in Table 1), the availability of new methods or technologies, and/or changes needed to make programs sustainable over the long term. Table 2 summarizes priority recommendations, and each is described more fully on pages following the table.

Table 2 - Recommendations for Monitoring in the Inland Bays (Categorized according to relative priority (*** = highest priority).

Objectives	Data to be Collected / Contribution to Other Programs	Responsible Entity/Entities	Future Frequency of Reporting/Use	How Future Data Will Be Shared / Reported	Funding and Timeline	Priority Level
1. DEVELOPMENT OF A NEW HYDRODYNAMIC/WATERSHED MODEL FOR THE INLAND BAYS						
Build a well-calibrated hydrodynamic/water quality model for the Inland Bays in order to address water quality threat of dissolved oxygen (DO).	<ol style="list-style-type: none"> 1) Will collect more recent sediment flux measurements, a sediment flux model, and explicit incorporation of benthic algae; 2) Will collect lag times for groundwater and soil phosphorus loads to be incorporated; 3) Will incorporate primary production and respiration rates into model calibration; 4) Will collect increased spatial resolution particularly in tidal headwaters; 5) Will incorporate meteorological records that were unavailable during calibration years (1998-2000); 6) Will incorporate re-assessment of nutrient loading to include potential groundwater discharge being explored by many DIB researchers since 2000. 	<p>Recommended partners: CIB STAC, University of Delaware, DNREC Division of Watershed Stewardship.</p> <p>Appointment of a standing STAC CCMP/Monitoring subcommittee will begin work in early 2018.</p>	X	X	<p>Building a well-calibrated model will take significant data, funding, and time.</p> <p>Will take several years and several million dollars.</p> <p>STAC subcommittee will engage stakeholders in early 2018 to plan subtasks, responsible entities, timeline, and funding strategy. Plan to be presented to STAC by June 2018.</p>	***
2. UPGRADE CITIZEN MONITORING PROGRAM DATABASE AND SERVE DATA TO PUBLIC ONLINE THROUGH STORET AND STATE WATER QUALITY PORTAL						
Develop a CMP data entry template and database processing tools that will allow submission of data to STORET through the EPA's Water Quality Exchange (WOX).	<ol style="list-style-type: none"> 1) Will ensure long-term sustainability of the Inland Bays Citizen Monitoring Program; 2) Will fill critical data needs for effective restoration and management of the estuary. 	The CIB, CMP, EPA Region 3, Center for Environ. Monitoring and Analysis (CEMA).	Data synthesized and reported by CIB every 5 years in State of the Bays reports. Included in biannual 305(b) reports.	Publicly available and searchable through a common online portal, such as the Delaware WQ Portal.	Estimated cost to complete is \$50,000. EPA has provided supplemental funding to the CIB (\$35,000) to begin work in FY2018	***

Table 2 - Recommendations for Monitoring in the Inland Bays (Categorized according to relative priority (*** = highest priority).

Objectives	Data to be Collected / Contribution to Other Programs	Responsible Entity/Entities	Future Frequency of Reporting/Use	How Future Data Will Be Shared / Reported	Funding and Timeline	Priority Level
3. LONG-TERM, CONTINUOUS MONITORING OF DISSOLVED OXYGEN AND CHLOROPHYLL						
Develop and deploy continuous monitoring networks to measure dissolved oxygen and chlorophyll in the Inland Bays, with a focus on tributaries.	1) Will increase sufficiency to detect rapidly changing or cycling conditions in the Bays or episodic events (ex. diel-cycling hypoxia, and phytoplankton blooms).	STAC, Inland Bays, University of Delaware CEOE, DGS.	Daily online reporting.	Data available through online portal.	Cost dependent on methods used and # of stations monitored. Based on similar NJ system, \$50,000/sensor installation cost, plus \$15-20,000/sensor/yr recurring maintenance.	***
4. MONITORING OF SUBMERGED AQUATIC VEGETATION IN THE INLAND BAYS						
Monitor and document acreage and distribution of SAV in the Inland Bays.	1) Will assist protection of any existing SAV beds; 2) Will help focus restoration efforts in areas where SAV beds are known to survive; 3) Will track progress in SAV restoration.	CIB, CERF SAV monitoring workgroup, MD Coastal Bays Program; EPA Region 3 expressed interest in collaborating on SAV monitoring.	Every 1 to 5 years, depending on resource availability.	Available in the State of the Bays report (every 5 years)	Funding impossible to determine until methodologies are established.	***
5. MONITORING OF LOCAL INDICATORS OF SEA LEVEL RISE, INCLUDING A FLOOD MONITORING NETWORK						
Expand the CFMS into the Inland Bays to partly fill the need for more local indicators of climate change and sea level rise	1) Will provide real-time tool to create flood inundation potential maps and time series of forecasted tidal predictions; 2) Will monitor sea-level rise and the development/validation of hydrodynamic models if maintained for an extended period of time.	CEMA, DeIDOT, CIB, DE Geological Survey (DGS).	Continuous.	Available to the public via online application.	Annual cost to maintain current CFMS network of approx. \$15,000; adding more sensors will increase cost. May be able to use existing DeIDOT telemetry system to expand the sensor network. Potential funding from DNREC Coastal Programs, NOAA, DEMA, Sussex County.	***

Table 2 - Recommendations for Monitoring in the Inland Bays (Categorized according to relative priority (*** = highest priority).

Objectives	Data to be Collected / Contribution to Other Programs	Responsible Entity/Entities	Future Frequency of Reporting/Use	How Future Data Will Be Shared / Reported	Funding and Timeline	Priority Level
6. CONTINUE MONITORING TIDAL FLUSHING AT THE INDIAN RIVER INLET						
Continue measurement of tidal prism at the Indian River Inlet.	<ol style="list-style-type: none"> 1) Will evaluate the volume of water passing through the inlet; 2) Will monitor the residence time of water within Inland Bays; 3) Data required for development of hydrodynamic model for the Bays. 	U.S. ACOE, Univ. of Delaware.	Every 5 years	State of the Bays Report (every 5 years).	Estimated cost \$40,000 per survey. U.S. ACOE and DelDOT possible funding partners.	**
7. MONITORING OF OYSTER RECRUITMENT AND GROWTH IN THE BAYS						
Develop a monitoring plan that regularly measures oyster recruitment, populations, and growth in all three bays.	<ol style="list-style-type: none"> 1) Will assist with understanding the dynamics of restored oyster populations and their ecosystem services under naturally variable conditions; 2) Will track progress of CCMP objective to enhance oyster populations in the Bays. 	CIB, Delaware State University, DE Sea Grant, DNREC Divisions of Watershed Stewardship and Fish & Wildlife.	Annually. Future indicator of restoration success.	Annual project reports published by the CIB. Potential future indicator for State of the Bays reports.	Estimated cost is \$30,000 annually. CIB should work with DSU to implement regular survey in 2018.	**
8. SHORELINE CONDITION AND MODIFICATION MONITORING						
Build upon and update shoreline inventories conducted by VIMS in 2006 (for Indian River Bay) and 2012 (for Rehoboth Bay).	<ol style="list-style-type: none"> 1) Update inventories for IR and Rehoboth Bays; 2) Complete inventory for Little Assawoman Bay; 3) Will help track success of Living Shoreline Initiative 	CIB, DNREC Wetlands Assessment and Monitoring Program, DNREC Subaqueous Section, VIMS.	Every 5-10 years.	Data made available through online mapping tool and project reports. Potential future indicator for State of the Bays reports.	\$125,000 every 10 years. Cost could be reduced by use of volunteers and/or drones in data collection.	**

Table 2 - Recommendations for Monitoring in the Inland Bays (Categorized according to relative priority (*** = highest priority).

Objectives	Data to be Collected / Contribution to Other Programs	Responsible Entity/Entities	Future Frequency of Reporting/Use	How Future Data Will Be Shared / Reported	Funding and Timeline	Priority Level
9. CONTINUE ANALYSES OF TIDAL MARSH ACREAGE AND CONDITION USING GIS METHODOLOGY ESTABLISHED IN 2104 RARE STUDY						
Use established methodologies for GIS analyses of aerial imagery and LULC data to monitor Inland Bays tidal marsh acreage/condition.	<ol style="list-style-type: none"> 1) Will help understand and track status and trends in overall extent and health of tidal marshes in the estuary; 2) Will assist with prioritizing future research and areas for restoration; 3) Continue to use tidal marsh acreage and extent of fractured pooling as indicators in State of the Bays reports. 	CIB, Univ. of Delaware Water Resources Agency.	Every five years.	State of the Bays Report (every 5 years).	Total cost per analysis estimated at \$37,000.	*
10. MONITORING OF ESTUARY ACIDIFICATION						
Collect data needed to understand proton fluxes and balances in the Inland Bays.	<ol style="list-style-type: none"> 1) Will allow assessment of long-term trends in acidification that may occur with climate change; 2) Will provide data to help model the interactions between hydrodynamics, eutrophication, and estuary acidification; 3) Will help understand potential impacts on shellfish in the Bays. 	Univ. of Delaware CEOE, DNERR, Mid-Atlantic Coastal Acidification Network	Continuous monitoring, reporting mechanism and frequency to be determined.	Continuous monitoring data could potentially be made available online. Will be shared in project technical reports and through STAC.	Estimated \$30,000 per station added installation cost annually if included as a component of a continuous WQ sensor network.	*
11. MONITORING OF RECREATIONAL BLUE CRAB AND HARD CLAM HARVESTS						
Obtain estimates of recreational crab and clam landings in the Bays.	<ol style="list-style-type: none"> 1) Data will help assess the health and status of shellfish populations and allow state to account for recreational harvest in management of these fisheries. 	DNREC Div. of Fish & Wildlife	Annually.	Annual reports from DFW. Potential future indicator for State of the Bays reports.	\$46,000 annually.	*

Table 2 - Recommendations for Monitoring in the Inland Bays (Categorized according to relative priority (*** = highest priority).

Objectives	Data to be Collected / Contribution to Other Programs	Responsible Entity/Entities	Future Frequency of Reporting/Use	How Future Data Will Be Shared / Reported	Funding and Timeline	Priority Level
12. DEVELOP A LIST OF EMERGING CONTAMINANTS RESEARCH AND MONITORING IN THE INLAND BAYS.						
<p>Build a baseline for determining which compounds have the highest potential for significant impact in the Bays, and prioritize future EC monitoring and research.</p>	<ol style="list-style-type: none"> 1) State to maintain a database of contaminants found in monitoring of water supplies; 2) STAC to build and maintain a list of research and monitoring activities investigating emerging contaminants in the Bays; 3) Will allow definition of the problem, recommendations for further monitoring, and promote source control. 	<p>STAC CCMP/monitoring subcommittee.</p>	<p>Biannually.</p>	<p>STAC white papers. Data to be evaluated and used for biannual updates to the IBEMP.</p>	<p>In-kind agency staff time.</p>	<p>*</p>

1. HYDRODYNAMIC/WATERSHED MODEL

The first state-of-the-art water quality modeling program implemented in Delaware's Inland Bays used a calibration database that included data from DNREC, USGS, US Army Corps of Engineers, University of Delaware researchers, citizen monitors and others (Cercio et al. 1994). The model was calibrated using data from 1988-1990 and is largely based on the same model used to manage water quality in Chesapeake Bay (CE-QUAL-W2) (Cercio and Noel 2005). This model included a mechanistic sediment flux model and even included a benthic algal model due to the shallow nature of the bays (Cercio and Seitzinger 1997).

In 2004, Entrix, Inc. and J.E. Edinger Associates developed a TMDL model, currently used for the Inland Bays (Entrix and JEEAI 2004). The model is a fully coupled 1-D watershed and 3-D hydrodynamic-water quality model called the Generalized Environmental Modeling Surface Water System (GEMSS). That model was primarily used to calculate water quality constituents such as nitrogen, phosphorus (particulate/dissolved, inorganic/organic) and dissolved oxygen. It was calibrated using data collected from 1998-2000.

Since 2000, there has been a significant increase in both understanding and data collection in Delaware's Inland Bays. In particular, the University of Delaware and DNREC have collected continuous data records for dissolved oxygen (DO) over many years and at many locations. DO is arguably the greatest potential water quality threat to Delaware's Inland Bays, with multiple fish kills attributable to hypoxia occurring most years. Substantial research efforts have also demonstrated reduced growth rates and behavioral avoidance of hypoxia by juvenile estuary dependent fishes that rely on the Bays for essential fish habitat.

For this reason, the CIB requested an independent assessment of the DO calculation in the GEMSS model (Brady, 2014). While the report focuses on DO, it notes that improvements in the understanding of nutrient loading and biogeochemical cycling will also be necessary to improve future model formulations. The conclusions of this report are that GEMSS is not effective at simulating DO (especially in Indian River and tributaries). The calibration and validation datasets from 1998-2000 included few to no substantive continuous DO records. Assessing model performance in relation to diel-cycling hypoxia is exceedingly difficult, and that was not the original intent of that modeling effort. DO data collected since 2001 contain DO fluctuations from 0% to 200% saturation in the headwaters of major creeks/tributaries, and the model output shows no such fluctuations. Brady's explanation for this is either: (1) diel-cycling hypoxia only became a significant feature of the water quality in Delaware's Inland Bays in 2001 or (2) the monitoring program only became robust enough to detect diel-cycling hypoxia in 2001. In either case, the proliferation of data and understanding since 2001 strongly argues for re-visiting the modeling framework for the Bays.

Recommendation: *There is a critical need for a predictive, coupled watershed, hydrodynamic, and water quality model for the Inland Bays that uses current and high-frequency data.*

Brady (2014) provided specific recommendations for future accurate simulations of diel-cycling hypoxia. These include:

- In the shallow Inland Bays estuary, benthic pelagic coupling between the water column and sediments is potentially a large source of oxygen demand. The current GEMSS model uses fluxes measured from 1992-1993 and 2001. There is no mechanistic sediment flux model associated with this modeling effort. More recent flux measurements, a sediment flux model, and explicit incorporation of benthic algae will almost certainly be necessary to complete nutrient budgets. There is also potential for the bottom sediment to play a role in time lags between the implementation of the Pollution Control Strategy and response in the estuary that cannot be explored in the current modeling framework. Lag times for groundwater and soil phosphorus loads should also be incorporated.
- Incorporation of primary production and respiration rates into model calibration. Seasonal respiration appears well calibrated, but daily respiration rates are clearly not large enough to generate hypoxia in the early to late morning. These rates can be estimated from high-frequency DO measurements.

- Increased spatial resolution particularly in tidal headwaters where recent fish tagging evidence has highlighted potential fish exposure mechanisms reliant on spatial gradients in DO. Headwaters are where improvements will be seen first, so an improved model will be a tool to test the effectiveness of best management practices.
- Incorporation of multiple meteorological records that were unavailable or offline during the calibration years (1998-2000) made available by the Delaware Environmental Observing System.
- Re-assessment of nutrient loading to include potential groundwater discharge being explored by many DIB researchers since 2000. The operation of the Millsboro Pond Outlet Aqualab provides more complete loading and loading process information, particularly for storms and storm periods. It also identifies significant diel DO swings in the pond. This is the largest point source of fresh water to the Indian River.

Building a well-calibrated model will require significant data, funding, and time. Development of a predictive model tool will need to plan for a minimum of several years for data collection and model building/calibration. The total cost is likely to be several million dollars. Thus it is essential that the effort be broken down into subtasks. The model development should be led by the CIB's STAC and include the following immediate actions:

- a. Appointment of a standing STAC CCMP/Monitoring subcommittee that will begin work in early 2018. This subcommittee will have among its responsibilities overseeing the implementation of the IBEMP recommendations, including model development as one of the highest priorities. The chair of this subcommittee will work closely with the STAC presiding officers and CIB staff to manage development of a new model.
- b. The subcommittee will engage stakeholders in early 2018 in order to put together a specific plan for model development consisting of: subtasks, responsible entities, timeline, and funding strategy. The subcommittee will present this plan to the full STAC by June 2018.

2. UPGRADE OF CITIZEN MONITORING PROGRAM DATABASE

Due to funding and staffing issues, the University of Delaware Citizen Monitoring Program (CMP) has struggled to keep up with nutrient analyses and management of data. The CIB recently has provided EPA Section 320 funds to assist the CMP with analysis of backlogged nutrient samples taken at nine stations that are key for producing *State of the Delaware Inland Bays* and *Your Creek* indicator reports. But a larger issue is the fact that the current Access database used by the CMP is unsustainable and incapable of exporting data that can be integrated with the state's water quality databases.

The CIB, the CMP, EPA Region 3, and DNREC have a shared goal converting the current CMP database to a new, supported, sustainable format that allows public, online access and queries of Inland Bays water quality data. Conversion would be made to a database that can export data into the EPA's STORET data warehouse, and allow sharing of data online via the Delaware Water Quality Portal (<http://demac.udel.edu/waterquality/>). The new database could include an online site for volunteer monitors to enter their measurements directly, with data validation, including access from mobile platforms.

Recommendation: It is recommended that the Delaware Environmental Monitoring & Analysis Center (DEMAC, University of Delaware), which manages the state's Water Quality Portal, work with the CMP and the CIB to develop a CMP data entry template and database processing tools that will allow submission of data to STORET through the EPA's Water Quality Exchange (WQX).

Specific recommended objectives and actions are:

1. *Development of an updated, supportable database structure for the CMP.*
 - a. Creation of a mechanism and framework whereby CMP data can be submitted.
 - b. Development of data ingestion software to input CMP data into a database or repository.
 - c. Creation of a QA/QC reviewer access procedure for quality control purposes.

2. *Online availability of CMP data to the public.*
 - a. Creation of a mechanism for automated submission of CMP data into STORET from the newly developed CMP database.
 - b. Ingestion of CMP data from STORET into the Delaware Water Quality Portal as an additional "station type" for easy public access to CMP data.

Once this is complete, legacy data should be converted, if possible, for inclusion into STORET.

Successful completion of this project will help ensure long-term sustainability of the Inland Bays Citizen Monitoring Program and fill critical data needs for effective restoration and management of the estuary. A robust, high-quality water quality data set for the bays, combining both volunteer and state-collected data, will be publicly available and searchable through a common online portal. Additionally, this effort will further a goal of both the CIB and EPA to have CMP data incorporated into water quality analyses and models managed by the state.

Funding: Estimated cost to complete Objectives 1 and 2 is \$40-50,000. EPA has provided supplemental funding to the CIB in the amount of \$35,000 to begin this work in FY 2018, and it is included as a project in the CIB's FY 2018 workplan.

3. CONTINUOUS WATER QUALITY MONITORING

Ambient water quality monitoring in the Bays has to date largely followed a traditional paradigm of discrete sampling to collect data from as many locations and as often as funding allows. Typical best-case sampling frequencies are weekly (for the Citizen Monitoring Program) or quarterly (for the State's GAMN network). While this type of monitoring program may be useful for long-term status and trend analyses, it does not provide sufficient resolution to detect rapidly changing or cycling conditions in the Bays or episodic events. Examples of these in the Inland Bays system include diel-cycling hypoxia (Tyler et al., 2009) and phytoplankton blooms.

Continuous monitoring is increasingly becoming a standard to characterize water quality in shallow coastal systems, where conditions can change frequently over time. Continuous monitoring is the sampling method of choice when water quality variations are to be characterized over time. Multiparameter sondes, for example, are increasingly being used to monitor water quality at fixed monitoring sites, to carry out vertical profiling, or to perform water quality mapping. Flow-through continuous monitoring stations also are developing and have been used at a few sites in the state. DNREC has deployed some continuous sensors at Millsboro Pond in the Bays watershed. New, autonomous platforms are available provide spatial resolution for parameters that cannot be sensed remotely.

Continuous monitoring in coastal environments can be challenging because of rapid biofouling from microscopic and macroscopic organisms, corrosion of electronic components from salt and high humidity, and wide ranges in values of field parameters associated with changing weather and tidal conditions. The sensors that are used to measure water-quality field parameters require careful field observation, cleaning, and calibration procedures, as well as thorough procedures for the computation and publication of final records. However, procedures and technologies for continuous water quality monitoring have evolved greatly in recent years, and continue to evolve. Emerging sensor technology broadens the variety of measurable chemical constituents and reduces the limits of detection. Because it has become possible to make near real-time water-quality monitoring data available on the Internet, continual progress is being made to improve applications and refine quality-control procedures.

Recommendation:

Continuous monitoring networks to measure dissolved oxygen and chlorophyll should be developed and deployed in the Inland Bays, with a focus on tributaries. This effort should build upon ongoing work in this area by Dr. Bill Ullman and others, and consider emerging, innovative technologies. The STAC should help guide the selection of sites and station configuration.

Funding:

Dependent upon the methods used and number of stations monitored. Annual cost to maintain the continuous monitoring station currently deployed at Massey's Ditch (USGS 01484680, measuring temperature, specific conductivity, DO, and pH) is on the order of \$50,000. A real-time estuarine water quality monitoring network with 7-8 stations was implemented in New Jersey by the Barnegat Bay Partnership and the NJDEP. The network includes both shore-mounted and buoy-mounted stations. Cited costs are approximately \$50,000 for installation of each new shore-based station, and \$80,000-100,000 in recurring maintenance and operation costs for all sondes per year.

4. MONITORING OF SUBMERGED AQUATIC VEGETATION

While listed as a major action in CCMP, submerged aquatic vegetation (SAV) restoration efforts in the Inland Bays have stalled. Due to high levels of nutrients, chlorophyll a, and macroalgae, SAV beds in the Inland Bays have all but disappeared. However, recent water quality analyses suggest that water quality has improved in many locations over the long term, and that macroalgae populations are much lower than previous levels in much of the Inland Bays. This suggests that SAV beds may be able to survive in the Inland Bays once again, and may currently exist in some areas that have seen improvements.

Recent surveys of SAV in the Bays have been limited to the Horned Pondweed meadows discovered a few years ago in Love Creek. No monitoring program currently exists for eelgrass or other baygrasses.

Recommendation:

Monitoring to document what SAV remains in the Inland Bays, and where it's located are necessary in order to protect any existing SAV beds, as well as focus restoration efforts in areas where SAV beds are known to survive. In addition, the recognition of SAV as an important carbon sink adds impetus to the need for accurate maps at a regional level.

Ideally the survey will be performed annually, but adjusted depending on resource availability. At a minimum it should be conducted every five years to correspond with indicator development for the *State of the Delaware Inland Bays* reports.

Recent technological advances present new options for monitoring and mapping SAV. Aerial surveys are one option. However, the current sparseness of SAV beds in the Inland Bays means that surveyors would be searching for much smaller patches than are typically identifiable by plane. Monitoring may also be conducted either with boats and divers, or with short-range, low elevation drone flights, checking areas for SAV presence and patch size. Divers and drone pilots could maximize their time by focusing efforts in areas where water quality meets SAV requirements as determined through the State of the Bays report and other water quality analyses. A workshop at the November 2017 conference of the Coastal and Estuarine Research Federation (CERF) will be devoted to sharing best practices for SAV monitoring, including new technology options.

Funding:

Funding required is impossible to determine until methodologies are established. EPA Region 3 has expressed interest in potentially collaborating on SAV monitoring.

5. LOCAL INDICATORS OF SEA LEVEL RISE

There is a critical need in the Inland Bays watershed to conduct monitoring geared specifically to address how sea level rise affects and is perceived by the public. The Inland Bays watershed is particularly vulnerable to sea level rise and its effect on the frequency and intensity of coastal flooding events, emphasizing the need for a modern, dependable coastal flood monitoring and warning system for the Bays' coastal communities.

The Delaware Coastal Flood Monitoring System (CFMS) was developed, jointly by the Delaware Geological Survey (DGS) and the Delaware Environmental Observing System (DEOS) at the University of Delaware, to provide water level predictions and flood potential for 15 communities on Delaware Bay

(<http://coastal-flood.udel.edu/>). The tool serves three primary functions: to send out warning alerts up to 48 hrs in advance of potential flood conditions, to provide access to current meteorological and hydrologic conditions, and to provide local tidal predictions and map their areas of impact.

The CFMS currently covers only the Delaware Bay coastline. Expansion of the system to the Inland Bays and Atlantic Coast of Delaware is planned by DEOS. However, the NOAA DBOFS operation model used currently does not work well for the Inland Bays system. A different hydrodynamic or statistical model is required.

In 2015, DEOS partnered with DGS and the CIB to conduct a three-year study of water level conditions at various locations in the Inland Bays. Water level sensors were installed in spring of 2015 and likely will be maintained until the summer of 2018. Data collected will help inform development of a flood prediction model specific to these bays.

Recommendation:

Expansion of the CFMS into the Inland Bays will at least partly fill the need for more local indicators of climate change and sea level rise. If a successful model is developed for the Bays, it will provide a publically-accessible, real-time tool to create flood inundation potential maps and time series of forecasted tidal predications. The CFMS itself is not meant to be a sea level rise tool, but rather its continued development is contingent upon the availability of tidal water level data from a sensing network like the one currently deployed in the Inland Bays. The data from that same network can lend itself towards sea level rise monitoring and the development/validation of hydrodynamic models if maintained for an extended period of time.

It is recommended that support for developing this tool for the Inland Bays be continued and prioritized. A water level/flood monitoring network should be permanently installed throughout the Bays. These data are needed not only for flood alert tools, but also for development of a new hydrodynamic model for the Inland Bays.

Priority should also be given to long-term local monitoring of other indicators of climate change, such as precipitation, air temperature, and growing season length. The Inland Bays watershed experiences a range of microclimate effects, so data collected at the coast, for example, (or from outside the watershed) cannot easily be extrapolated to inland locations.

Funding:

The annual cost to maintain the current network of CFMS sensors is approximately \$15,000, including upkeep of a borrowed (DelDOT-owned) RTK-GPS system. Installation of new sensors would add more cost, so maintaining what is there already is more cost effective, as long as the locations are answering the pertinent science questions. DelDOT's telemetry system might be used to expand the network.

Currently the CFMS is funded through grants from the DNREC Delaware Coastal Management Program and the Delaware National Estuarine Research Reserve, with funding from NOAA's Office of Ocean and Coastal Resource Management. Opportunities for additional funding from NOAA, DEMA, and Sussex County should be explored.

6. INDIAN RIVER INLET TIDAL FLUSHING

Since it was stabilized in the 1930s, the Indian River Inlet has deepened over time, passing greater volumes of water and increasing the tidal range of the Bays. This has led to long-term increase in salinity of the Bays and contributed to degradation of marshes. Increased tidal flushing through the Inlet also results in greater flushing of nutrients from the system.

Inlet flushing is one of the important indicators included in the *State of the Delaware Inland Bays* reports. Flushing is estimated through tidal prism calculations. The tidal prism is defined as the volume of water leaving a channel on an ebb tide cycle. The following general equation is used:

$$P = H A$$

where H is the average tidal range and A is the average surface area of the basin. Area is calculated through a series of transects and using bathymetry data for depth.

Tidal prism calculations for the Indian River Inlet are not made with regularity. Calculations were made in 1939, 1952, 1969, 1975, 1976, 1978, 1979, 1981, 1985, 1986, 1988, and 2004. Bathymetry data was collected in 1999 and 2004. The most recent measurements were performed by Ocean Surveys, Inc. (Saybrook, CT).

Tidal prism calculations can be used not only to evaluate the volume of water passing through the inlet, but also the residence time of water within the Inland Bays. Evaluation of the Indian River Inlet flushing indicator during preparation of the *2016 State of the Delaware Inland Bays* report revealed a need for dedicated funding to regularly assess the inlet flushing. As the inlet deepens and widens the volume of saltwater will increase and leads to a cascade of ecological impacts. In addition, the change in channel morphology may lead to structural problems with the inlet bridge. 2004 data were provided by the Army Corps of Engineers, Coastal Planning Section.

Recommendation: It is recommended that the state work with the Army Corps to repeat these measurements every five years. Because of its interest in resiliency of coastal roadways and long-term integrity of the inlet and its bridge, DelDOT may be able to contribute to this effort.

Funding: The tidal prism data collected in 2004 was one portion of a larger data collection effort funded under a single contract. The Army Corps estimates the tidal prism portion of the work cost roughly \$30,000. The tidal prism data collection effort involved hourly boat-mounted ADCP surveys measured over a 25-hour period along five transects, as shown in Figures 3 and 4. Today, a similar effort would likely cost closer to \$40,000.




FIGURE 3—Hydraulic measurements made at the Indian River Inlet in 2004, showing current measurement transects.




FIGURE 4 – Hydraulic measurements made at the Indian River Inlet in 2004.

7. MONITORING OF OYSTER RECRUITMENT AND GROWTH IN THE BAYS

A major goal of the Center is to restore a sustainable population of native oysters in the Inland Bays. Oyster restoration/enhancement projects (such as living shorelines, oyster reefs, and the CIB’s oyster gardening program) and commencement of oyster aquaculture in the Bays will all contribute to this goal. Currently, however, no regular monitoring of oyster population, distribution, or recruitment in the Bays is occurring. Such monitoring is necessary to evaluate the effectiveness of restoration efforts.

A critical need also exists for studies on post-settlement growth and survival of oysters in the Inland Bays system. Existing literature documenting growth rates for Eastern Oysters is largely from studies of large, protected, or hatchery-spawned animals. Multi-year, population-level estimates of wild growth in local bay waters are necessary in order to understand the dynamics of restored oyster populations, and their ecosystem services, under naturally variable conditions. Such studies have very recently begun in Delaware Bay (Munroe et al., 2017).

Recommendation:

The CIB currently is drafting a Shellfish Enhancement Action Plan that will include recommendations for specific restoration projects such as oyster reefs. All implementation projects should, if feasible, be monitored for at least three to five years for oyster survival, growth, reef height and size (if applicable), disease, and recruitment.

In addition, however, it is recommended that regular, long-term surveys of oyster populations and recruitment be developed for all three Bays. A plan for this monitoring should be a component of the Shellfish Enhancement Action Plan.

A current research project being conducted under the guidance of Dr. Gulnihal Ozbay at Delaware State University may provide a template for this monitoring program. The focus is to develop repeatable methods that can be used later to assess the impact that aquaculture and hatchery raised oysters have on

the local wild oyster population. Components of this effort include: (a) Standardized transect surveys on riprap–armored shorelines; (b) spat collectors deployed at locations throughout the Bays; and (3) genetic analyses of spat, to determine the diversity and probable parent populations.

Funding:

DSU has estimated the cost of a long-term monitoring program to be approximately \$30,000 annually. CIB and DSU should work together to identify funding to implement a survey in 2018.

8. MONITORING OF SHORELINE CONDITION AND MODIFICATION

The Center is engaged in an initiative to maximize the use of living shorelines stabilization techniques in order to protect the water quality and habitat of the estuary. The initiative is a focus of the Comprehensive Conservation and Management Plan (CCMP) for the Inland Bays and is also a priority for the State.

A study to assess the shoreline conditions of Rehoboth and Indian River Bays was conducted by the Virginia Institute of Marine Science (VIMS), Center for Coastal Resources Management, with support from DNREC's Wetlands Monitoring and Assessment Program and the Center for the Inland Bays. Data for Indian River Bay were collected in 2006; data for Rehoboth Bay were collected in 2012. The spatial data collected in the study were used to build a public web-based mapping and analysis interface (http://cmap.vims.edu/ShlInv/Delaware/Delaware_shlinv.html). The assessment was not completed for Little Assawoman Bay at the time, due to funding constraints.

The data developed for the inventory were based on a three-tiered shoreline assessment approach. This assessment characterized conditions that can be observed from a small boat navigating along the shoreline. Hand-held GPS units and GPS registered videography were used to collect data on conditions observed in the field. The three tiered shoreline assessment approach divided the shore zone into three regions: 1) the immediate riparian zone, evaluated for land use; 2) the bank, evaluated for height, stability, cover and natural protection; and 3) the shoreline, describing the presence of shoreline structures for shore protection and recreational use.

A complete and up to date shoreline condition database is an important tool for evaluating success in meeting the goal of reducing hardened shorelines and increasing the extent of natural/living shorelines in the Bays. If maintained long-term, the data can be used to develop useful shoreline condition indicators for the Inland Bays. The online analysis tool can assist land owners, marine contractors, and State regulators to identify locations where shoreline restoration is needed and what type of shoreline restoration method would be most appropriate. In particular, the data may be used to help CIB and others prioritize sites for installation of living shoreline restorations in all three Inland Bays.

Recommendation:

It is recommended that a shoreline inventory be completed for Little Assawoman Bay, and that the shoreline data collected in 2006 for Indian River Bay be updated within the next one to two years. In order to use the shoreline data to develop an environmental indicator for the Bays, the data will need to be updated periodically. Ideally this would happen every five years, to correspond with release of the *State of the Delaware Inland Bays* reports.

Funding:

Total cost for the original inventory of Rehoboth and Indian River Bays was ~\$125,000. In 2015, VIMS provided a cost estimate of ~\$67,000 to complete the inventory for Little Assawoman Bay and update Indian River Bay data.

The Center recommends that this work be completed more cost effectively by using in-house staff and volunteers to collect shoreline photographs and field data from boats. Use of drones may also facilitate data collection. Center staff, or a contractor, would then update the mapping layers and report the results. The Barnegat Bay Partnership recently developed a citizen science program called "Paddle for the Edge," to monitor shoreline condition. Data are collected from kayaks and canoes using a

smartphone app. A similar program could easily be developed for the Inland Bays.

9. CONTINUATION OF TIDAL MARSH ACREAGE/CONDITION AS ENVIRONMENTAL INDICATORS

Monitoring the extent and condition of the tidal marshes of the Inland Bays is critical for determining overall health of the estuary and to track trends in its condition. By understanding changes over time it is possible to identify potential areas of concern and prioritize future research and remediation efforts.

The *2016 State of the Delaware Inland Bays* report included new salt marsh acreage and condition (fractured pooling) indicators. These indicators were developed using data generated by a Regional Applied Research Effort (RARE) study conducted by the University of Delaware and the Center for the Inland Bays (Jo et al, 2014). The study documented the areal extent of the marshes of the Inland Bays at a number of intervals between 1937 and 2007 using geospatial analyses of aerial photography, State of Delaware wetland maps, and Landsat Thematic Mapper satellite imagery. Historic trends in the extent of vegetated marsh, fractured pooling, ditching, and wetland/upland boundary hardening provide an indication of the general health of the tide marsh system. The RARE study established a methodology for continued analysis of status and trends.

Recommendation:

In order to continue to use tidal marsh acreage and extent of fractured pooling as environmental indicators for the Bays, analyses of updated aerial imagery and land cover data should be repeated at least every five years, using the established GIS methodology. The University of Delaware Water Resources Agency (WRA) is best suited, at this time, to conduct these analyses.

Funding:

To perform the analysis, including data compilation and generation, aerial photointerpretation, GIS processing of files, generation of data for trend analysis, and reporting of status and trend results, the WRA estimates a total cost of \$21,100 to \$37,700.

10. ESTUARY ACIDIFICATION

The drivers, patterns, and ecological impacts of acidification in estuaries are not well understood, due to dramatic spatial and temporal variation in the processes that control pH in near-shore environments. Upwelling/overturn, tides, eutrophication, and watershed alteration are expected to interact with increasing atmospheric carbon dioxide and warming waters in complex ways. Proton fluxes may vary seasonally or with weather patterns. Biological impacts may also vary. Yet there is evidence that even a slight increase in acidification disrupts recruitment and growth of shellfish. Responses of other organisms to acidification are less clear.

Currently no long-term acidification monitoring is being conducted in the Inland Bays. Whether or not it is a significant concern is unknown. Monitoring pH in estuaries is not straightforward. Other National Estuary Programs have deployed cutting-edge systems to monitor long-term coastal acidification trends, including San Francisco Bay, Santa Monica Bay, Tampa Bay, Massachusetts Bay, Casco Bay, Barnegat bay, Long Island Sound, and Corpus Christi Bay. In addition, EPA has measured acidification in the Delaware River and Estuary as well as the Mid-Atlantic waters. Dr. Bill Ullman's group at the University of Delaware currently is studying acidification in the Murderkill Estuary, using a continuous, stable, high frequency pH sensor for estuarine and marine applications.

The Mid-Atlantic Coastal Acidification Network (MACAN) effort may help inform this monitoring (<http://midacan.org>). MACAN intends to develop research priorities and a coastal acidification monitoring framework for the Mid-Atlantic region in the near future.

Recommendation:

Collection of data needed to understand proton fluxes and balances in the Inland Bays should be included as a component of a continuous water quality monitoring system. Efforts should be made to upgrade planned continuous monitoring stations with sensors to monitor pH and carbonate system parameters. State of the art sensors suitable for use in brackish or marine waters will be required.

Funding:

Barnegat Bay Partnership estimates an added cost of \$30,000 per station to install sensors for carbonate parameters at their network stations.

11. RECREATIONAL BLUE CRAB AND CLAM HARVESTS

DNREC's Division of Fish and Wildlife indicates that obtaining estimates of recreational Blue Crab and Hard Clam harvests is one of the highest priorities for fisheries research in the Inland Bays. Currently no information exists on what quantity of shellfish are being recreationally landed from the Bays. This information would be used to assess the health and status of those shellfish populations and would allow the state to better account for recreational harvest in their management.

Recommendation:

The Division of Fish and Wildlife should develop an ongoing recreational Hard Clam and Blue Crab harvest survey in the Inland Bays similar to the MRIP survey currently used by NOAA recreational fish catch. This survey would rely on a field, 'harvester-intercept' interview survey that records catch rates for species; and a telephone (or mail survey) that is designed to estimate effort. Average catch rates would be applied to the effort estimates to generate landings per year. These surveys would be conducted annually.

Funding:

In 2009, the state contracted with MARCO to conduct this type of survey to generate a recreational landings estimate for Blue Crabs in the state. The cost was \$46,000. Cost for the proposed survey is expected to be similar.

12. EMERGING CONTAMINANTS

Wastewater Management Objective #2 of the 2012 Addendum to the CCMP is "Examine emerging contaminants entering the Inland Bays and engage the regulatory community and general public in education and source reduction." Action 1A is a symposium be held that "identifies emerging contaminants, their sources, and potential effects." A statewide symposium on emerging contaminants was co-sponsored by the CIB and the University of Delaware on March 13, 2017. Proceedings are available at www.inlandbays.org/events/emerged-symposium/. Contaminants of concern discussed included pharmaceuticals, personal care products, perfluoroalkyl and polyfluoroalkyl Substances (PFAS), polybrominated diphenyl ethers (PBDE), phytoestrogens, oxybenzone (sunscreen), microplastics, and protons (coastal and estuarine acidification). Monitoring, regulation, and research needs were also discussed. Limited monitoring of these compounds has occurred in the Inland Bays study area.

Recommendation:

DNREC's Division of Water and the Delaware Health and Social Service's Division of Public Health report contaminants found during monitoring of water supplies to the U.S. EPA, although the EPA does not have the resources to make regulatory determination on all of them. The State of Delaware should continue to maintain a database of contaminants found and inform EPA of the findings.

The STAC's CCMP/Monitoring subcommittee should also maintain a list of research and monitoring activities investigating emerging contaminants in the Inland Bays. This will build a baseline for determining which compounds have the highest potential for significant impact in the Bays, at which point reports can be prepared to define the problem, recommend further monitoring and research efforts, and promote source control.

Funding: In-kind agency staff time.

REFERENCES

- Appledorn, M.V., and M.E. Baker. 2011. Report to the Delaware Center for the Inland Bays: Methodological Description and Medium-Resolution Results of Spatial and Temporal Changes Riparian Buffers and Implications for Stream Nitrate Concentrations. University of Maryland, Baltimore County, February 2, 2011.
- Baker, M.E., D.E. Weller, and T.E. Jordan. 2006. Comparison of Automated Watershed Delineations: Effects on Land Cover Areas, Percentages, and Relationships to Nutrient Discharge. *Photogrammetric Engineering and Remote Sensing* 72(2):159-168.
- Batiuk, R., P. Bergstrom, M. Kemp, E. Koch, L. Murray, C. Stevenson, R. Bartleson, V. Carter, N. Rybicki, J. Landwehr, C. Gallegos, L. Karrh, M. Naylor, D. Wilcox, K. Moore, S. Ailstock, and M. Teichberg. 2000. Chesapeake Bay submerged aquatic vegetation water quality and habitat-based requirements and restoration targets: A second technical synthesis. CBP/TRS 245/00. EPA/903/R-00/014. Annapolis, MD: U.S. Environmental Protection Agency, Chesapeake Bay Program.
- Brady, D.C. 2014. TMDL Model and Data Evaluation for Delaware's Inland Bays: Modeling Diel-cycling Hypoxia in Delaware's Inland Bays.
- CB&I Environmental & Infrastructure, Inc. 2014. Long Term Stewardship Plan Burton Island Historical Ash Disposal Area Indian River Generating Station Dagsboro, Delaware, DNREC Site Number DE-1399. CB&I Project No. 1009684052, Prepared for Indian River Power LLC, October 2014, 34 pp.
- Cerco, C. F., B. Bunch, M. A. Cialone, and H. Wang. 1994. Hydrodynamic and Eutrophication Model Study of Indian River and Rehoboth Bay, Delaware. U.S. Army Corps of Engineers, Vicksburg, MS.
- Cerco, C. F. and M. R. Noel. 2005. Incremental improvements in Chesapeake Bay Environmental Model Package. *Journal of Environmental Engineering-ASCE* 131:745-754.
- Cerco, C. F. and S. P. Seitzinger. 1997. Measured and modeled effects of benthic algae on eutrophication in Indian River Rehoboth Bay, Delaware. *Estuaries* 20:231-248.
- Cormier, S., S. Norton, G. Suter, and D. Reed-Judkins. Stressor Identification Guidance Document. U.S. Environmental Protection Agency, Washington, DC, EPA/822/B-00/025, 2000.
- Coxe, R. 2011. Historical Analysis and Map of Vegetation Communities, Land Covers, and Habitats of Thompson Island Nature Preserve, Sussex County, Delaware: Lewes and Rehoboth Canal and Rehoboth Bay Watersheds. Delaware Natural Heritage and Endangered Species Program, November 17, 2011, 93 pp.
- Coxe, R. 2012a. Historical Analysis and Map of Vegetation Communities, Land Covers, and Habitats of Assawoman Wildlife Area, Sussex County, Delaware: Little Assawoman Bay and Indian River Bay Watersheds. Delaware Natural Heritage and Endangered Species Program, July 12, 2012, 289 pp.
- Coxe, R. 2012b. Historical Analysis and Map of Vegetation Communities, Land Covers, and Habitats of Delaware Seashore State Park, Sussex County, Delaware: Indian River Bay, North Atlantic Strand and Rehoboth Bay Watersheds. Delaware Natural Heritage and Endangered Species Program, July 23, 2012, 261 pp.
- Coxe, R. 2012c. Historical Analysis and Map of Vegetation Communities, Land Covers, and Habitats of Fenwick Island State Park, Sussex County, Delaware: Assawoman Bay, Little Assawoman Bay, and Southern Atlantic Strand Watersheds. Delaware Natural Heritage and Endangered Species Program, July 24, 2012, 182 pp.

Coxe, R. 2012d. Historical Analysis and Map of Vegetation Communities, Land Covers, and Habitats of Cape Henlopen State Park, Sussex County, Delaware: Breakwater Harbor, Broadkill River, Lewes-Rehoboth Canal, North Atlantic Strand, and Rehoboth Bay Watersheds. Delaware Natural Heritage and Endangered Species Program, October 8, 2012, 289 pp.

Coxe, R. 2013. Historical Analysis and Map of Vegetation Communities, Land Covers, and Habitats of the James Farm Ecological Preserve, Sussex County, Delaware, 1954-2012: Indian River Bay. Delaware Natural Heritage and Endangered Species Program, October 28, 2013 (updated September 11, 2014), 158 pp.

Delaware Center for the Inland Bays. 2004. Delaware Inland Bays' environmental indicators. Rehoboth Beach, DE.

Delaware Department of Natural Resources and Environmental Control. 2008. Delaware Wetland Conservation Strategy.

Delaware Center for the Inland Bays. 2011. State of the Delaware Inland Bays. Rehoboth Beach, DE. 63 pp.

Delaware Center for the Inland Bays. 2012. Addendum to the Delaware Inland Bays Comprehensive Conservation and Management Plan. 58 pp. <https://www.inlandbays.org/wp-content/documents/Final%209-27-12%20CIB%20CCMP.pdf>

Delaware Center for the Inland Bays. 2016. State of the Delaware Inland Bays. Rehoboth Beach, DE. 70 pp.

Delaware Department of Natural Resources and Environmental Control's Whole Basin Management Team. 2001. Inland Bays/Atlantic Ocean Basin assessment report. Doc. 40- 01/01/01/02. Dover, DE.

Delaware Department of Natural Resources and Environmental Control. 2008. Inland Bays pollution control strategy. Dover, DE.

Delaware Department of Natural Resources and Environmental Control. 2011. Delaware Wetland Monitoring Strategy. Dover, DE, March 2011.

Delaware Department of Natural Resources and Environmental Control. 2015. State of Delaware Draft 2016 Assessment, Listing and Reporting Methodologies Pursuant to Sections 303(d) and 305(b) of the Clean Water Act.

Delaware Sea Level Rise Advisory Committee. 2011. Preparing for Tomorrow's High Tide. A progress report of the Delaware Sea Level Rise Advisory Committee to the Delaware Coastal Programs Office, Delaware Department of Natural Resources and Environmental Control, Dover, DE.

Delaware State Game and Fish Commission, State Park Commission, State Planning Office, Marine Laboratories of the University of Delaware, Water and Air Resources Commission, and Delaware Geological Survey. 1969. Environmental study of the Rehoboth, Indian River and Assawoman Bays. Presented to Governor Russell W. Peterson. Department of Natural Resources and Environmental Control, Dover, DE.

Entrix Inc., J.E. Edinger Associates Inc., and Gahagan and Bryant Associates Inc. 2004. Hydrodynamic and water quality modeling and feasibility analysis of Indian River, Rehoboth Bay and Little Assawoman Bay: Delaware's Inland Bays Flushing Study Project.

FDA. 1998. Bacteriological Analytical Manual, 8th Edition, Revision A. Online version <https://www.fda.gov/Food/FoodScienceResearch/LaboratoryMethods/ucm2006949.htm>.

- Gibson, J. R., M. T. Barbour, J. B. Stribling, J. Gerritsen, and J. R. Karr. 1996. Biological criteria—technical guidance for streams and rivers, revised edition. EPA 822-B-96-001, Office of Water, U.S. Environmental Protection Agency, Washington, D.C., USA.
- Greene, R. 2013. Assessment of Biota and Sediment Data Collected by the Center for the Inland Bays in Conjunction with the Burton Island Bioaccumulation Study. Delaware Department of Natural Resources and Environmental Control, Dover, DE, May 10, 2013, 26 pp.
- Hess, G. K., R. L. West, M. V. Barnhill III, and L. M. Fleming. The Birds of Delaware. University of Pittsburgh Press, Pittsburgh, Pa. 635pp
- Jacobs, A., A. Rogerson, D. Fillis, and C. Bason. 2009. Wetland condition of the Inland Bays watershed. Volume 1. Delaware Department of Natural Resources and Environmental Control, Watershed Assessment Section, Dover, Delaware, USA.
- Jacobs, A.D. 2010. Delaware Rapid Assessment Procedure Version 6.0. Delaware Department of Natural Resources and Environmental Control, Dover, DE. 36pp.
- Jo, Y. H., K. R. Philipp, R. T. Field, A. R. Homsey, C. Bason, and K. Somers. 2014. Acreage and condition trends for marshes of Delaware's Inland Bays as an environmental indicator and management tool: A report to the Center for the Inland Bays to identify and document recent trends in the salt marshes of Rehoboth, Indian River and Little Assawoman Bays.
- Moulton S. R., J.G. Kennen, R. M. Goldstein, J. A. Hambrook. 2002. Revised Protocols for sampling algal, invertebrate, and fish communities as part of the National Water Quality Assessment Program. United States Geological Survey, Open-File Report 02-150. Reston Virginia Office
- Munroe et al., 2017. Early post-settlement growth in Wild eastern oyster (*Crassostrea virginica* Gemlin 1791) populations. *Estuaries and Coasts* 40:880-888.
- Price, K.S., L.M. Valdes, B. Glazer, R. Tyler, S. Runyan, and M. Timmons. 2001. An examination of the use of the biological communities as indicators of nutrient enrichment in Middle Atlantic Coastal Estuaries-Delaware Component Final Report 1998-1999. Submitted to the Delaware Center for the Inland Bays. University of Delaware, Lewes, DE.
- Rattner, B., T.G. Bean, R.S. Lazarus, P.C. McGowan, D.D. Day, R.W. Scarborough, and K. Fleming. 2017. Contaminant Exposure and Productivity of Osprey (*Pandion haliaetus*) Nesting in Delaware Bay and River in 2015. Proceedings Delaware Estuary Science and Environmental Summit 2017, Cape May, NJ, January 22-25, 2017, p.86.
- Riedel, G., and B. Wilson. 2013. Final Report: Assess the Potential for Accumulation of Toxic Trace Elements in Biota near Burton Island Ash Disposal Site Indian River Bay, Delaware. Delaware Center for the Inland Bays, Rehoboth Beach, Delaware, 78 pp.
- Rogerson, A., A. Howard, and A. Jacobs. 2009. Wetlands condition of the Inland Bays watershed. Volume 2. Delaware Department of Natural Resources and Environmental Control, Watershed Assessment Section, Dover, Delaware USA.
- Scarborough, R.W., and M.G. Mensinger. 2005. Evaluation of the Storm Water Sediment Control Forebay at Anchorage Canal, South Bethany, DE. Delaware Department of Natural Resources and Environmental Control, report submitted to the Delaware Department of Transportation, August 2005.
- Shaw, 2008. Facility Evaluation Report, Indian River Generating Station, Burton Island Old Ash Landfill, Millsboro, Delaware, Site Number DE-1399. March 2008.
- State of Delaware, 2014. Surface Water Quality Standards: Title 7 Natural Resources and Environmental Control, Delaware Administrative Code - 7401

Tyler, R.M., D.C. Brady, and T.E. Targett. 2009. Temporal and spatial dynamics of diel-cycling hypoxia in estuarine tributaries. *Estuaries and Coasts* 32(1):123-145.

Ullman, W.J., J.R. Scudlark, and J.A. Volk. 2010. Standard Operating Procedure for the Calculation of N and P Deposition from the Atmosphere to Waters of Delaware's Inland Bays. Report to the Center for the Inland Bays, July 2010. University of Delaware School of Marine Science Policy and the Delaware Department of Natural Resources and Environmental Control, Lewes, DE. 11 pp

U.S. Environmental Protection Agency. 1997. *Field and Laboratory Methods for Macroinvertebrate and Habitat Assessment of Low Gradient, Nontidal Streams*. Mid-Atlantic Coastal Streams (MACS) Workgroup, Environmental Services Division, Region 3, WV. 23 pp

Valdes-Murtha, L. 1997. Analysis of critical habitat requirements for restoration and growth of submerged vascular plants in the Delaware and Maryland Coastal Bays. M.S. Thesis, College of Marine Studies, University of Delaware.

Weston. 1995. Appendix F. Characterization report. A study of the status and trends in the Inland Bays. Delaware Inland Bays Comprehensive Conservation and Management Plan. Delaware Inland Bays program.

APPENDIX A
2012 CCMP ADDENDUM ACTIONS AND PERFORMANCE MEASURES

2012 CCMP ADDENDUM ACTIONS AND PERFORMANCE MEASURES

Focus Area: Nutrient Management	
Objective 1. Monitor the effectiveness of the nutrient management program and CAFO regulations, and suggest and implement revisions as	
Actions	Performance Measures
A. Annually report on watershed agricultural BMP implementation including PCS goals for cover crop acreage, manure storage sheds, and manure relocated or put into alternative use.	Publish annual reports on agricultural BMP implementation.
B. Target and prioritize BMP implementation to areas of the watershed where they will be most efficient and effective.	
B1. Use GIS and BMP performance data to determine the locations of BMPs in the watershed by BMP type resulting in the most cost effective nutrient reductions.	A workgroup produces a report targeting BMPs by type and location.
B2. Cost share providers prioritize assistance for targeted BMPs and track implementation	Amount spent on targeted practices relative to non-targeted practices.
C. Secure and leverage funding for BMPs.	
C1. Conduct a workshop to examine and enhance BMP financing strategies.	Recommendations for improving BMP financing strategies developed.
C2. Utilize DNREC's Water Quality Improvement Project Sponsorship Program (WQIPSP) to leverage funding for BMPs.	WQIPSP dollars spent on BMP implementation.
D. Promote and reward those in the agriculture sector who are good stewards of the environment.	Members of the agricultural sector are recognized publically for their innovation and BMP implementation.
E. Improve nutrient management of developed lands through research and education to better quantify and reduce nutrient loads.	
E1. Conduct watershed specific analysis to determine nutrient loading to the Bays from developed lands under different management practices.	A report is produced on nutrient loading from developed lands including education and management recommendations.
F. Develop program to educate the general public and landscapers on the benefits of reducing fertilization and improving fertilization practices.	Number of individuals and landscapers educated on the benefits of improving fertilization practices.
Focus Area: Wastewater Management	
Objective 1. Examine, improve and update existing on-site wastewater treatment and disposal regulations and their enforcement.	
Actions	Performance Measures
A. Annually assess and update information on regulatory initiatives in the onsite wastewater sector.	
A1. Continue and report on DNREC holding tank inspection program.	Regular reports on the holding tank inspection program are publicly available.
A2. Report compliance with DNREC pump-out and inspection requirements for septic systems on properties that are sold.	Regular compliance reports are publicly available.
A3. Verify that all new and replacement septic systems in the Inland Bays watershed are required to meet all regulatory performance standards.	Regular reports on replacement septic systems are publicly available.
B. Promulgate and enforce revisions to DNREC's onsite wastewater treatment and disposal regulations.	Regulations are promulgated and enforced.
C. Ban permanent holding tanks in the watershed.	A ban is in place and remains as such.
Objective 2. Examine emerging contaminants entering the Inland Bays and engage the regulatory community and general public in education	
Actions	Performance Measures
A. Conduct a symposium that identifies emerging contaminants, their sources, and their potential effects.	Emerging contaminant symposium held and findings conveyed.
B. For emerging contaminants with the highest potential for significant environmental impact, prepare reports to define the problem and promote source control.	Number of emerging contaminants reported upon.
C. Determine the need for regulations to reduce the threat of identified emerging contaminants.	Position on regulations provided by regulatory agencies.
D. Inform the public about the potential threats, challenges, and solutions to identified emerging contaminants.	A public education campaign on emerging contaminants is developed and implemented.

Objective 3. Promote the use of regional wastewater treatment and disposal systems within designated growth zones over multiple small	
Actions	Performance Measures
A. Develop a wastewater planning committee comprised of DNREC, Sussex County, utility industry representatives, and other stakeholders to coordinate the treatment and disposal of wastewater from new and existing developments based on the TMDLs of receiving waters.	The wastewater planning committee is formed and meets regularly.
B. Conduct workshop to share new technology and incentives for increasing the beneficial reuse of wastewater.	Workshop results in an increase in the number of beneficial re-use projects.
C. Require surface water assessments that clearly demonstrate how all proposed wastewater treatment and disposal systems will help to meet TMDLs for receiving waters.	Regulations or permit requirements are implemented that require consistency with TMDLs.
D. Enforce the waters of Exceptional Recreational and Ecological Significance (ERES) provisions of the State Water Quality Standards requiring the least environmentally damaging disposal alternatives for wastewater.	ERES provisions are enforced for wastewater disposal.
E. Develop a nutrient budget for wastewater to determine existing and projected total wastewater loads to receiving waters.	Nutrient budget for wastewater is developed and used for planning and permitting.
F. Research the attenuation of nutrients and contaminants released from different types of on-site wastewater systems along flowpaths to receiving waters.	Information on attenuation developed and used to guide permitting.
Stormwater Management	
Objective 1. Reduce nutrient contributions from stormwater to help achieve TMDLs.	
Actions	Performance Measures
A. Complete the revision and implementation of sediment and stormwater regulations.	Revised sediment and stormwater regulations are promulgated.
B. Create stormwater management facilities and source reduction strategies for 4,500 acres of urban and residential lands developed pre-1990.	Acres developed pre-1990 treated by stormwater retrofits.
C. Provide assistance to local governments and HOAs to draft ordinances that minimize new and reduce existing impervious surfaces.	The number of governments and HOAs assisted with impervious surfaces ordinances.
D. Engage corporate partners to include Green Infrastructure practices in new and redevelopment projects.	Number of corporate partners adopting Green Infrastructure practices.
E. Encourage Sussex County and/or municipalities to create a stormwater utility to fund maintenance and retrofits.	Stormwater utility(ies) created.
F. Develop and implement a lines and grades/drainage code for Sussex County.	Sussex County adopts a lines and grades/drainage code.
G. Develop impervious surface coverage targets to protect aquatic life and urge their inclusion into county and municipal comprehensive plans.	Maximum impervious surface coverage targets developed. Number of comprehensive plans including maximum impervious surface coverage targets.
Water Quality Management	
Objective 1. Update the Inland Bays estuarine and watershed models with the latest scientific understanding and best available data, and	
Actions	Performance Measures
A. Update the Inland Bays estuarine water quality and hydrodynamic model.	An updated model populated with the best available data is available for use.
B. Update the Inland Bays watershed nutrient loading model.	An updated model with the best available data is available for use.
C. Utilize updated estuarine and watershed models to evaluate if existing TMDLs are adequate to achieve water quality standards for nitrogen and phosphorus.	A report is produced. TMDL regulations are updated as needed.
Objective 2. Report on the implementation of the PCS, revise and prioritize remaining actions, and devise an implementation plan to meet the	
Actions	Performance Measures
A. Produce initial report on PCS implementation and identify barriers to implementation.	Initial report on PCS implementation is published.
B. Revise PCS goals as needed, incorporating any revisions to the TMDLs.	PCS goals revised as needed incorporating any revisions to the TMDLs.
C. Develop an implementation plan for remaining PCS actions that includes a time frame for completion, interim goals, and identified implementation funding sources.	A revised PCS with implementation plan is published.
D. Produce annual PCS progress reports including a yearly determination of the nutrient loads to the Bays and their tributaries relative to their TMDLs.	Annual PCS progress reports are generated.

Objective 3. Review and revise State and local standards for ground and surface water protection.	
Actions	Performance Measures
A. DNREC reviews their technical standards for ground and surface water protection at five-year intervals.	Review is publicly available.
B. Obtain and review County standards for ground and surface water protection.	Such a review is publicly available.
Objective 4. Quantify the transport of contaminants from Indian River Power Plant (IRPP) coal ash landfills to receptors in the aquatic	
Actions	Performance Measures
A. Study the transport of contaminants to aquatic life near the IRPP to inform the Voluntary Cleanup and the Natural Resources Damage Assessment Processes for the Burton Island coal ash landfill.	Final reports made available to the public.
B. Model transport scenarios of contaminants from the IRPP coal ash landfills to environmental receptors based on various levels of sea level rise and severe storm impacts.	A report on this scientific research is made available to the public.
Objective 5. Reduce nutrient input to residential canals and lagoons.	
Actions	Performance Measures
A. Treat or remove greywater discharges into tributaries, canals, and lagoons.	Number of greywater discharges treated or removed.
B. Filter runoff from roofs, driveways, and other impervious surfaces.	Number of projects implemented.
C. Provide and disseminate educational material for homeowners on reducing fertilizer inputs to tributaries, canals, and lagoons.	Materials are distributed.
D. Examine dead-end canals to determine if any could benefit from low-cost solutions to increase flushing.	Candidate sites for additional tidal flushing are identified.
Objective 6. Re-assess water quality monitoring efforts for their representativeness and capacity to detect trends, then develop recommendations	
Actions	Performance Measures
A. Aggregate historic and contemporary water quality monitoring data and metadata into one publicly accessible database.	Database is available and updated annually.
B. Conduct a long-term trend analysis of water quality parameters.	The trend analysis is completed and published.
C. Develop recommendations to improve efficacy of monitoring efforts to detect trends.	Recommendations report is published.

Managing Living Resources and Their Habitat	
Objective 1. Promote recurrence of bay grasses.	
Actions	Performance Measures
A. Conduct an education initiative on the benefits and importance of re-establishing submerged bay grasses.	Number of individuals educated about bay grasses and their restoration.
B. Map areas of the Bays that have habitat characteristics supportive of the re-establishment of bay grass species that have been identified as suitable candidates for restoration.	A report including data layers and maps is produced.
C. Convene a bay grass restoration workgroup to develop a bay grass restoration, protection, and monitoring plan.	Baygrass restoration workgroup is active. A plan is developed.
D. Implement the bay grass restoration, protection, and monitoring plan.	Number of acres successfully restored. Annual restoration and monitoring reports are produced.
Objective 2. Halt the continued loss of wetlands and reverse these loss trends by promoting projects to mitigate for previously lost wetlands.	
Actions	Performance Measures
A. Bring regulation of freshwater wetlands, including isolated wetlands, under State jurisdiction and permitting.	State legislation is passed and regulations are adopted.
B. Identify candidate sites for the creation and restoration of wetlands.	Numbers and acreages of sites identified.
C. In accordance with the Inland Bays Pollution Control Strategy (PCS), create or restore wetlands on areas previously converted to cropland.	The PCS goal of restoring 4,147 acres is met.
D. Protect and enhance/restore additional wetland acreage.	Number of acres protected. Number of acres enhanced/restored/created per year.
E. Encourage the planting of trees and other plants adjacent to all wetlands.	Number or acres of planted next to wetlands.
F. Revise the existing Sussex County Ordinance on water quality buffers to be in line with the CIB's Recommendations for a Water Quality Buffer System.	A revised ordinance is adopted.
G. Engage the state Natural Areas Advisory Council to help identify freshwater wetlands that should be purchased as preserves.	A prioritized list of properties to be purchased is produced.
H. Develop a living shoreline initiative to maximize the amount of natural Bay shorelines.	
H1. Assess and report on the condition of shorelines in the Inland Bays.	Shoreline assessment reports are produced.
H2. Conduct an education and outreach program on shoreline function and management alternatives for shoreline property owners.	Number of shoreline property owners informed.
H3. Conduct living shoreline demonstration projects to train installation and maintenance contractors.	Length of living shoreline enhancements or stabilizations installed. Number of contractors trained.
H4. Demonstrate innovative living shoreline stabilization techniques utilizing bay grasses, shellfish, and other native biota where feasible.	Number of demonstration projects completed.
H5. Support legislative and/or regulatory changes needed to require that living shoreline techniques be employed wherever feasible for shoreline stabilization.	Legislation is passed or regulations are updated.
Objective 3. Provide access for native migratory fish to upstream areas for use as spawning and/or nursery sites.	
Actions	Performance Measures
A. Conduct a migratory fish passage restoration feasibility and planning study.	The study is completed .
B. Implement fish passage restoration projects.	Number of passage projects completed. Number of miles of fish habitat restored.
C. Monitor fish passage restoration success.	Annual reports are produced that document the number or percentage of target migratory fishes utilizing the passages.
D. Conduct education and outreach efforts on the importance of migratory fishes and the benefits of fish passage restoration.	Number of people informed.
Objective 4. Eliminate once through cooling at the Indian River Power Plant (IRPP).	
Actions	Performance Measures
A. Track progress of IRPP compliance with DNREC agreements for removal of Unit 3 water withdrawals by January 1, 2014.	The only water being withdrawn at IRPP is to compensate for evaporative loss at the cooling tower.

Objective 5. Increase the economic and environmental benefits of shellfish.	
Actions	Performance Measures
A. Increase the acreage of approved shellfishing waters.	
A1. Examine water quality data for the past 5 years to determine if areas of the Bays could be re-opened to shellfish harvest.	Number of acres reclassified from closed to approved or seasonally approved.
A2. Determine the sources of contamination that presently constrain the opening of additional shellfishing areas.	A listing of the sources of contamination is available for public scrutiny.
A3. Develop and implement a strategy to address contaminant source reduction so that additional shellfishing waters may be opened.	The strategy is completed and implementation is underway .
B. Enhance populations of eastern oysters.	
B1. Create additional hard bottom areas suitable for oyster recruitment or planting of oyster spat.	Acres of suitable hard bottom areas created.
C. Promote and encourage shellfish aquaculture in the Inland Bays.	
C1. Convene a team of state and federal regulatory representatives and stakeholders to produce the scientific, educational, and policy groundwork necessary to develop legislation and regulations that govern shellfish aquaculture in the Inland Bays.	Legislation favorable to aquaculture is passed. PM C1.2 Regulations governing aquaculture are promulgated.
C2. Provide financial incentives for new aquaculturists.	Financial incentives are competitive with other states.
C3. Provide technical support and education to aquaculturists.	Technical guidance specific to the Inland Bays is published.
Objective 6. Monitor and control the spread of invasive species within the Bays and their watershed.	
Actions	Performance Measures
A. Map the known distributions of invasive species of concern in the watershed.	A map is created and publicized.
B. If needed, support implementation of policy designed to curb the spread of invasive species.	The legislative/and/or regulatory process is engaged to limit the spread of invasive species.
Planning for Climate Change	
Objective 1. Integrate projected sea level rise into land use planning and proposed development to protect shore zone ecosystems and bay	
Action	Performance Measures
A. Work with the County and municipalities to incorporate sea level rise into comprehensive plans.	The County includes sea level rise in its comprehensive plan. Percentage of bayside municipalities that include the projected impacts of sea level rise in their comprehensive plans.
B. Conduct a sea level rise vulnerability analysis specific to the Inland Bays watershed that includes potential impacts to both green and gray infrastructure.	Report of analysis is publicly available.
C. Implement the recommendations of the State Sea Level Rise Advisory Committee relative to the Inland Bays when they become available.	Percentage of total recommendations implemented.
D. Model the distribution of tidal wetlands under different sea level rise scenarios to guide land use and protection decisions that maximize future tidal wetland extent.	Number of additional acres that would need to be acquired or protected.
E. Track shifts of dominant aquatic species potentially caused by climate change through the use of previous and recent surveys.	Generation of a list of species affected that is regularly updated.
F. Include climate change and sea level rise information in public outreach and education efforts.	Number of individuals informed about climate change and sea level rise.

Coordinating Land and Water Use Decisions	
Objective 1. Involve all levels of government to obtain commitments for coordination of land use decisions that minimize environmental	
Action	Performance Measures
A. Designate the Inland Bays watershed as a 'Critical Environmental Area' and manage the watershed for nutrient reductions consistent with TMDL load reductions or reductions attributed to best available technologies.	Designation of the Inland Bays watershed as a 'Critical Environmental Area'.
B. Request that representatives of all levels of government sign a letter of understanding that their land use decisions will minimize environmental impact to existing aquatic resources in the watershed.	Such a letter is drafted and signed by appropriate agencies represented on the CIB Board of Directors.
Objective 2. Provide maximum protection of waterways, forested stream corridors, groundwater, natural areas, open space, tidal and non-	
Action	Performance Measures
A. Water quality buffers shall be clearly demarcated, designated, and recorded on final site plans or final major subdivision plats and demarcated on the ground with signs or other kinds of markers.	This requirement is included in a revised Sussex County ordinance.
B. Maintain land presently classified as open space under County or municipal ordinances or codes to minimize nutrient loading to the Inland Bays estuary.	County and municipal officials are educated on the need for such maintenance.
C. Update and implement the Inland Bays Habitat Protection Plan.	Number of acres protected through acquisition or easement.
D. Use the Delaware Ecological Network and other appropriate information source to prioritize the preservation of key habitat in the Inland Bays drainage system.	Number of acres protected.
Objective 3. Update and implement the Inland Bays Water Use Plan.	
Action	Performance Measures
A. Assess implementation progress of the Water Use Plan and revise remaining and new actions.	Water Use Plan Update is published.
B. Focus outreach on increasing waterway safety and channel marking.	Waterway maintenance improves.
C. Focus on low impact water use activities.	The public is informed about the availability of low impact water use activities.
D. Continue marine spatial planning efforts to maximize aquatic resources and minimize water use conflicts.	
D1. Develop a publicly accessible marine spatial planning database.	A database is publicly available.
D2. Provide educational and planning forums on spatial aspects of water uses.	Number of individuals attending forums.
Outreach and Education	
Objective 1. Increase the visibility of the CIB and its mission.	
Action	Performance Measures
A. Assess the CIB brand and communications strategy to determine effectiveness.	
A1. Conduct surveys to gather data on citizen perceptions and understanding of issues of concern in the watershed.	Surveys are developed and implemented.
B. Identify and implement high priority Work Plan/CCMP actions and develop outreach and education campaigns.	Campaign(s) are developed and implemented.
Objective 2. Educate stakeholders in the watershed about their impacts on water quality in the Bays and how they can help.	
Action	Performance Measures
A. Develop and deliver watershed education programs for children.	
A1. Programs for school age children are offered at the James Farm Ecological Preserve.	Number of students attending programs.
A2. Offer community outreach and education to children, families, and visitors at the Bethany Beach Nature Center.	Programs offered at BBNC.
A3. Continue to offer watershed education at schools through the Schoolyard Habitats Program.	Number of schools reached annually at their schools.
B. Administer a Speakers Bureau.	Number of speaking engagements annually.
C. Continue to promote the 1000 Rain Gardens for the Inland Bays program.	A demonstration rain garden is established in every incorporated town in the watershed. Rain Garden booklets are distributed at demonstration sites.

Objective 3. Communicate with stakeholders through a variety of media; to promote public involvement and influence behaviors, attitudes	
Action	Performance Measures
A. Continue to develop and administer a website as a primary vehicle for disseminating information	A comprehensive website is maintained.
B. Incorporate social marketing and enhanced use of media into CIB's communications strategy	Social media is used to maintain regular communication with Board and Board Committees, volunteers, Friends of the Bays, and elected and public officials.
C. Edit and disseminate a newsletter and annual report.	The Inland Bays Journal is published and distributed three times each year and an annual report is published annually.
D. Create and disseminate printed marketing materials such as brochures, postcards, flyer exhibits and signage to address specific education/outreach needs to target audiences.	Printed materials and exhibits are produced each year for priority issues and projects.
E. Maintain relationships with local media outlets and reporters and disseminate press releases and photos for their use.	Press releases are issued for events and selected projects.
Objective 4. Encourage more stakeholder support through volunteerism.	
Action	Performance Measures
A. Direct a volunteer program that provides citizens a formal track to partner with the CIB.	Volunteer opportunities are developed and volunteers receive regular invitations to assist.
B. Involve volunteers and stakeholders in demonstration projects that model desired changes in practices and citizen science research.	Number of volunteer hours worked each year.
Objective 5. Communicate environmental results to inform legislators and raise citizen awareness about the state of the Inland Bays and its	
Action	Performance Measures
A. Results of Inland Bays environmental studies or projects are published.	A "State of the Inland Bays Report" is published and disseminated every five years. Press releases are issued to provide new information to the media.
B. Communicate the benefits to economic development, tourism, recreation and quality of life of achieving water quality goals.	Increase in participation from the business community.

APPENDIX B
ENVIRONMENTAL MONITORING PLAN REVISION PROCESS AND MEETING NOTES

ENVIRONMENTAL MONITORING PLAN REVISION PROCESS

Ambient Water Quality Monitoring Workshop

As part of the IBEMP update, the Center for the Inland Bays (CIB) hosted a facilitated workshop on July 29, 2015 to discuss ambient water quality monitoring in the Inland Bays. The workshop was facilitated by a contractor, Jim Eisenhardt, of RK&K. Primary goals of the workshop were to:

1. Review current ambient water quality monitoring programs and identify strengths, weaknesses, and areas of improvement;
2. Develop a process to transfer existing and new University of Delaware Citizen Monitoring Program (CMP) data to a public database such as STORET;
3. Include CMP data effectively into the combined Watershed Assessment Report (305(b)) and Determination for the Clean Water Act Section 303(d) List of Waters Needing TMDLs;
4. Evaluate the effectiveness, capacity, and costs of the DNREC water quality portal (developed by DEMAC) to provide DNREC and CMP data to researchers and the public; and
5. Improve the participation of organizations in the review and interpretation of Inland Bays environmental indicator reports.

Prior to the workshop, participants were asked to provide input on key questions to help guide discussion; compiled results are included below (pages B-3 through B-17). Participants represented the following organizations/agencies: CIB, Delaware Department of Natural Resources and Environmental Control (DNREC), University of Delaware, and the United States Geological Survey (USGS). Representatives from the United States Environmental Protection Agency (EPA) Region III were not in attendance; however, Bill Richardson of the Office of Standards, Assessment and TMDLs provided input prior to the workshop.

Workshop notes are included below (pages B-18 through B-25).

Scientific and Technical Advisory Committee Input and Review

At its meeting on September 18, 2015, the STAC discussed the IBEMP update. The discussion was facilitated by Jim Eisenhardt (RK&K) and the Center's Science and Restoration Coordinator, Marianne Walch. The focus of the discussion was existing monitoring programs and anticipated monitoring needs.

A questionnaire similar to the one used for the facilitated workshop was sent to key STAC members prior to the meeting. Compiled results of that questionnaire (pages B-26 through B-32) and notes from the discussion held at the STAC meeting (page B-33) are included below.

Compiled Questionnaire for WQ Monitoring Workshop Meeting

29 July 2015

In order to gather information to help guide our discussions at the July 30th workgroup meeting, we ask that you provide answers to the following questions about your own agency/organization's monitoring programs in the Inland Bays, as well as your thoughts about improving the Center for Inland Bays (CIB) monitoring plan¹.

Attendees/Responders

- BR: Bill Richardson, EPA Region 3, Office of Standards, Assessment and TMDLs
- DW: Dave Wolanski, DNREC, Watershed Assessment
- HM: Hassan Mirsajadi, DNREC, Division of Watershed Stewardship, Watershed Assessment and Management Section
- RG: Rick Greene, DNREC, Division of Watershed Stewardship, Watershed Assessment Section
- RT: Robin Tyler, DNREC, Division of Water/Environmental Laboratory Section
- JY: Joanna York, University of Delaware, School of Marine Science and Policy
- SA: Scott Andres, Delaware Geological Survey/Chair CIB STAC
- BU: William (Bill) Ullman, University of Delaware, School of Marine Science and Policy, Oceanography Program
- JF: Joe Farrell, University of Delaware, Delaware Sea Grant, UD Citizen Monitoring Program
- EW: Edward Whereat, University of Delaware, UD Citizen Monitoring Program

¹Per email from Chris Bason on 22 July: Is there anyway EPA can communicate its desire to have citizen science data operationally included into the determination of water quality impairment (or lack thereof)? Important role for EPA to encourage States to do this. Maybe there will still be ways in which EPA can support our effort to do this in DE and I think it would be fantastic if DE and Region III could be a national leader in this. Nancy Laurson [EPA HQ]: Monitoring Plan should fit the local needs.

Per Bill Richardson: It would be EPA's preference for any data collected by the citizen scientists working with CIB be used in DE's Integrated Reports (IRs).

1. What areas, based on your experience/expertise, are the most important to focus on in a water quality monitoring plan for the Inland Bays and (very briefly) why?

- BR:
 - Eutrophication; habitat and wetland loss.
 - Bacteria for protection of recreation.
- DW: The DNREC monitoring stations are pretty extensive and cover the largest waterbodies with enough stations to give us big picture conditions, provide some modelling data and track trends in the watershed as a whole.
- HM:
 - Water quality monitoring at headwaters and smaller streams.
 - Groundwater quality and movement.
 - Ecological conditions including habitat and biology.
 - Wetland assessment.
 - Since we have an acceptable level of water quality monitoring of larger streams and Bays that is currently conducted by DNREC and Citizen groups, the above four areas may need additional attention.
- RG:
 - Nutrients and algal productivity (Why: Eutrophication, dissolved oxygen, fisheries, benthic habitat and aesthetics).
 - Bacteria (Why: Shellfish harvesting and primary contact recreation).
- RT: I think that the most important areas (factors/variables) to focus on within the water quality element of the Inland Bays Monitoring Plan are those presently being focused upon – dissolved oxygen, nutrients, chlorophyll, and pathogen indicator bacteria.
- JY: Monitoring of chlorophyll, DO, and nutrients. Potentially also biomass of macroalgae, depending on balance between micro- and macro-algal primary producers. Could also consider light—even via Secchi disk.
- SA:
 - Spend monitoring dollars wisely. This can be accomplished if you:
 - 1) Clearly articulate the goals and objectives of monitoring.
 - 2) Develop protocols (including QA/QC) and document metadata consistent with the protocols.
 - 3) Communicate 1) and 2) to the monitoring program staff and volunteers
 - Recognize and avoid mission creep. This dilutes the effort and spreads staff too thin.
 - Do what is necessary to have monitoring data incorporated in or linked to larger state, regional, or national datasets. “Spending monitoring dollars wisely” will guide the program to this objective.
- BU:
 - Monitoring needs to yield data in a form that can be usefully related to both status and fluxes. Fluxes are harder to determine on ecologically relevant time scales, but these fluxes are needed to appropriately set goals for long term

preservation. Spot samples are easy to collect, but much harder to use as these spot analyses need to be integrated into flow models to yield fluxes.

- We have been successfully using continuous automated sensors in the Indian River, Nanticoke River and Murderkill River watersheds to supplement spot measurements for a number of years and have been using these data to estimate fluxes and to answer specific scientific questions in these systems.

- JF:

- Dissolved oxygen as good overall indicator of living resource and Bay health.
- Nutrients (N and P) load and concentration - as long term indicator of whether Bay conditions are improving.
- Pathogen indicator – for recreation water quality and safe shellfish consumption.

- EW:

- Causes of eutrophication, i.e. nutrient levels - needed to assess trends.
- Biological responses to eutrophication, i.e. low dissolved oxygen levels, reduced water clarity, high chlorophyll levels, (harmful algae blooms?) - needed to assess trends.
- Recreational contact safety - total Enterococcus levels - recreational use and public interest.
- Habitat conditions are important, but difficult to address.

2. What monitoring activities/data collection is your organization undertaking in the Inland Bays that could be part of the CIB WQ monitoring network?

- BR: EPA's national survey may collect Inland Bay sites, but the survey is only conducted every five years. The data may not be sufficient to supplement CIB needs.
- DW:
 - DNREC publishes the monitoring plans online:
<http://www.dnrec.delaware.gov/swc/wa/Pages/WaterQualityMonitoring.aspx>
 - In lieu of Table 1, I am attaching our most up to date monitoring plan² which is being updated now. No significant changes are expected at this time.
- RG:
 - Routine monitoring of selected metals at multiple stations (see monitoring plan provided by Dave Wolanski for details).
 - Toxics: Work is done when a specific need arises and is justified scientifically.

² State of DE Ambient Surface Water Quality Monitoring Program – FY 2015. Available online at:

<http://www.dnrec.delaware.gov/swc/wa/Documents/WAS/State%20of%20Delaware%20FY%202015%20Ambient%20Surface%20Water%20Quality%20Monitoring%20Program%20.pdf>.

- Toxics in Biota Monitoring Program collects and tests fish tissue and shellfish samples for chemical contaminants.
- Toxics in Sediment Monitoring Program assesses toxic contaminants in sediment samples (mostly collected by other programs).
- SA: I cannot overstate the importance of collecting good quality data on the flow of water. Water flow is the single most important variable in computing pollutant loads.
 - Coordinate and pass funding to U.S. Geological Survey
 - Being done with DEOS/DEMAC
 - Being done with DEOS
- BH: Our group does not have any present monitoring efforts in the Inland Bays watershed.

PLEASE COMPLETE TABLE 1 AT THE END OF THE QUESTIONNAIRE. Feel free to provide additional comments here.

3. What water quality monitoring programs are planned for the future that might be applicable to the Inland Bays?

- BR: EPA funds a National Coastal Conditions Assessment (NCCA) every five years. The NCCA is ongoing this summer in 2015. The NCCA is a probabilistic survey and there are normally a few sites located within the DE Inland Bays. DNREC staff are provided grant funding to collect DE samples.
- DW: See response to question #2.
- RG: The Watershed Approach to Toxics Assessment and Restoration (WATAR) might be applicable to the Inland Bays in the future.
- RT: No knowledge of new programs.
- BU: We have no current plans for the Inland Bays watershed, although, we have submitted a proposal with Scott Andres, Delaware Geological Survey, to instrument and monitor the effects of the Wolfe Neck Regional Wastewater Treatment Plant on nutrient concentrations and loads leaving this site for surface waters of the Lewes-Rehoboth Canal. This work, if funded, would begin in Spring 2016.
- EW: Nothing applicable at this time.

PLEASE COMPLETE TABLE 2 AT THE END OF THE QUESTIONNAIRE. Feel free to provide additional comments here.

4. What areas, based on your experience/expertise, are missing from the current Inland Bays WQ monitoring plan?

- BR: EPA would recommend CIB coordinating with DNREC to ensure adequate data are collected for use in DE's biannual Integrated Report (305(b) and 303(d)). EPA can assist with DNREC-CIB discussions if needed.
- HM: The 4 areas mentioned in question #1 above.
- RG:
 - Continuous monitoring of dissolved oxygen, temperature, pH and conductivity.

- Habitat types and quality.
- RT: Establishing bacterial source tracking as a routine element of water quality monitoring would seem to have value because it may dampen much of the present accusation/denial conflict regarding such sources and give more pointed and effective direction to pollution abatement efforts. Also, if bacteria contributions are linked to human sources then it is reasonable to assert that those sources are also accountable for some larger proportion of the loading of other factors of interest such as organic matter and associated nutrients (i.e., actions corrective to the input of pathogen indicator bacteria are also likely to yield improvements in the levels of organic matter and nutrients).
- SA: The current plan relies almost entirely on DNREC action and political pressure. There is an overwhelming influence of the TMDL, PCS, and 305b programs on monitoring and the data that are regularly assessed and reported on by DNREC staff. Changing any aspect of monitoring or assessment has consequences for hard-won progress with the TMDL and PCS. The fear of consequences has stopped ideas for new ways to collect or assess data.
- BH: I think that too little is known about the variable impacts of agricultural and domestic wastewater on total nutrient loads to the Inland Bays. With population increases, I am concerned that total loads to surface waters will increase, even as better treatment and disposal options are put into place. I think that we need to have better estimates of these loads and develop robust models for estimating future loads from these sources as population increases in the future.
- JF: Not sure.
- EW:
 - In general, inadequate frequency and spatial coverage.
 - Episodic events (extreme rain, wind and tide events).
 - Winter conditions.
 - Targeted sampling to address specific issues or determine effectiveness of BMPs.
 - Toxicity studies of fish, shellfish, and sediments are limited

5. What additional WQ monitoring activities would your organization benefit from if CIB could include in their monitoring plan? Please feel free to think outside of the box, without regard to availability of funding, including new or innovative technologies.

- BR: Continuous monitoring of DO and pH to evaluate nutrient impacts (if not on-going).
- BR: Research on possible impacts from ocean acidification.
- DW: Continuous monitors might be valuable for evaluating DO criteria in tributaries.
- RG: An innovative study of engineered denitrification of groundwater.
- RT: The DNREC Laboratory is exploring the efficacy of doing bacterial source tracking testing as one of its services. The feasibility of successfully doing so is contingent upon having a steady enough flow of samples (work) to make the venture viable. Establishing

bacterial source tracking as a routine test within long-term monitoring and special project sampling efforts is essential and a commitment from the Inland Bays Monitoring Plan would be helpful in advancing this.

- JY: Nutrient monitoring. Perhaps in situ monitoring instrument - similar to the LOBO instrument that is installed at Bowers Beach (Bill Ullman) or the instruments maintained by the DNERR. Expensive, but provides LOTS of data.
- SA: Deployment of automated WQ sensors. In addition to standard physical sensors, nitrate sensors provide invaluable data.
- BU: We have considerable and growing expertise in the use of continuous water quality monitoring devices for coastal watersheds and estuaries. I think that it is more likely that we could assist the CIB and the State than vice versa, but we would be interested in expanding the use of these systems and developing automated applications of these data for managers.
- JF: Phytoplankton/HAB monitoring plan for recreational and commercial shellfish harvesting/aquaculture.
- EW:
 - Continuous monitoring in some tributaries.
 - Bacterial source tracking.
 - Toxicity of harmful algae.

6. Does your organization have historical WQ or other monitoring data that were not included in the original monitoring plan that would be useful for future WQ monitoring activities or for use as State of the Bays indicators?

- BR: Possible NCCA data, but for only a few sites.
- DW: DNREC publishes all its data through STORET, online at: <http://www.epa.gov/storet/>
- HM: We have several years of high frequency water quality and nutrient data from Millsboro Pond outlet.
- RG: Possibly.
- RT: Yes.
- JY: Probably not. I assume folks from UD have been engaged with plans at the CIB and State level and would have given input.
- SA: The Monitoring Plan does not include a concise bibliography, database, or list of data. It is possible that data held by the Delaware Geological Survey are missing, but I don't have enough time to do this detective work by July 28.
- BU: I believe all of our data were made available to DNREC and CIB. Not all of the data collection was part of the initial Monitoring Plan. Much of the data that we collected historically, have also been published.
- JF: More than a decade of comprehensive phytoplankton/harmful algal bloom (HAB) data for Inland Bays.
- EW:
 - Bacteria (TE) levels.
 - Select harmful algae species.

7. Does your organization anticipate having sufficient future funding to carry out WQ monitoring activities that CIB has included in the monitoring plan to date? Do you anticipate future funding for additional data collection parameters beyond the current monitoring criteria included in CIBs plan?

- BR: No.
- DW: Funding is always challenging. The State budget is going to be interesting the next few years.
- HM: Our surface water quality monitoring program has changed from what is in the CIB plan based on our emerging and changing needs. However, our current level of monitoring is much more than what is in the CIB plan. And, we expect to have sufficient funding to continue our current level of monitoring into the future.
- RG: Unknown.
- RT: The “Priority Basin Monitoring” (aka General Assessment for 305(b) Water Quality Inventory reporting) is anticipated to continue at least at the level of effort of the past 25 years. Biological (macroinvertebrate) monitoring is likely to continue, probably at five-year intervals. Other Clean Water Act mandated monitoring such as NPDES will also continue as will other permit-related programs in the Division of Water.
- JY: Funding is dependent on successful grant applications. It is hard to anticipate beyond a two-year timespan.
- SA: The plan has NO activities for my agency only work that might be done if resources were available. Table 2 lists potential data collection efforts.
- BU: All of our work required extramural funding. Other than the proposal described in item 3, above, we have no additional plans for monitoring activities in the Inland Bays.
- JF: We will need support to continue some of our past monitoring efforts that are no longer funded e.g. nutrients.
- EW:
 - To some degree, depending on scope.
 - Doubtful at this time.

8. Do you have any recommendations for additional funding sources for WQ monitoring activities?

- BR: None at this time.
- DW: No.
- HM: EPA, NOAA, NSF.
- RG: No.

- RT: Increases in environmental monitoring funding levels via the government sector seem unlikely in the present economic and political climate, absent a crisis such as that triggered by Pfiesteria almost 20 years ago. More productive perhaps would be grants from the private sector, particularly large foundations or endowments.
- JY: Collaborative funding proposals to the EPA/NSF/NOAA for longer term funding, potentially including the CIB, academic researchers, and citizen groups.
- SA:
 - Private entities could purchase monitoring equipment for CIB to use or borrow for monitoring (see question #5).
 - EPA NEP funds could potentially fund migration of the CMP data into STORET. From there, the data are accessible to everyone through multiple channels.
- BU: There is very little funding available for just monitoring. However, data collected as part of scientific research can also serve a secondary monitoring objective. Both DNREC and Sea Grant have funded such dual-use projects in the past.
- JF: I wish we did.
- EW:
 - DNREC/State should devote more funds to monitoring.
 - CIB should devote more funds to monitoring.

9. Other than the University of Delaware Citizen Monitoring Program, are you aware of any volunteer organizations/programs that are currently active and could be included in future Inland Bays WQ monitoring for supplemental information?

- BR: No.
- DW: No.
- HM: No.
- RG: No.
- RT: No.
- JY: No.
- SA: No.
- BU: No.
- JF: South Bethany Water Quality Committee.
- EW: I am not aware of other volunteer organizations/programs, perhaps with exception of the South Bethany Water Quality Committee, which falls under our umbrella, but has certainly become semi-independent and has been involved in CIB BMP projects.

10. For the data you discuss in this document, is your organization just collecting raw data or are you also processing the data and developing trends from the data?

- BR: Raw data only. Reports from the NCCA are written at a national scale.
- DW: We collect and process the data for a number of programs and purposes.
- HM: Our Section develops the monitoring plan and reviews and analyzes the data.
- RG: Watershed Assessment has compiled and assessed toxics data generated through its own monitoring and monitoring performed by others to produce a better understanding of an issue (e.g., arsenic in surface water, ground water, soils, sediments, fish tissue and air).
- RT: Mostly data collection, with some analysis.
- JY: I also process the data and interpret mechanisms and trends.
- SA: We also process and interpret data.
- BU: We are only interested in projects where monitoring data can be put to scientific use.
- JF: We provide semi-monthly summary reports during summer and are working on providing trend information, but we need additional staff support and resources to develop and maintain trend data in more accessible form.
- EW: Our organization has tended to focus on collection of raw data, but we do process and report data (reports issued regularly during summer, and data summaries submitted for 305b report), and have participated in prior CIB's State of the Bays reports.

Table 1

Attendee	Program/Monitoring Activity	Purpose/users	Where? (bay(s), stream(s), watershed(s), etc.)	Years collected, frequency of sampling	Parameters measured	Current/future status	Funding Status/Sources
HM	Surface water quality monitoring – Fixed Station, grab samples	Assessment of surface water quality conditions	About 24 stations in the tributaries and main Bays	30+ years of data with different frequencies Monitoring frequency during FY 2016 is monthly	About 30 parameters including nutrients, organic, physical, bacteria, etc.	Will monitor monthly for 2 yrs, then will monitor 6 times/yr for the next 3 yrs	State General Fund, EPA 106 grant
HM	Continuous WQ monitoring (datasonde)	To monitor short term changes of dissolved oxygen and other parameters	Millsboro Pond outlet Massey Ditch	3+ yrs 15 minutes data collection	DO, temp, salinity, pH	Will continue monitoring during FY 2016	EPA 106 grant
HM	Stream gages	Measuring stream discharge	Millsboro Pond outlet Beaverdam ditch near Millville	20+ yrs	Stream discharge	Will continue during FY 2016	State General Fund, EPA 106 grant
HM	Tide gages	Measuring	Rosedale Beach	20+ yrs	Tidal elevation	Will continue	State General

		tidal elevations	Indian River Bay near Bethany Beach Beaverdam ditch near Millville			during FY 2016	Fund, EPA 106 grant
RG	Toxics in Biota Monitoring (i.e., testing of fish and shellfish for chemical contaminants)	To determine need for fish consumption advisories	Indian River Bay Rehoboth Bay Millsboro Pond	Infrequent	PCBs, dioxins/furans, OC pesticides, PAHs, arsenic, and mercury	Low priority compared to State waters with known toxic impacts	State & Federal
RG	Toxics in Sediment Monitoring & Assessment (data collection leveraged through dredging projects and EPA Coastal Assessment Program)	Assess ecorisk to benthic aquatic life and potential human health risk through sediment to biota transfer	Indian River Bay Rehoboth Bay	Infrequent	PCBs, OC pesticides, PAHs, and metals	Low priority compared to State waters with known toxic impacts	State & Federal
RT	General water sampling	Environmental Modeling and Trend analysis – Watershed Stewardship Section.	Tidal and Nontidal waters of Indian River Bay, Rehoboth Bay and Little Assawoman Bay	Over 30 years of sampling. Frequency varies from monthly to quarterly depending upon Statewide needs of the	General Water chemistry, Field parameters, bacteria, and chlorophyll	Ongoing - stable	EPA via Watershed Stewardship

				overall Surface Water Monitoring Plan			
RT	Biology sampling (macroinvertebrates)	Status of environmental condition Environmental Laboratory Section	Nontidal waters of Indian River Bay, Rehoboth Bay, and Little Assawoman Bay	25 years of sampling Sporadic, present intent is to sample random sites within the watershed every 5 years	Macroinvertebrates, general water chemistry, field parameters, and chlorophyll	Sporadic - unstable	EPA via Watershed Stewardship
JY	Water column sampling	Research projects	Guinea Creek	1 year, sporadic	NO3, NH4, PO4, chlorophyll a	Project ends 1/31/16	DE Sea Grant/NOAA
JY	Ground water sampling	Research projects	Indian River Bay	2 years, sporadic	NO3, NH4, PO4, N isotopes	Project ended 1/31/15	DE Sea Grant/NOAA
SA	1) Stream and tide gaging	Ambient hydrologic monitoring	IRB, RB, LAB, tributaries	1980's - present	Tide height, stream discharge	Always a challenge to acquire funds	State and federal pass through sources
SA	2) Coastal flooding	DNERR, DEMA, DEOS, CIB	IRB, RB	2014 - 2016	Tide height	One and done	NERR
SA	3) WQ portal	DNREC	Statewide		WQ	?	DNREC
SA	4) Evapotranspiration	DEOS	RB watershed at Warrington Farm	2015-	Atmospheric Water flux	?	DNREC/DEOS/DGS

SA	5) Groundwater level and salinity	DNREC/DGS	Multiple wells	1980's - present	Water level, temperature, and salinity	?	DNREC/DGS
JF	UD Citizen Monitoring Program	DNREC, CIB, public	Inland Bays	Since 1991, year round, weekly in summer	DO, Secchi, salinity, nutrients, Chlorophyll a, and TSS	Continued nutrients is contingent on CIB or other funding support	DNREC
JF	Phytoplankton (HAB) monitoring	Public health, DNREC	Inland Bays	Since 2001	Screen and enumerate potentially harmful algal species	Continuing	DNREC
JF	Recreational Water (BEACH) monitoring	Public health, DNREC	Inland Bays	Since 2003	Total enterococcus	Continuing	DNREC/EPA
EW	Volunteer field data	Impairment in DO and water clarity/DNREC, CIB, and public	Inland Bays and tributaries	25 years	DO and Secchi depth	Ongoing	Funded/DNREC
EW	Nutrient samples	Impairment in nutrient, chlorophyll and TSS levels DNREC, CIB, volunteers, and UD researchers	Inland Bays and tributaries	20 years	Nitrate plus Nitrite Ammonium DIP Chlorophyll TSS	Ongoing, but cutting back on sites and outsourcing analysis	Funding for analyses on year to year basis/CIB

EW	Bacteria samples	Recreational contact safety/public, CIB, DNREC, and UD	Inland Bays and tributaries	11 years	Total Enterococcus	Ongoing, expect to continue	Funded/DNREC & EPA
EW	Harmful algae samples	AULS and public safety/DNREC, UD, public	Ocean beach, Inland Bays and Tributaries	14 years	HAB species, fresh and marine General phytoplankton community	Ongoing, expect to continue	Funding limited/DNREC & EPA

Table 2

Attendee	Program/Monitoring Activity	Purpose/users	Where? (bay(s), stream(s), watershed(s), etc.)	Duration, frequency of sampling	Parameters measured	Anticipated Start Date	Funding Status/ Sources
RG	Watershed Approach to Toxics Assessment and Restoration (WATAR)	See attached Work Plan	Watersheds with known toxics impacts (does not include the IBs)	One-time, comprehensive synoptic survey, with follow-up investigation if justified	PCBs, dioxins/furans, OC pesticides, PAHs, and mercury in surface water, sediment, and biota	Not scheduled within next 2 years	State & Federal
SA	DE Sea Grant proposal	Ullman, Andres	Groundwater project in RB watershed	2 yrs – multiple/day	Hydrology, nutrients, and DOC	Jan 2016	DE Sea Grant
SA	Groundwater salinity	DGS/DNREC	Multiple wells	ongoing	salinity	?	State of DE capital budget or see Table 1, SA reponse 5)

Delaware Center for the Inland Bays
Notes from Monitoring Plan Workgroup Meeting, 30 July 2015

Attendees:

CIB – Marianne Walch
RKK – Jim Eisenhardt, Larry Trout, Leslie Jamka
DNREC – Robin Tyler, David Wolanski, Michael Bott, Debbie Rouse, Hassan Mirsajadi, John Schneider
University of Delaware (UD) – Joanna York, Kevin Brinson, Tina Callahan, Ed Whereat, Bill Ullman, Joe Farrell, and Scott Andres
USGS – Judy Denver

Introduction

- Monitoring Plan for the Comprehensive Conservation and Management Plan (CCMP)
 - Measures effectiveness of CCMP
 - Written in 1995¹
 - Revised by Robin Tyler (DNREC) in 1996
 - Charged by EPA to update by 1 Oct 2015
 - 2012 CCMP addendum: New goals and strategies to be incorporated into the Monitoring Plan
 - Changes since 1996: New data, programs, partnerships, needs, technologies, understanding of the Inland Bays, TMDLs, etc.

CIB needs/goals

- Status and trends of the Inland Bays
- Identify data needs/gaps
- Identify/access best available data
- Coordination
- Public education/engagement/perception

Objectives of facilitated discussion

- Obtain input from partners
- Monitoring needs/goals
- Identify strengths/weaknesses/gaps
- How best to house/share/archive data
- Identify and prioritize funding needs/opportunities
- Maintain perspective of “importance”; everyone thinks their work is the most important

Parties that should be at the workshop, but are not represented

- Agriculture sector
 - Jennifer Volk, Environmental Quality Extension Specialist (invited)
 - Delaware Department of Agriculture (DDA)
 - Laura Torres, Delaware Nutrient Management Program
 - Laura Match

¹Per Robin, data well pre-dated 1995.

- Scott Blair
- Sussex County
 - Mike Izzo, County Engineer
 - Heather Sheridan, Director of Environmental Services
- EPA Region III
 - Mike Hoffman (invited)
 - Bill Richardson² (invited)
- DNREC Division of Fish and Wildlife
 - Initiate discussion/review of Monitoring Plan prior to 1 Oct vs. leave placeholders

History and status

- Intern updating datasets
 - Brian Glaser compiled/maintained list of historical studies/reports until 1996
 - Does CIB have this list?
- Folks in the room have considerable experience and long-term involvement
- EPA wants DE to take the lead in estuary management
- Big questions asked in 1996 and addressed with monitoring data
 - Eutrophication, habitat, and wetland loss
 - Dissolved oxygen, nutrients, chlorophyll, and pathogen indicator bacteria
- Continue to monitor for core data

Goals

- How best to dot the i's and cross the t's for EPA *and* obtain data important to DE?
 - Short-term: Submit revised Monitoring Plan to EPA by 1 October 2015³
 - What is needed to update the Monitoring Plan?
 - Use 2011 State of the Delaware Inland Bays as a starting point
 - Report trends; never say “we are there” – insinuates no need for funding
 - Highlight what is being done well and areas that are deficient or need improvement
 - Use the Monitoring Plan to introduce long-term needs and potential management resources
 - Long-term: What do we want to know? What are the big questions monitoring should answer? What needs to be monitored?
 - Continue current monitoring; expand to include new data, such as upper watershed
 - Identify action areas and short/medium/long-range goals
 - What data do we have/need?
 - Can we improve what we have?
 - Are there additional data that can be collected under existing monitoring?
 - Do we know target goals?

²Submitted questionnaire.

³Maintain the Monitoring Plan as a living document with opportunities to update.

- Do we understand system enough to know goals/needs?
- Different areas have different goals⁴
- Monitor water quality or water quality indicators?
 - What are other monitoring criteria?
 - What are other indicators of estuary health?
- Broader scope than just water quality: Need to capture chemical, physical, biological data
- Changing needs
 - As initial problems are addressed and the Inland Bays improve, other/secondary problems become apparent
 - Recognize evolutionary changes
 - Is monitoring capturing data?
 - If not, how best to capture?
 - Restoration is really renovation: Bays should look better, but not necessarily what they looked like in the past
 - Do we know what they looked like?
 - Is current level of monitoring sufficient to see/show changes?
- Big picture/think outside the box
 - Key concerns/players/milestones
 - Opportunity to really make a difference
 - Communicate information to future generations
 - Keep science going: Change the lingo, monitoring *is* the science
 - Effective mechanisms for data sharing and collaboration
 - Creative approaches to funding monitoring initiatives

Funding

- Issues
 - Identifying and obtaining funding for monitoring is difficult
 - EPA will not fund monitoring
 - Most states do limited monitoring
 - Need creative approach to fundraising
 - Science often done “EPA’s way” to standardize data/collection for statistical purposes
 - Change is difficult to see
 - Key: Create strategy to motivate change
 - Market the collaborative/collective approach to increase options/opportunities/success
 - CIB is hiring a water quality manager that could manage a grant
 - Leverage research/resources of others
 - Current funding for on-going activities
 - Funding is continuously decreasing with inflation
- Entities
 - Delaware is a small state; how best to market and secure funding?
 - Corporate sponsorships: Walmart, WWTPs, power plants, artesian water, etc.

⁴For example, seeing the stream bottom can be good, but in wetlands, water clarity is bad.

- Private parties including non-profits/foundations
- Kickstarter
- Strategies
 - Avoid using the word monitoring in proposals
 - “Sell” scientific question that can be answered by monitoring data
 - Clearly state why data are needed/utility of data
 - Partner vs. compete with the Chesapeake Bay

Questionnaire compilation/discussion

- How might the intensity, duration, and frequency of events drive the Monitoring Plan? How have these changed over time?
 - Twenty years ago, primarily spot monitoring (exception of pH and conductivity)
 - No option for continuous monitoring
 - Important questions to answer
 - Timescales needed to answer these questions
 - “Vat” of data
 - Need to synthesize and apply to answer bigger questions such as climate change
 - Are there things we should be monitoring, such as water depth?
- How are our actions impacting the Bays?
- Would we know improvement if we saw it?
- Answered lots of questions posed in 1995/1996
- What/where are the critical needs/trends that should be monitored?
 - Bacteria
 - EPA has guidance for the protection of recreational waters
 - CIB is concerned with health risks, which are becoming increasingly important
 - Tests are expensive, but people want these data
 - Look for pathogens vs. indicators
 - Sub-watersheds
 - Streams vs. larger bodies of water
 - Use local studies to inform larger questions
 - Stressors
 - Stressors are changing; point sources have decreased
 - Implications for monitoring
 - Monitoring may lead to identification of new management issues
 - Management practices
 - Example: Monitoring of BMPs
 - Privacy concerns, lack of focus, small scale
 - Need aggregated, doable, monitoring strategy
 - Lack before and after data
 - Know Inland Bays system now much better than in 1995
 - Lots of data
 - Third generation of modelling
 - Problem: Minimal, and/or anecdotal, historical data from the 1950s/60s
 - Look for trends in all applications

- Example: 305b reporting includes downstream monitoring that summarizes trends
- 1999 – 2013 trends
 - Nitrogen down, phosphorus up/down
 - Slow, but steady improvement
- New sources of contamination
 - Chemical indicators for small source monitoring
 - New technology for monitoring
 - Effect on aquatic health
- How to handle changes/trends that have yet to hit the Inland Bays
 - What are strong indicators of health in the Inland Bays?
 - Hard to quantify if no historical data
- Short vs. long-term monitoring
 - Limited utility of short-term monitoring; need long-term monitoring to see changes, which only manifest with time
 - Consider scale/frequency of sampling/monitoring
 - Intensive monitoring for a year vs. every five years
 - Advantages/necessity of more frequent monitoring
 - Monitoring indicators vs. trends
 - CIB monitoring interests may not match DNREC monitoring interests
 - National vs. state-specific focus
 - DNREC only has one station in the Inland Bays
 - Continue base monitoring of Inland Bays, but add more specific monitoring upland (sub-basins/watersheds)
- Groundwater
 - Is DNREC monitoring groundwater?
 - CCMP goal: Groundwater monitoring for saltwater intrusion
 - Good assessment tools for groundwater, but expensive and difficult
 - Note in Monitoring Plan
 - Lack clear understanding of land-based wastewater; some polluter-based monitoring
 - Target groundwater collection over time
 - Could do more with base flow sampling or mine existing data for flow
 - Need to understand processes and re-sample in networks not sampled recently
 - Jen Volk (UD) does continuous stormwater monitoring
 - Other sampling efforts
 - North East Water Resources Network (NEWNet)
 - Researchers in Rhode Island, Delaware, and Vermont are using sensors in streams to measure water depth, temperature, dissolved oxygen and organic matter, nutrients, and cloudiness
 - National Estuary Research Reserve
 - Network of 28 coastal sites designated to protect and study estuarine systems
 - NOAA funded; each site managed by state agency or university

- Maybe options to collaborate
- Data
 - Availability, accessibility, maintenance, integrity
 - Sharing mechanisms
 - STORET (STORage and RETrieval) data warehouse is EPA's repository for water quality, biological, and physical data
 - Available to state environmental agencies, EPA and other federal agencies, universities, private citizens, etc.
 - At this point, DNREC is the only one populating STORET
 - Historically, difficult to use
 - Greatest challenges
 - How/where to house data: STORET, Delaware Environmental Observing System (DEOS), other?
 - Accessibility is essential to long-term value/utility
 - Need solution for broader datasets
 - DEOS: Data aggregator of continuous data for Delaware
 - Provides interface
 - Mapping application for water quality data (pulls from STORET)
 - Other data considerations
 - Management
 - Who can manage? Maintain?
 - Need single entity to coordinate
 - Identify/include special/one-time studies
 - How to most effectively and efficiently share data?
 - Metadata to ensure longevity
 - How to avoid losing data/datasets?
 - How to maintain integrity?
 - How to handle studies with no digital data?
 - Utilization
 - Everyone is looking at their own data
 - How to aggregate for analysis?
 - No one is synthesizing, integrating, or compiling data; very time-intensive activity
 - How best to do this?
 - Who should/could do this?
 - CIB is only one of many users
 - Need common time stamp
 - Need universal format/standardization
 - How can CIB use data most effectively?
 - Collection
 - Define protocols in a specific way
 - Account for different collection strategies: Fixed sampling locations vs. collection within a box
 - Can existing datasets be tweaked to meet current data gaps/needs?
 - Needs
 - Continuous long-term datasets at fixed points

- Non-continuous periodic sampling, maybe with mobile sensors
 - Automated mechanism to report continuous data results vs. raw data
 - No need to sample pH in saltwater
 - Need minimum/maximum levels of dissolved oxygen
 - Are continuous concentrations of nitrogen and phosphorus necessary?
 - What is the necessity/utility of high density data streams?
 - Climate: Groundwater flow, depth, and inundation
 - Lacking data for acute/small-scale events
- Monitoring technology
 - Technology has changed drastically over the past 20 years
 - Increasingly cost-effective option to use portable sensors for single monitoring event or leave in place for extended monitoring
 - Time and cost savings: Instruments will defray analytical costs over time
 - Data available on the spot with no analytical needs
 - In two to five years, cost-effective option to purchase mobile equipment
 - Provides option for automated, continuous sampling
 - Disadvantages
 - Equipment needs to be maintained and calibrated
 - May not get all the data, such as enterococcus
 - Scott Andres is a co-PI for water quality sampling technology project
 - Need initial start-up funds to purchase equipment
 - DNREC has pool of equipment and experiences personnel within the state
 - Sensor capability
 - Some can collect temperature and salinity data needed for the hydrodynamic model
 - Cannot collect total nitrogen/phosphorus, but can collect nitrate
 - Could equipment be modified to fit need?
- Citizen Monitoring Data (CMD)
 - Questions of variability, quality, etc.
 - Volunteers are all trained and many have years of experience
 - CMD near shore stations vs. DNREC off shore stations
 - Shoreline data are heterogeneous; need large numbers to be meaningful
 - Huge volume of data with considerable buy-in/community support
 - Beneficial to add total nitrogen and total phosphorus
 - CIB wants to include CMD in Monitoring Plan; no one else is collecting these data
 - What can be done to increase credibility?
 - Example: Community Collaborative Rain, Hail and Snow Network (CocoRaHS)
 - How best to integrate?
 - Extensive QA/QC
 - Reports archived on website
 - Volunteer monitoring reports are a good example of data compilation
 - Should data be reviewed prior to posting online?

- Need to figure out the best way to get the data into STORET
 - These data are very important to the CIB for trends
- How to expand citizen monitoring activities?
 - How best to manage and oversee activities and data?
 - Is engagement an issue?
 - Can we request volunteers do specific things?
 - How best to expand capacity?
- Non-monitoring related needs
 - Searchable library/archive of historical reports, data summaries, etc.
 - Housed at CIB; does CIB have the capacity to maintain?
 - DNREC Watershed Assessment and Management Section moving; great opportunity to scan documents and get them online
 - Need someone to sort through historical data
 - Need summaries or keywords searchable in pdf image
 - Kent Price's student maintained list of reports/datasets until 1972ish
 - Who has this list? DNREC? CIB?
 - Accuracy
 - Not a critical issue unless data are used for regulatory purposes
 - Plot data over time/space; if consistent, accuracy is good
 - Stored electronically (STORET? If not, where?) and link to historical report
 - What is in it for CIB? Trends?
 - Consider compartmentalizing tasks for internships, etc.
 - Target specific sources

Questionnaire for CIB CCMP Appendix Monitoring Plan Update - General

Responders

- DJ: Doug Janiec, Sovereign Consulting Inc., Natural Resources Program Manager, Senior Restoration Ecologist
 - LT: Lauren Torres, Delaware Department of Agriculture, Nutrient Management Program
 - JD/MN: Judy Denver/Mark Nardi, USGS, MD-DE-DC Water Science Center
 - JC: John Clark, DNREC, Fish and Wildlife
 - Jim Sadowski
-

1. Did you participate in preparation/meetings/workshops for initial 1995/1996 CCMP Monitoring Plan?

- DJ: No.
- LT: No, I did not.
- JD: Yes, to some extent, but don't remember much about it.
- JC: No, I think Roy Miller and Jeff Tinsman participated for Fisheries.

2. What areas, based on your experience/expertise, are the most important to focus on in a monitoring plan for the Inland Bays and (very briefly) why?

- DJ: Habitat loss and gain (e.g., shoreline). Habitat loss is a top of pyramid indicator, which includes the net outcome of many primary indicators working together in a complex system.
- LT: Based on our experience, DDA believes that the most important areas to focus on in a monitoring plan for the Inland Bays are septic's, cesspools, and the expansion of sewer treatment plants based on the associated population growth in the Inland Bays watershed. DDA will continue to implement The Nutrient Management Program and the Concentrated Animal Feeding Operation (CAFO) Program to oversee nutrient handling activities associated with animal production, specifically poultry production, and on farm nutrient management practices.
- JD/MN:
 - Monitoring to understand water quality in the tidal portion of the bays for assessing current conditions and trends where samples have been collected long-term by DNREC and others should be continued to assess the health of the Bays.
 - The addition of upstream sites for long-term monitoring of non-tidal fresh surface-water quality would be very useful to track changes in water quality related to changes in land use and land management which could lead to a better understanding of the system as a whole. A monitoring network in small sub-watersheds is suggested. To track and understand nitrogen, which is mostly from groundwater, several sites that were selected and sampled in the late 1990s and early 2000s (by USGS, DGS and UD) could be used to develop a network of non-tidal sites and sampling could be repeated during winter base-flow conditions every year. A subset of these sites could be selected as a base

for this network and, and, at a minimum, could be sampled during winter baseflow for nitrogen inputs, but it would be better to do at least season sampling and include ecological sampling at least during the warm seasons. This network could also be used to track and better understand sediment and phosphorus inputs.

- JC: My experience/expertise is with fish, so I would focus on fish and water quality, but water quality and fish habitat depend on land use in the watershed, so I would have a hard time coming up with a single most important focus.
- JS: The Indian River Power Plant has shut down three cooling units and the remaining unit, with the cooling tower, has not been running very much. It was always said that the plant was impacting the Indian River and Bay but the 316a and 316b studies showed "no appreciable harm". I think the Center should further investigate what affects, if any, shutting down of the once through units has had on the Indian River and Bay. What has happened to all the species, increased/decreased/remained the same? Are the species changing? Are they different during the different seasons due to the temperature changes? What are the effects of the temperature changes? What is the effect of the reduced flow in Island Creek? The list can go on and on, but I think the Indian River System would be a great lab to see what affects a power plant shutting down has on the ecosystem.

3. What areas, based on your experience/expertise, are missing from the current monitoring plan?

- DJ: More detailed tracking on shoreline changes and its contribution to sediment loads in the Bays.
- LT: Based on our knowledge, we think it would be helpful for an evaluation of the total number of septic systems, and cesspools that are in existence in the Inland Bays Watershed, and how many of those are planned to be converted to the city sewer systems to be included in the new monitoring plan. Additional data could include the age and types of systems, and the estimated nutrient load from those sources, and subsequent reductions from conversions. Additionally, a summary of how this monitoring plan and all of these activities relate to the State of Delaware's Watershed Implementation Plan along with the TMDL's would help explain how all these activities tie together.
- JD/MN: See above answer for surface water.
- JC: Given that the Inland Bays has one of the healthiest hard clam populations of any coastal lagoon system on the Atlantic Coast, hard clam monitoring would be a good addition.

4. What monitoring activities/data collection is your organization undertaking in the Inland Bays that could be part of the CIB monitoring network?

- DJ: None, yet.
- LT: Currently, none, but see new project below.
- JD/MN:
 - Data are currently being collected at 7 stations in the Inland Bays watershed for flow, tide, and/or real-time monitoring of specific conductance, dissolved oxygen, pH, and temp.
 - The USGS has setup an ephemeral storm surge network (deployed only during events) comprised of 7 sites. Six of the sites measure water level and specific conductance; one of the sites is a rapid deployment gage (RDG) that measures and transmits in real time water level, wind speed and direction, barometric pressure, and air temperature. We will work with DeIDOT on RDG siting.

5. What monitoring programs are planned for the future that might be applicable to the Inland Bays?

- LT: DDA and USGS will be beginning a new study monitoring shallow ground water wells on a 5 year basis. See details in Table 2.
- JD/MN: Most of the current USGS monitoring in the Inland Bays is done in collaboration or cooperation with other agencies including DGS, DNREC and Del DOT and is likely to continue in the near future. Future plans depend on program directions of all agencies involved and are not known now.

6. For the data you discuss in this document, is your organization just collecting raw data or are you also processing the data and developing trends from the data?

- LT: The data is being collected currently to establish baselines, and the future data will be used to develop trends.
- JD/MN: All data currently collected by USGS are published and available on the USGS NWIS webpage under the “Atlantic Coastal Bays” tabs (http://waterdata.usgs.gov/de/nwis/current/?type=flow&group_key=basin_cd). In limited areas the USGS is working with cooperating agencies to analyze the data. All data collected by the USGS are subject to QA/QC and processing procedures prior to public release, except in limited cases where partner agencies have direct data access.
- JC: We process our trawl data and report trends in overall catch and trends in juvenile abundance indices.

7. What additional monitoring activities would your organization benefit from if CIB could include in their monitoring plan? Please feel free to think outside of the box, without regard to availability of funding, including new or innovative technologies.

- LT: Monitoring activities that would benefit the DDA, and all organizations, would be data that indicates the source of impairments and the age of the water.
- JC: The volunteer seining survey complements our trawl survey. I think it should be continued.

8. What do you, or your organization, feel are the critical indicators of health for the Inland Bays?

- DJ: Fisheries, shellfish, and SAVs, shorelines.
- LT: Critical indicators for the health of the Inland Bays are nutrient concentrations, dissolved oxygen, bacteria counts, submerged aquatic vegetation, and water clarity.
- JD/MN: SAV's, dissolved oxygen, water clarity, biological diversity. Underpinning all of these is water chemistry of both groundwater and surface water inputs which from a monitoring point of view are very important to the Inland Bays system that has limited exchange with the Atlantic.
- JC: Water quality, fish and other aquatic organism communities, hard clams, etc.

9. Does your organization have historical monitoring data that was not included in the original monitoring plan that would be useful for future monitoring activities or for use as State of the Bays indicators? Please list them.
- DJ: No.
 - LT: We do not have historical monitoring data.
10. Does your organization anticipate having sufficient future funding to carry out monitoring activities that CIB has included in the monitoring plan to date? If not, what level of additional funding is needed?
- LT: We do not have monitoring activities currently listed in the monitoring plan.
 - JC: Fisheries will continue the trawl survey.
11. Do you anticipate future funding for additional data collection parameters beyond the current monitoring criteria included in CIBs plan?
- LT: We anticipate the funding for the new project between USGS and DDA.
 - JC: No.
12. Do you have any recommendations for additional funding sources for Inland Bays monitoring activities?
- DJ: No.
 - LT: Not currently.
 - JC: No.
13. Are you aware of any volunteer organizations/programs that are currently active and could be included in future Inland Bays monitoring for supplemental information?
- DJ: No.
 - LT: I am not aware of any other organization or programs currently active in the Bays other than the Inland Bays, "Your Creek" projects.
 - JC: No.

Table 1

Responder	Program/Monitoring Activity	Purpose/users	Where? (bay(s), stream(s), watershed(s), etc.)	Years collected, frequency of sampling	Parameters/species measured	Current/future status	Funding Status/ Source
JC	Division of Fish and Wildlife Juvenile Finfish Trawl Survey	This survey estimates year class strength of various fish and invertebrate species and assesses abundance and community structure	12 stations sampled in Indian River and Indian River and Rehoboth Bays	1986 to present Sites are sampled monthly from April through October	Fish and invertebrates are sorted and counted by species A subsample of each species is measured Surface water quality is taken at each station	DFW plans to conduct this survey indefinitely	Secure funding through the Federal Aid in Sport Fish Restoration Program

Table 2

Responder	Program/Monitoring Activity	Purpose/users	Where? (bay(s), stream(s), watershed(s), etc.)	Duration, frequency of sampling	Parameters measured	Anticipated Start Date	Funding Status/ Source
DJ	More real-time monitoring, such as the York Canal Project		Near oyster cages and floating wetlands				
	Update the IB GIS AP and shoreline migration rates						
	Monitoring for project success/failure, not just bay-wide status/trends						
LT	USGS/DDA Shallow Ground Water Monitoring Program	Establish trends in nutrient levels in shallow ground water (40 ft or less) To test nutrient loading BMPs	Statewide, approximately 70 total wells (all dedicated established monitoring wells)	Every 5 years	N, P, age dating ions, land use tracer ions	Fall of 2014 & fall of 2015 to establish baselines First trend assessment will be Fall 2020	USGS and DDA matches
JC	Division of Fish and Wildlife hard clam density survey	Determine whether hard clam density is compatible with shellfish aquaculture on a given acre of bottom	Indian River and Rehoboth Bays	Sampling began in 2014 to determine hard clam density in acreage that DFW attempting to get permitted for shellfish aquaculture leases	Number of hard clams per square yard, bottom type	2014	DFW funds

Notes from STAC Meeting Discussion of Monitoring Plan September 18, 2015

Suggestion to review the monitoring plans of other programs:

- Other NEPs
- Chesapeake Bay Program
- National Coastal Assessment

Areas to focus on:

- *Bill Ullman*: Figure out how people use the estuary; focus on areas of the estuary that are most used
- *Robin Tyler*: Minimizing and tracking disruptive effects on the system
- *Jenn Volk*: Land use, BMPs, sea level rise, future land use
- *Andrew Homsey*: Tidal wetlands condition
 - Sea level rise, degradation, shorelines, vegetation changes
 - Aerial photography
 - Ability of wetlands to migrate
- Habitat Loss: GIS, remote sensing
- *Scott Andres*: Population, demographics and projections; this has a large impact
- Set aside areas to protect important resources
- *Joanna York*: Must consider projections to guide monitoring plan. Use models, think strategically about monitoring needed based on projections.
- Cost/benefit analyses (economics, ecosystem services). What has been spent already, what will be lost.
- Land surface loadings, changes (agriculture, wastewater treatment, development, etc.)
- *Robin Tyler*: Long-term seaweed monitoring – an important indicator of eutrophication. Last done in 2012. Suggest doing it every five years?
- Monitoring in South Bethany:
 - ~12 Citizen Monitoring Program stations there (salinity, temp, clarity, DO); three sites monitor bacteria and nutrients; three continuous monitors
 - The town collects and analyzes data
 - Local tide gages are monitored and analyzed (sea level rise)
- *Jim Sadowski*: There has been no follow-up on actions taken in the Bays (e.g. NRG plant changes, dredging) to determine if expected impacts have happened. Only short-term studies. We must know if management actions have worked.
- Monitoring to determine the impact of oyster farming.
- *John Clark*: DFW tracks fish kills (since 1981)
- Microbial source tracking
- Phytoplankton monitoring: biomass (chl A), trends, changes in community composition

Historical Data Sources: South Bethany WQ Committee, Robin Tyler and DNREC Environmental Laboratory Section, Division of Watershed Stewardship, Bill Ullman

Key to coordination is educating all partners on what others are doing. CIB needs to do this on an ongoing basis, through workshops, symposia, special STAC meetings, etc.

APPENDIX C

DESCRIPTIONS OF EXISTING LONG-TERM MONITORING PROGRAMS IN THE INLAND BAYS STUDY AREA

C.1	Monitoring of Surface Water	
C.1.1	State of Delaware Ambient Surface Water Quality Monitoring Program.....	C-2
C.1.2	University of Delaware Citizen Monitoring Program.....	C-7
C.1.3	State Fecal Coliform Monitoring Program	C-19
C.1.4	USGS-DGS Stream and Tide Gaging Program	C-19
C.1.5	State Biological Assessment of Streams Program	C-22
C.1.6	Toxics Monitoring	C-23
C.2	Monitoring of Groundwater	
C.2.1	Delaware Groundwater Monitoring Network.....	C-26
C.2.2	Delaware Agricultural Shallow Groundwater Monitoring Network	C-28
C.2.3	Public Supply Wells.....	C-30
C.3	Monitoring of Wastewater	
C.3.1	Point Source Wastewater Discharges	C-31
C.3.2	Land Application of Wastewater	C-32
C.4	Monitoring of Atmospheric Deposition	
C.4.1	Atmospheric Deposition of Nutrients.....	C-34
C.5	Monitoring of Wetlands	
C.5.1	DNREC Wetlands Assessment and Monitoring	C-35
C.5.2	Long-term Inland Bays Salt Marsh Monitoring.....	C-38
C.6	Monitoring of Living Resources and Habitats	
C.6.1	National Aquatic Resource Surveys	C-40
C.6.2	Statewide Vegetation Community and Land Cover Mapping Project	C-44
C.6.3	Seaweed Monitoring.....	C-45
C.6.4	Coastal Finfish Assessment Survey	C-46
C.6.5	Volunteer Inshore Fish and Blue Crab Survey	C-48
C.6.6	Recreational Fishing Surveys	C-49
C.6.7	Hard Clam Surveys	C-50
C.6.8	Volunteer Horseshoe Crab Survey and Tagging Program	C-52
C.6.9	Delaware Breeding Bird Atlas	C-53
C.6.10	Mid-winter Waterfowl Surveys	C-55
C.6.11	Bald Eagle and Osprey Nesting Surveys.....	C-56

C.1 MONITORING OF SURFACE WATER

C.1.1. STATE OF DELAWARE AMBIENT SURFACE WATER QUALITY MONITORING PROGRAM

Description/Objective(s)

The Delaware Department of Natural Resources and Environmental Control (DNREC) maintains a statewide General Assessment Monitoring Network (GAMN) of 134 stations, of which 24 are located in the Inland Bays (Figure C.1). GAMN stations are considered long-term stations, and data collected support compilation of Watershed Assessment Reports as mandated by the Clean Water Act under section 305(b).

All GAMN stations are monitored for temperature, salinity, dissolved oxygen, pH, alkalinity, hardness, chloride, chlorophyll, biological oxygen demand, total suspended solids, turbidity, organic carbon, nutrients, and Enterococcus bacteria. Tidal waters and lakes/ponds are also monitored where and when possible for water clarity (Secchi depth) and light attenuation. Some nontidal and tidal stations are further monitored for metals, while some nontidal stations are monitored for biology/habitat.

Monitoring frequency at GAMN stations follows a 5-year rotating basin schedule in which every station is monitored monthly for 2 years and monitored every other month for the remaining 3 years. Each station is monitored for conventional parameters such as nutrients, bacteria, dissolved oxygen, pH, alkalinity, and hardness (see Tables C.1 and C.2. for a complete list of parameters). Some stations are also monitored for dissolved metals.

Responsible Organization and Contact

The collection of ASWQMP samples and field data is conducted by the DNREC, Division of Water, Environmental Laboratory Section (ELS). The analysis of samples and generation of analytical results is also done by the ELS, with exception of some tests which are outsourced to selected laboratories that have EPA-approved Quality Assurance Management Plans.

Contact: Kathy Knowles, DNREC Environmental Laboratory Manager
kathy.knowles@state.de.us, 302-739-9942




FIGURE C.1 -- Map showing DNREC GAMN monitoring stations in the Inland Bays.

DNREC's Division of Watershed Stewardship, Watershed Assessment Section, analyzes the data to (1) define the water quality status and trends for each sub-basin and (2) compare the data with water quality standards to assess designated use support as mandated by Section 305(b) of the CWA.

Contact: David Wolanski
david.wolanski@state.de.us, 302-739-9939

Data Collection Cost

The annual cost of water quality monitoring of sites within the Inland Bays Watershed is between \$100,000 to \$170,000, depending on frequency of monitoring (monthly or every other month). Frequency of monitoring follows a 5-year state-wide rotating basin schedule. Per this schedule, during every 5- years, most monitoring sites in the Inland Bays are monitored monthly for 2 years and are monitored every other month for 3 years.

Data Use

GAMN data are used to:

- Describe general water quality conditions of the State's surface waters;
- Identify long term trends in water quality;
- Determine the suitability of Delaware waters for water supply, recreation, fish and aquatic life, and other uses;
- Calculate annual nutrient loads and track progress toward achieving Total Maximum Daily Loads (TMDLs) targets; and
- Evaluate the overall success of Delaware's water quality management efforts.

The findings are reported biannually to the EPA in the Water Quality Inventory Report as mandated by Section 305(b) of the Clean Water Act (CWA), and are used to identify and prioritize water-quality limited waters as mandated by Section 303(d) of the CWA, as well as waters of high quality. Since the late 1990's the data have been used to develop and calibrate TMDL models, and in the future will be used to gage the success of TMDL-based Pollution Control Strategies.

Data from this program also are used by the Center to prepare environmental indicator reports, including the pentennial *State of the Delaware Inland Bays* reports.

Record of Collection

See Table C.1 for the record of collection at each Inland Bays station.

Collection Method

Water sample collection:

All water sample collections are conducted by the DNREC Field Services Branch according to the Environmental Laboratory's *Operational Procedure, Surface Water Sampling*,

Continuous Monitoring:

A portable, automated on-site laboratory was deployed from 2005-2011 at the outlet to Millsboro Pond in order to define inorganic nitrogen and phosphorus loads entering the Inland Bays via the nontidal segment of Indian River.

DNREC has been developing a network of water quality monitoring stations at which data is collected continuously for dissolved oxygen concentration and percent saturation and other parameters (temperature, specific conductivity, pH, and salinity) that exhibit substantial fluctuation over short time scales such as the diel (24-hour) cycle or in response to weather conditions. This monitoring is conducted using YSI 6-series multi-parameter sondes. Measurements are taken at least every 15 minutes when the instruments are deployed. A rotating basin approach in the state is planned by DNREC.

Currently the only continuous monitoring station is deployed in Massey's Ditch (USGS 01484680), at the Massey's Landing fishing pier. Water quality data has been collected at this site since November 2011 (<https://waterdata.usgs.gov/usa/nwis/uv?01484680>). It is funded jointly by USGS and DNREC.

Analysis Methods

See Table C.2 for a summary of analytical methods used for each parameter.

Data Location

The GAMN data are entered into the STORET database, and are publically available via the Delaware Water Quality Portal (<http://demac.udel.edu/waterquality/>).

Management Goal

Management goals are defined by the TMDLs approved for each watershed.

Hypothesis and Test Statistics

Methodologies for analyses are defined in the *State of Delaware Draft 2016 Assessment, Listing and Reporting Methodologies Pursuant to Sections 303(d) and 305(b) of the Clean Water Act*, available from the Division of Watershed Stewardship.

TABLE C.1. -- Summary of DNREC GAMN Stations in the Inland Bays

Station ID	Location		Period of Record		Sampling Frequency*
	Latitude	Longitude	Start	Latest	
312011	38.557982	-75.089363	1998	2016	b
311041	38.455828	-75.206459	1998	2016	b
310121	38.521498	-75.133453	1998	2016	a
310071	38.478040	-75.055024	1998	2016	b
310031	38.483943	-75.117882	1998	2016	b
310011	38.453054	-75.064774	1998	2016	b
309041	38.564236	-75.274149	1998	2016	b
308371	38.721514	-75.206296	1999	2016	b
308361	38.545671	-75.163470	1999	2016	b
308341	38.622818	-75.256184	1998	2016	b
308281	38.638595	-75.316098	1998	2016	b
308091	38.547228	-75.243750	1998	2016	b
308071	38.594322	-75.290992	1998	2016	a
308051	38.640505	-75.173960	1998	2016	b
308031	38.671799	-75.185884	1998	2016	b
306341	38.583450	-75.224515	1998	2016	b
306331	38.587743	-75.204563	1998	2016	b
306321	38.609540	-75.067228	1998	2016	a
306181	38.590102	-75.248384	1998	2016	b
306121	38.606510	-75.111310	1998	2016	a
306111	38.625578	-75.099793	1998	2016	b
306091	38.649121	-75.109355	1998	2016	b
305041	38.772986	-75.132947	1998	2016	b
305011	38.708891	-75.093022	1998	2016	b

* a: Station is monitored monthly every year.

b: Station is monitored six times a year for three years, then twelve times a year for two years.

TABLE C.2 -- Water quality parameters analyzed at DNREC GAMN Stations and analytical methods.

Parameter	Method Reference (EPA)	Reporting Level
Water Column Nutrients		
Total Phosphorous	EPA365.1 M	0.005 mg/l P
Soluble Ortho-phosphorus	EPA365.1	0.005 mg/l P
Ammonia Nitrogen	EPA350.1	0.005 mg/l N
Nitrite+Nitrate N	EPA353.3	0.005 mg/l N
Total N*	SM 4500 NC	0.08 mg/l N
Carbon and Organics		
Total Organic Carbon	EPA415.1	1 mg/l
Dissolved Organic Carbon	EPA415.1	1 mg/l
Chlorophyll-a (Corr)	EPA 445.0	1 µ/l
Biochemical Oxygen Demand		
BOD ₅ , N-Inhib (CBOD)	SM20 th ed-5210B	2.4 mg/l
BOD ₂₀ , N-Inhib (CBOD)	SM20 th ed-5210B	2.4 mg/l
General		
Dissolved oxygen – Winkler	EPA360.2	0.25 mg/l
Dissolved oxygen – Field	EPA360.1	0.1 mg/l
Total Suspended Solids	EPA160.2	2 mg/l
Alkalinity	EPA310.1	1 mg/l
Hardness	EPA310.2	5 mg/l
Field pH	EPA150.1	0.2 pH units
Conductivity - Field	EPA120.1	1 µS/cm
Salinity	SM20 th ed-2520B	1 ppt
Temperature	EPA170.1	°C
Secchi Depth	EPA/620/R-01/003	meters
Light Attenuation	EPA/620/R-01/003	%
Turbidity	EPA180.1	1 NTU
Chloride	EPA325.2	1 mg/l
Bacteria		
Enterococcus	SM20 th ed-9230C	1 cfu/100 ml

3.1.2 UNIVERSITY OF DELAWARE CITIZEN MONITORING PROGRAM

Description/Objective(s)

University of Delaware Sea Grant Marine Advisory Service's Citizen Monitoring Program (CMP) was formed in 1991 to support the newly-formed Inland Bays Estuary Program (<http://citizen-monitoring.udel.edu/>). Through this citizen science program, more than 300 trained volunteers have collected samples at water quality monitoring sites throughout the Inland Bays, and provided important data - including dissolved oxygen, dissolved inorganic nitrogen, dissolved inorganic phosphorus, water clarity, bacteria levels, and other environmental data. Supplemental NEP grants have helped grow shorter-term special interest monitoring programs, including harmful algal species and dissolved oxygen measurements taken from boats.

Combined with the state's fixed monitoring stations, the additional resolution provided by the more than 15 years of data, from over 30 sampling sites in the Inland Bays, has resulted in a long-term, robust, high-quality data set that provides scientists and resource managers with a clearer picture of the bays' health and the trend information needed to understand and manage the ecosystem.

Responsible Organization and Contact

The program is managed by the University of Delaware Sea Grant Marine Advisory Service at the College of Earth, Ocean, and Environment in Lewes.

Contact: Edward Whereat
whereat@udel.edu, 302-645-4252

Data Collection Cost

Original support for the CMP came through the National Estuary Program, but since 1994 the Citizen Monitoring Program has received an appropriation from the Delaware General Assembly through a Memorandum of Agreement with DNREC. Additional support currently comes from a variety of sources, including: Sea Grant; DNREC; the Center for the Inland Bays; U.S. EPA; the Delaware Estuary Program; the National Fish and Wildlife Foundation; and local communities.

Reported total cost of the program in federal fiscal year 2016 was \$102,737. This included \$34,177 federal funding, \$45,000 state funding, and \$23,560 private in-kind.

In 2015 and 2016, the CIB has covered the cost for nutrient analyses to be conducted by the University of Maryland's Chesapeake Biological Laboratory. Total annual cost for these analyses is approximately \$4,000.

Data Use

CMP data are used by the CIB for status and trend analyses in the CIB's 'State of the Inland Bays' reports, which are published every five years. In addition, CMP data are being used to develop indicator reports for individual tributaries in the CIB's 'Your Creek' project.

Bacteria and harmful algal bloom data collected by the CMP are used by DNREC to support the Delaware Shellfish Monitoring Program and to assess bacteria levels and trends in the state's coastal watersheds.

Record of Collection

See Table C.3 for the record of collection at each Inland Bays station.

Collection Method

Data are collected by trained volunteers at sites assigned to them by the CMP Program Manager. Table C.3 summarizes the CMP monitoring sites that currently are used to assess Inland Bays water quality status and trends. Figures C.2 to C.5 provide a maps of CMP sites for individual indicator parameters. Additional sites are monitored in the Bays, but only those that provide long-term data used for Center environmental indicator reports are included here.

Sampling methods are detailed in a guidebook, *Inland Bays Citizen Monitoring Program Volunteer's Water Quality Monitoring Manual* (revised 2017). They are summarized below:

<u>Parameter</u>	<u>Collection Frequency</u>	<u>Method/Instrument</u>
Temperature	Weekly	Thermometer
Turbidity	Weekly	Secchi Disk
Dissolved Oxygen	Weekly	Micro-Winkler Titration
Salinity	Weekly	Hydrometer
pH	Weekly	Digital Meter
Rainfall	Daily	Rain Gauge
Nitrates	Biweekly	Grab Sample/Lab Analysis
Orthophosphates	Biweekly	Grab Sample/Lab Analysis
Bacteria	Biweekly	Grab Sample/Lab Analysis
Phytoplankton	Biweekly	Grab Sample/Lab Analysis

Analysis Methods

Currently, nutrient analyses are subcontracted to the University of Maryland Chesapeake Biological Laboratory in Solomons, MD. Other laboratory analyses are conducted in-house at the University of Delaware, with assistance from trained volunteers.

See Table C.4 for a summary of analytical methods used for each parameter.

Note that DIP is labeled as Orthophosphate in the state's GAMN dataset. DIN is calculated by summing nitrate+nitrite and dissolved ammonia.

Data Location

Data are stored in a database at the University of Delaware College of Earth, Ocean, and Environment in Lewes. Semi-monthly reports are posted online at <http://www.citizen-monitoring.udel.edu/reports/>. Currently the data cannot be queried or downloaded online. Data are available by request to the CMP Program Manager.

Management Goal/Hypothesis and Test Statistics

Data are included as an Appendix to the state's Combined 305(b) Report and 303(d) List, but are not used directly in listing determinations.

The CIB uses CMP data to develop an indicator known as the Water Quality Index (WQI) for eelgrass reestablishment. The WQI is created by relating the values of DIN, DIP, Chlorophyll a, and Secchi depth to the eelgrass restoration criteria developed in the Chesapeake Bay as refined for the Delmarva Coastal Bays (Valdes-Murtha 1997, Batuik et al. 2000). Thresholds are DIN = 0.14 mg/L, DIP = 0.01 mg/L, Chlorophyll a = 15 micrograms per liter, and Secchi depth = 2.2 ft.

All data collected for these parameters that followed the established criteria for inclusion are scaled linearly using the following Excel function:

$$=TREND(calcs!E$2:F$2,calcs!C$4:D$4,F2)$$

E2 and F2 represent the top and bottom of the scale desired (0 and 1), and C and D represent the 95th percentile of all data for that parameter and the threshold value for that parameter. F represents the actual parameter value. For Secchi depth, the 5th percentile is used instead of the 95th percentile, because a larger number represents a better Secchi depth, unlike the other parameters where a larger value indicates a worse environmental condition. After scaling linearly, any values below 0 are changed to 0, and any values exceeding 1 are changed to 1. Then, for each row, the four parameters are added

together, and the sum was divided by 4 to get a mean WQI for each sampling event at each station. The mean annual WQI values are used for Mann Kendall trend analyses.

Indicator	Criteria for Indicator Status			
	Very poor water quality	Does not support eelgrass reestablishment	May support eelgrass reestablishment	Supports eelgrass reestablishment
Water Quality Index for Eelgrass Reestablishment (range=0 to 1)	<0.75	0.75 to <0.9	0.9 to <1.0	1.0

TABLE C.3 – Summary of Citizen Monitoring Program Stations in the Inland Bays. These are stations currently sampled regularly, as of 2016.

Notes: ¹ Stations identified by the CIB as top priority to continue, for use in the State of the Delaware Inland Bays reporting.

² Station was sampled fewer than five times for any parameter in 2015.

Station ID	Location		Period of Collection		Parameters Collected									
	Latitude	Longitude	Start Date	End Date	Chl a	DIN	DIP	DO	pH	Salinity	Secchi Depth	Total <i>Enterococcus</i>	TSS	Water Temp
IR02 ^{1,2}	38.59453	-75.20838	2000	2002										
IR04 ¹	38.59583	-75.20500	2000	2015	•	•	•	•		•	•	•	•	•
IR11	38.61683	-75.10223	2000	2015	•	•	•	•		•	•	•	•	•
IR12	38.58853	-75.07288	2001	2015				•		•	•			•
IR20 ¹	38.58260	-75.15509	2000	2015		•	•							
IR24	38.58212	-75.26920	2013	2015				•		•	•	•		•
IR26	38.58323	-75.26492	2015	2015				•		•	•			•
IR29	38.57123	-75.08472	2005	2015				•		•	•			•
IR32	38.57069	-75.08449	2002	2015				•		•	•	•		•
IR36	38.58125	-75.08710	2004	2015				•		•		•		•
IR38 ¹	38.56158	-75.19947	2006	2015	•	•	•	•	•	•	•	•	•	•
IR39	38.60893	-75.06465	2006	2015				•		•	•			•
IR50	38.55575	-75.08852	2002	2015				•		•	•			•
IR73	38.54901	-75.07046	2009	2015				•		•	•			•
LA03 ¹	38.48538	-75.07735	2000	2015	•	•	•	•		•	•	•	•	•
LA09 ¹	38.48362	-75.11880	2000	2015	•	•	•	•		•	•	•	•	•
LA29	38.47675	-75.13791	2015	2015				•		•	•	•		•
LA45 ^{1,2}	38.46103	-75.05778	2004	2013										
LA46 ¹	38.45497	-75.05833	2010	2015	•	•	•	•		•		•	•	•
LA48	38.46199	-75.05355	2011	2015				•		•	•	•		•
RB06 ¹	38.63828	-75.16918	2002	2015	•	•	•	•	•	•	•	•	•	•
RB06A	38.64065	-75.17410	2002	2015						•		•		•
RB07 ¹	38.66194	-75.13222	2000	2015		•	•							
RB34 ¹	38.70262	-75.16075	2002	2015	•	•	•	•		•	•	•	•	•

Station ID	Location		Period of Collection		Parameters Collected									
	Latitude	Longitude	Start Date	End Date	Chl a	DIN	DIP	DO	pH	Salinity	Secchi Depth	Total <i>Enterococcus</i>	TSS	Water Temp
RB38	38.67794	-75.13603	2000	2015						•				
RB64	38.69909	-75.11242	2005	2015				•		•				•
RB80	38.71046	-75.17502	2014	2015				•		•		•		•
RB90	38.66555	-75.18235	2014	2015				•	•	•		•		•
SB01	38.51901	-75.05558	2008	2015	•	•	•	•		•	•	•	•	•
SB02	38.51851	-75.06083	2006	2015				•		•	•			•
SB04	38.51793	-75.05603	2007	2015				•		•	•	•		•
SB05	38.51783	-75.06050	2009	2015				•		•	•			•
SB06	38.51744	-75.05539	2012	2015				•		•	•	•		•
SB07	38.51490	-75.05993	2006	2015	•	•	•	•		•	•	•	•	•
SB09	38.51033	-75.05633	2008	2015				•		•	•			•
SB10E	38.52083	-75.06113	2002	2015				•		•	•			•
SB10W	38.52057	-75.06350	2002	2015				•		•	•			•
SB12	38.51356	-75.06233	2002	2015				•		•	•			•

TABLE C.4 – Summary of Citizen Monitoring Program Analytical Methods

Parameter	Method Reference (EPA)	Reporting/Quantitation Limit
Water Column Nutrients – These analyses are currently performed by the UMCES CBL NASL		
Soluble Ortho-phosphorous	EPA365.1	0.0025 mg/l P
Ammonia Nitrogen	EPA350.2	0.01 mg/l N
Nitrite + Nitrate N	EPA353.2	0.0035 mg/l N
General		
Dissolved oxygen – Winkler	EPA360.2	0.25 mg/l
Dissolved oxygen – Field	EPA360.1	0.1 mg/l
Chlorophyll-a	EPA445.0	1 µg/L
Total Suspended Solids	EPA160.2	2 mg/l
Field pH	EPA150.1	0.2 pH units
Conductivity - Field	EPA120.1	1 µS/cm
Salinity	SM20 th ed-2520B and 2520C	1 ppt
Temperature	EPA170.1	0.5 °C
Secchi Depth	EPA/620/R-01/003	0.1 meters
Bacteria		
Total Enterococcus	SM20 th ed-9230C	1 cfu/100 ml



FIGURE C.2 – Map showing locations of Citizen Monitoring Program stations that collect nutrient data.




FIGURE C.3 – Map showing locations of Citizen Monitoring Program stations that collect dissolved oxygen data.



FIGURE C.4 – Map showing locations of Citizen Monitoring Program stations that collect Secchi depth data.



FIGURE C.5 – Map showing locations of Citizen Monitoring Program stations that collect Total *Enterococcus* data.

C.1.3 STATE FECAL COLIFORM MONITORING PROGRAM

Description/Objective(s)

In order to regulate shellfish harvest areas, the State has monitored levels of total coliform bacteria and, beginning in 2016, fecal coliforms in all three Inland Bays. The goal of this program is to ensure that waters that are approved for shellfish harvest meet water quality standards based on routine water quality assessments. Additionally, this program can be used to justify the opening of previously closed areas for harvest should the data reflect the area is safe, and close previously open areas in the event of high coliform levels.

Responsible Organization and Contact

DNREC Division of Watershed Stewardship

Contact: Michael Bott, Environmental Scientist
Michael.Bott@state.de.us, 302-739-9939

Data Collection Cost

Not available.

Data Use

Data are used primarily by DNREC Division of Watershed Stewardship's Shellfish Program to assess the suitability of the Inland Bays waters for shellfish harvest.

Record of Collection

2008 to present. Fecal coliform collection began in 2016, with plans to succeed total coliforms as the method to assess the suitability of waters.

Collection Method

Ten times per year, 52 sites in Rehoboth and Indian River Bay are sampled and analyzed for fecal coliforms. Little Assawoman Bay is sampled nine times a year at nine sites.

Analysis Methods

Analyses use mTEC agar (M-198) membrane filter medium, for enumerating fecal coliforms in marine and estuarine waters (FDA, 1998).

Data Location

Data are managed and stored by the DNREC Division of Watershed Stewardship Shellfish Program

Management Goal

Standard for total coliforms: no more than 10% or 90th percentile, of past 30 samples exceed 330 mpn/100mL; geometric mean shall not exceed 70mpn/100mL over the past 30 samples.

C.1.4 USGS-DGS STREAM AND TIDE GAGING PROGRAM

Description/Objective(s)

The US Geological Survey (USGS), in cooperation with the Delaware Geological Survey (DGS) through a State-Federal partnership program, operates and maintains stream and tide gages throughout Delaware. The stream gage network is a component of the National Streamflow Information Program (NSIP), a program that provides real-time and long-term current and historical streamflow information that is not only accurate and unbiased, but also meets the needs of many users.

Currently there are nine station locations in the Inland Bays; three of these measure flow (Table C.5 and Figure C.6).

Responsible Organization and Contact

The stations located in the Inland Bays are managed by the MD-DE-DC Water Science Center, 1289 McD Drive, Dover, DE 19901. Phone 302-734-2506

Data Collection Cost

Funding for the three drainage stations is provided by DNREC through the DGS. Funding for the operation of the Inland Bays tidal stations are funded by DNREC through the DGS, and the Delaware Department of Transportation. The USGS provides match funding for the streamgage portion of the program.

Data Use

The Delaware Stream and Tide Gage network provides the hydrologic and water quality information necessary to aid in defining, using, and managing surface and groundwater resources. The data are used for a multitude of purposes, including, but not limited to, long-range water resources planning and management, short-term resource management, evaluation of drought-no drought conditions, allocation of water resources for public, industrial, commercial, and irrigation water supplies, flood forecasting and warning, bridge and culvert design, hazard spill response and mitigation, analysis of sea level rise, recreation, and floodplain mapping. The stream and tide data are also utilized in existing real-time early warning systems related to potential flooding, and storm/coastal erosion throughout Delaware. The warning systems are used by the DGS, Delaware Emergency Management Agency, all three county emergency management offices, most municipalities, the National Weather Service, the Office of the State Climatologist, and others.

Record of Collection

See Table C.5.

Collection Method

Water-stage recorder gages. The peak tidal stage that is recorded by each gage is the elevation of water above the North American Vertical Datum of 1988. Stations have a USGS satellite data-collection platform.




FIGURE C.6 – Map showing locations of Tide Gauges in the Inland Bays.

Analysis Methods

Once a complete day of readings are received from a site, daily summary data are generated and made available online. USGS finalizes data at individual sites on a continuous basis as environmental conditions and hydrologic characteristics permit.

Data Location

Stream and tide gage information are available at <http://waterdata.usgs.gov/de/nwis/current/?type=flow>. Data from USGS stream and tide gage networks in Delaware are also available through the Delaware Environmental Observing System (DEOS) site: <http://www.deos.udel.edu/data/>.

TABLE C.5 – Inland Bays Stream and Tide Gage Stations

Station No.	Station Name	Latitude	Longitude	HUC-8	Parameters	Period of Record
01484525	Millsboro Pond Outlet	38°35'40.4"	75°17'27.7"	02060010	Discharge, Gage height	May 1986 to September 1988. March 1991 to current year.
1484695	Beaverdam Ditch near Millville	38°31'17.2"	75°08'00.2"	02040303	Discharge, Gage height	August 1998 to current year.
01484540	Indian River at Rosedale Beach	38°35'29.5"	75°12'41.7"	02040303	Discharge, Gage height	April 1991 to current year.
01484549	Vines Creek near Dagsboro	38°33'23.0"	75°12'11.4"	02040303	Tide elevation	Annual maximum, water years 1985-97. May 2015 to current year.
01484670	Rehoboth Bay at Dewey Beach	38°41'39.2"	75°05'03.2"	02040303	Gage height, tide elevation	April 1985 to September 1997; November 2000 to current year.
01484683	Indian River Bay Inlet near Bethany Beach	38°36'35.4"	75°04'04.8"	02040303	Gage height, tide elevation	June 1988 to June 1989, April 1991 to December 2010, November 2011 to current year.
01484690	Unnamed Ditch on Fred Hudson Rd at Bethany Beach	38°33'19.6"	75°03'48.4"	02040303	Tide elevation	May 2015 to current year.
01484696	Jefferson Creek at South Bethany	38°30'48.8"	75°03'44.4"	02040303	Gage height, tide elevation	July 1999 to current year.
01484701	Little Assawoman Bay at Fenwick Island	38°27'17.9"	75°03'30.0"	02040303	Gage height, tide elevation	October 1999 to current year.

C.1.5 STATE BIOLOGICAL ASSESSMENT OF STREAMS PROGRAM

Description/Objective(s)

This long-term project collects biological and habitat data from nontidal wadable streams in order to relate water quality conditions with biological integrity. Eligible stations must (1) be completely nontidal, (2) have perennial flow, and (3) be uninfluenced by elevated temperature resulting from lentic discharge (i.e. millpond, stormwater pond, etc). The biological data consists of two instream matrices: macroinvertebrate, and periphyton (first initiated in spring - 2005). The habitat data consists of instream and riparian zone matrices.

Beginning in 2006, and every other autumn thereafter, biological (macroinvertebrate), habitat and chemical sampling has occurred under baseflow conditions at 50 stations located along streams that have been placed on the 303(d) list due to impaired biology or habitat. This bi-annual sampling rotates by county, major basin, or both. Conjunctional chemical sampling will be conducted. The data are evaluated to determine whether any form of impairment still exists at each respective station. If impairment is concluded, then effort will be made to identify the cause/s. The procedure for identifying causation is not yet fully outlined but will likely follow the EPA Stressor Identification Guidance Document (Cormier et al, 2000).

In years between 303(d) sampling, up to 50 GAMN stations are be sampled. The biological and habitat methodology is the same as used for the 303(d) sampling.

The immediate objective of this sampling is to determine the overall biological condition of nontidal streams in Delaware. The extended objective is to identify trends in biological condition in these waters.

Responsible Organization and Contact

DNREC Division of Water, Environmental Laboratory Section

Contact: Kathy Knowles, Laboratory Manager
Kathy.knowles@state.de.us, 302-739-9942

Data Collection Cost

Not available.

Data Use

All analytical results are provided to the Watershed Assessment Section (WAS). The biological assessment data are used to:

- Define current water quality conditions.
- Identify and define long-term trends in water quality.
- Determine the suitability of Delaware waters for designated uses (e.g. water supply; recreation; fish, aquatic life and wildlife) as specified in the Delaware Surface Water Quality Standards.
- Determine whether the water quality standards are being met.
- Identify and prioritize high quality and degraded waters.
- Support the Total Maximum Daily Load Program.
- Evaluate the overall success of Delaware's water quality management efforts.

The 305(b) Report is submitted biannually by the WAS to the EPA as mandated by the Clean Water Act (CWA), and the findings are used to identify and prioritize water-quality limited waters as mandated by Section 303(d) of the CWA, as well as waters of high quality. Since the late 1990's the data have been used to develop and calibrate TMDL models, and in the future will be used to gage the success of TMDL-based Pollution Control Strategies.

Record of Collection

2000 to present

Collection Method

Biology and habitat sampling is done in accordance with methods defined in USEPA (1997). Biology samples are collected at coastal plain sites using a D-framed net.

Periphyton sampling is conducted according to the USGS, National Water Quality Assessment Program (Moulton et al. 2002). Samples will be collected from natural substrates, sticks and/or macrophytes (coastal plain), and rocks (piedmont). Only sticks that have obviously been in the water for an extended period (weeks to months) will be sampled.

Analysis Methods

The field preserved macroinvertebrate samples are outsourced for subsampling and identification to the lowest practical taxon. Level of identification for each phylum is as follows;

- *Arthropoda* genus / some species
- *Annelida* genus / some species
- *Mollusca* genus / some species
- *Bryozoa* family / some genus (statoblasts)
- *Platyhelminthes* genus / some species
- *Cnidaria* genus

For analytical purposes, the species composition and abundance data will be reduced to the genus level. A multi-metric approach will be used to calculate a biological index (BI) for each sample which is expressed as a percentage of the ecoregion reference values (see Gibson 1996). Based on the BI, the site will then be categorized according to condition (i.e. excellent, good, moderately degraded, severely degraded).

The periphyton samples also are outsourced for identification.

Data Location

All completed field-generated and laboratory-generated data are entered into the DNREC ELS Laboratory Information Management System (LIMS).

Management Goal

The data are evaluated to determine whether any form of impairment still exists at each respective station. If impairment is concluded, then effort will be made to identify the cause/s. The procedure for identifying causation is not yet fully outlined but will likely follow the EPA Stressor Identification Guidance Document (USEPA, 2000).

Hypothesis and Test Statistics

From each set of triplicate results (three stations sampled in triplicate for macroinvertebrates and periphyton) a coefficient of variation is developed. The range of these three coefficients of variation is regarded as the within-station spatial variability of the biological community across the entire study area.

C.1.6 TOXICS MONITORING

Ambient Water Quality Monitoring, Heavy Metals

Most of the state's monitoring of toxics has focused on the Delaware Estuary basin. Ongoing monitoring of toxics in the Inland Bays includes sampling and analysis of copper, lead, zinc, and arsenic as part of the state's Ambient Surface Water Quality Monitoring Program (Table C.6). This table is current as of FY 2016. Other more intensive monitoring and assessment of toxics in water, sediment and biota in the Inland Bays has occurred in the past, but these programs have not been continued long-term.

TABLE C.6 – State monitoring of toxics in surface water the Inland Bays.

Station Location	Station ID	Cu, Pb & Zn	As
Burton Pond @ Rt. 24	308031	●	
Millsboro Pond @ Rt. 24	308071	●	
Pepper Creek @ Rt. 26 (Main St.)	308091	●	
Blackwater Creek @ Omar Rd. (Rd. 54)	308361	●	
Dirickson Creek @ Old Mill Bridge Rd. (Rd. 381)	310031	●	
Buntings Branch @ Rt. 54 (Polly Branch Rd.)	311041	●	
Guinea Creek @ Banks Rd. (Rd. 298)	308051	●	
Whartons Branch @ Rt. 20 (Dagsboro Rd.)	309041	●	
Lewes & Rehoboth Canal @ Rt. 9	305041	●	
Little Assawoman Bay @ Rt. 54 (The Ditch)	310011	●	
White Creek @ mouth of Assawoman Canal	312011	●	
Bundicks Branch @ Rt. 23	308371	●	
Beaver Dam Ditch @ Beaver Dam Rd. (Rd. 368)	310121	●	
Cow Bridge Branch @ Zoar Rd. (Rd. 48)	308281	●	
Swan Creek @ Mount Joy Rd. (Rd. 297)	308341	●	
Lewes & Rehoboth Canal @ Rt. 1	305011	●	
Indian River Inlet @ Coast Guard Station	306321	●	●
Rehoboth Bay @ Buoy 7	306091	●	●
Masseys Ditch @ Buoy 17	306111	●	●
Indian River Bay @ Buoy 20	306121	●	●
Indian River @ Buoy 49 (Swan Creek)	306181	●	●
Indian River @ Island Creek	306331	●	●
Island Creek upper third	306341	●	●
Little Assawoman Bay Mid-bay (Ocean Park Lane)	310071	●	

EPA Environmental Monitoring and Assessment Program

The EPA has some data available on toxics for fish samples and sediment samples from the Inland Bays. Those samples were collected as part of the Environmental Monitoring and Assessment Program (EMAP), which was a research program run by EPA's Office of Research and Development to develop the tools necessary to monitor and assess the status and trends of national ecological resources. EMAP collected field data from 1990 to 2006. Data are available online (<https://archive.epa.gov/emap/archive-emap/web/html/index-37.html>). Monitoring of the nation's aquatic resources is now being routinely conducted by the National Aquatic Resource Surveys, run by EPA's Office of Water (<https://www.epa.gov/national-aquatic-resource-surveys>). More information on this program is available in this appendix (Section C.1.6).

Monitoring of Burton Island Coal Ash Disposal Area

The Center collected biota and sediment samples from the Inland Bays in 2012 to assess whether material eroding off and/or transported from the Burton Island former coal ash disposal site in upper Indian River is contributing to significant accumulation of toxic trace elements in the local aquatic environment.

Mummichogs (*Fundulus heteroclitus*), ribbed mussels (*Geukensia demissa*), and surface sediment were collected from five locations along the southern shoreline of Burton Island within Island Creek. Sites were purposely selected to coincide with locations where previous sampling or observations indicated release from Burton Island through either erosion or shallow groundwater transport. Locations within Pepper Creek were sampled for sediment and biota to serve as controls against which the Burton Island results could be compared.

The study concluded that existing conditions and concentration levels of trace elements found in the *Geukensia*, *Fundulus*, and sediment samples currently did not warrant an expansion of sampling to evaluate the potential ecological impacts of bioaccumulation (Riedel and Wilson, 2013). The future conditions of the island could change due to rising water levels and/or changes in the rate of pore water movement; because of this, it is recommended that tissue and sediment samples be periodically sampled and analyzed (in methods consistent with this study) to evaluate any changes in the prevalence and concentration of trace elements and metals through bioaccumulation in the surrounding biota.

A Long-Term Stewardship (LTS) Plan for the ash disposal area was prepared for Indian River Power LLC in 2014 (CB&I Environmental and Infrastructure, Inc., 2014). The LTS Plan includes monitoring requirements for the remedial actions at the site. Field and laboratory data are to be provided electronically to DNREC-SIRS. The monitoring requirements are:

- Annual groundwater monitoring to assess any changing physical conditions from those used to develop the site conceptual model of groundwater flow and to assess any changing chemical conditions used to evaluate the potential for human health and ecological risk.
- Annual monitoring of sediment quality immediately offshore of the site. Shoreline sediment sampling events will consist of a visual shoreline survey near the time of low tide, sample collection and documentation, and laboratory analysis. Shoreline sediment samples will be analyzed for arsenic, barium, and selenium as these metals were identified as sediment constituents of concern (COCs) in the risk assessment (Shaw, 2008).

Requests from the Center for sampling of additional heavy metals in groundwater and sediment, as well as for biological sampling, were not included in the Plan. Nor were numeric criteria based on ecological standards for all constituents of concern, or what might constitute an increase from baseline conditions in these parameters that would require any further remedial action.

Fish and Shellfish Tissue Monitoring

Statewide fish tissue monitoring has been conducted since 1992, and data have resulted in issuance of consumption advisories for Striped Bass and Bluefish caught in the Inland Bays. DNREC currently is conducting a large study of contaminant levels in Bluefish and Striped bass that may result in an adjustment to the coastal advisory for these species. Data collection will not be complete until sometime later in 2017.

DNREC conducts three types of monitoring in support of its chemical contaminants in fish and shellfish program. Tier I screening samples are collected in areas where data gaps exist, or where the existing data are very old. Tier II intensive samples are collected in situations where the results of Tier I monitoring indicate a need to better characterize the extent and magnitude of contamination (and support a risk assessment/advisory decision); and last, advisory follow-up sampling after an advisory has been issued and there's a need to track changes over time. DNREC has performed both Tier I and Tier II sampling of contaminants in fish and shellfish in the Inland Bays. Finally, the EPA, as part of its National Coastal Assessment program, has collected and analyzed fish samples from the Inland Bays for toxics.

None of the data and assessments to date have led the State to conclude that an advisory is needed in the Inland Bays as a result of contaminant sources within the Inland Bays. There is, however, an advisory on coastal species such as Bluefish and Striped Bass that migrate in and out of the Inland Bays.

C.2 MONITORING OF GROUNDWATER

C.2.1. DELAWARE GROUNDWATER MONITORING NETWORK

Description/Objective(s)

Long time-series of water levels in major aquifers serve as critical baseline data for resource management and analyses of aquifer response to pumping, climatic variability, drought hazards, seawater intrusion, and interaction with streams and their ecosystems. The Delaware Geological Survey (DGS) currently monitors groundwater levels in a network of 90 plus wells in Delaware (Figure C.7). 15 of the wells are located in the Inland Bays watershed. The data are maintained in a relational database and served to stakeholders via a web interface.

DGS is automating data collection, reduction, and archiving to increase efficiency and quality control while sustaining growth of the statewide network over time. This supports evaluation of the long-term availability and sustainability of the groundwater supply, management of the resource, and a myriad of uses by the environmental management, engineering, and science communities.

In anticipation of needs to monitor for saltwater intrusion, DGS installed salinity sensors in 2015 in wells at three locations, Indian River Inlet, Fenwick Island Seashore State Park, and Woodland Beach Wildlife Area.

Groundwater quality data are collected when funded. DGS is seeking resources to expand monitoring to include routine groundwater quality sampling and analyses.

Responsible Organization and Contact

Delaware Geological Survey

Contact: A. Scott Andres, DGS
asandres@udel.edu, 302-831-0599

Data Collection Cost

Estimated capital cost for wells in eastern Sussex County is ~\$815,000.

Data Use

This monitoring program supports evaluation of the long-term availability and sustainability of the groundwater supply, management of the resource, and other uses by the environmental management, engineering, and science communities.

Record of Collection

Records for the oldest wells go back as far as 1957. Details on each well, along with records of collection, are available on the DGS website at <http://www.dgs.udel.edu/datasets/recent-and-historical-groundwater-level-data>.

Collection Method

The DGS database holds over 16 million instrument measured and 31,000 manual measured groundwater levels. Instrumentation has generated approximately 51,000 daily temperature and 9400 specific conductance records.

Analysis Methods

Data are reviewed by DGS for quality control and then are made available to stakeholders through DGS online sources. Hydrograph data from stream gages in the area have been utilized to compare surface-water baseflow to nearby groundwater levels and to assess the impacts of pumping.

Data Location

DGS manages and stores the data. Data can be viewed or downloaded at <http://www.dgs.udel.edu/datasets/recent-and-historical-groundwater-level-data>.

Management Goal

The network supports evaluation of the long-term availability and sustainability of the groundwater supply, management of the resource, models of sea-level rise impacts, and many other uses by the environmental management, engineering, and science communities.




FIGURE C.7 - Map of wells included in the Delaware Groundwater Monitoring Network in 2015.

C.2.2 DELAWARE AGRICULTURAL SHALLOW GROUNDWATER MONITORING NETWORK

Description/Objective(s)

Studies in the Delmarva Peninsula have demonstrated that groundwater in shallow unconfined aquifers near agricultural lands is susceptible to contamination from chemicals, including nutrients, applied at the land surface. The agricultural community in Delaware has been working with various State and Federal government agencies to apply a number of conservation practices intended to reduce the amount of nitrate reaching the water table beneath agricultural land in support of Delaware's nutrient reduction goals. However, changes in nutrient management practices on the land surface may take decades to improve water quality in groundwater discharge to Delmarva streams because of groundwater residence times.

To understand if these changes in agricultural practices are reducing nitrate concentrations in shallow groundwater, and eventually in drinking water and streams, a long-term groundwater monitoring program is being implemented by the USGS, in partnership with the Delaware Department of Agriculture. This network of wells is located in areas with young, oxic, shallow groundwater, overlain by agricultural land, where change will be seen most clearly in a relatively short timeframe.

Much of the documented variability in nitrate concentrations on the Delmarva Peninsula included results from networks with wells in both oxic and anoxic aquifer condition. To maximize the discriminatory power of the statistical tests, the network will have a large sample size (up to 50 wells) and only include wells in oxic aquifer conditions.

Responsible Organization and Contact

USGS and Delaware Department of Agriculture

Contact: Brandon Fleming, USGS Hydrologist
bjfleming@usgs.gov, (443) 498-5561

Data Collection Cost

Not available.

Data Use

Results of nitrate analysis from this study are intended to provide a baseline data set which, if supplemented by sampling results collected under similar hydrologic conditions in future studies, could be used to observe trends in nitrate concentrations.

Record of Collection

October 2014 to present.

Collection Method

A groundwater monitoring network (Figure C.8), consisting of existing shallow wells from the Delaware Department of Agriculture's Pesticide network and USGS networks, has been established based on geochemical and land-use characteristics including:

- The presence of oxic aquifer conditions
- Geochemical indicators of agricultural land use based on major ion and nutrient concentrations
- Location of wells with respect to agricultural lands
- Depth of wells, and
- Available age dates for groundwater.

Analysis Methods

Samples have been collected for analysis of nutrients, major ions, and groundwater age and the results will be used to characterize groundwater quality and compare to existing groundwater quality data.

Data Location

U.S. Geological Survey, 5522 Research Park Drive, Baltimore, MD 21228

Management Goal

Evaluation of trends in nitrate concentrations.

Hypothesis and Test Statistics

To evaluate changes in nitrate concentrations between 1980 and 2014, statistical matched pair tests will be applied to new nitrate analyses from wells sampled during the study and historical analyses (where available).




FIGURE C.8 - Map of wells included in the Delaware Agricultural Shallow Groundwater Monitoring Network.

C.2.3 PUBLIC SUPPLY WELLS

Description/Objective(s)

Groundwater is the primary source of drinking water in the Inland Bays watershed. Although the Clean Water Act focuses primarily on the quality of navigable [surface] waters, Section 106(e) requires that groundwater quality be reported "...to the extent practicable." Thus, along with the biannual Combined 305(b) Report and 303(d) List developed for Delaware, a concurrent 305(b) Groundwater-Quality Assessment Based on Public-Well Data report is produced. The primary purpose of this report is to summarize and report raw or apparently raw groundwater-quality data collected from public water-supply.

Per U.S. EPA guidance, data are evaluated with respect to hydrogeologic setting and water-quality criteria where possible. The scope is limited to available data obtained from two primary sources: the Department of Health and Social Services (DHSS) Safe Drinking Water Information System (SDWIS) database and DNREC's Source Water Assessment and Protection Program (SWAPP) database. These databases are supplemented with data collected by Tidewater Utilities, Inc.

Responsible Organization and Contact

DNREC Division of Water, Water Supply Section, Groundwater Protection Branch

Contact: Joshua Kasper
Joshua.kasper@state.de.us, 302-739-9945

Data Collection Cost

Not available.

Data Use

Biannual reporting on ambient groundwater quality in compliance with the Clean Water Act.

Record of Collection

2001 to present.

Collection Method

Groundwater quality is assessed based on pre-existing information stored in the SDWIS and SWAPP databases. Queries extract SDWIS records of raw or apparently raw groundwater-quality data collected from public water-supply systems during the reporting period. For the most recent (2016) assessment, supplemental groundwater-quality data for 90 public wells were obtained from a private water utility, Tidewater Utilities, Inc. The records include well details such as DNREC ID, depth, geographic coordinates, geologic formation, aquifer, and aquifer type.

Analysis Methods

Access is used to link and extract data from SDWIS and the SWAPP databases. For wells with more than one analysis of a given analyte, results are averaged. Analytes not detected above laboratory quantitation limits ("nondetects") are treated as zeros in all calculations.

Results are evaluated with respect to Primary Maximum Contaminant Levels (PMCLs), Secondary Maximum Contaminant Levels (SMCLs), and Health Advisories (HAs) for public water-supply systems. Because only raw or apparently raw groundwater-quality data are evaluated, the results may not be representative of finished or treated water delivered to consumers. Therefore, an exceedance of a drinking-water standard does not necessarily indicate that a public water-supply system is not in compliance.

Data Location

DNREC's Source Water Assessment and Protection Program (SWAPP) database.

Hypothesis and Test Statistics

Where possible, data are evaluated with respect to aquifer type (i.e., unconfined, confined, semi-confined, fractured-rock, or karst). Some data also are evaluated with respect to sample depth. Evaluation of trends (e.g., concentration vs. depth) in this assessment are qualitative and not statistically derived. ArcMap is used for the spatial analysis of groundwater data. Tabulated statistics are the result Excel calculations. Golden Software, Inc.'s Grapher version 11 is used to construct percentile diagrams. Values plot as outliers on percentile diagrams if either of the following criteria are met:

$$\text{Value} < \text{QL} - 1.5 \times \text{IQR} \quad \text{or} \quad \text{Value} > \text{QU} + 1.5 \times \text{IQR}$$

Where:

IQR is the interquartile range (i.e., the difference between the 75th and 25th percentiles)

QL is the lower quartile or 25th percentile (i.e, the bottom of the box in Figure 4)

QU is the upper quartile or 75th percentile (i.e., the top of the box in Figure 4)

Differences between tabulated statistics and corresponding percentile diagrams are the result of differences in the computational methods of Excel and Grapher.

C.3 MONITORING OF WASTEWATER

C.3.1. POINT SOURCE WASTEWATER DISCHARGES

Description/Objective(s)

Point source discharges directly to surface water are regulated under the National Pollutant Discharge Elimination Systems (NPDES) Permits Program. Monitoring of these discharges is required by law.

The NPDES Program controls point source discharges to surface waters and land, respectively, by means of a permit which establishes the parameters, limits, schedules, and conditions for each discharge. Also included in this program are the compliance/monitoring reports prepared by the permittee; the surveillance, sampling and inspection of facilities; and an enforcement element. Limits are established based on minimum technology-based standards set pursuant to federal and State laws and regulations. More restrictive controls may be established, if deemed necessary, to meet surface water quality standards.

At this time, there are three point source wastewater facilities discharging to the Inland Bays: City of Rehoboth Beach WWTP (discharging to Rehoboth Bay), the City of Lewes WWTP (discharging to the Lewes-Rehoboth Canal), and the Allen Harim facility near Millsboro (discharging to Wharton Branch). The City of Lewes discharges only 2.5% of its effluent to the Inland Bays.

Responsible Organization and Contact

The DNREC Division of Water, Surface Water Discharges Section

Contact: Program Manager Compliance & Enforcement Branch
glenn.davis@state.de.us, 302-739-9946

Data Collection Cost

The costs of monitoring and reporting of results are borne by the permitted facilities. Funding sources for DNREC operations include Federal Section 106 and 205(g) excess funds, as well as State funding and permit fees. Total cost is not available.

Data Use

Discharge monitoring data are used to ensure that point source discharges into the Bays and tributaries of any pollutant, or combination of pollutants, meet all the applicable requirements under Clean Water Act Sections 301, 302, 306, 307, 308 and 403.

Record of Collection

Varies by permitted facility.

Collection Method

The effluent limitations, frequency of monitoring, parameters tested, and other special conditions vary between the individual facilities according to the requirements specified within their permits. Requirements may also vary between individual outfalls within a given facility. The permits are valid for five years, but may be administratively extended.

All of the discharge facilities monitor flow, and eutrophication indicators such as biological oxygen demand, total suspended solids, total phosphorus, total nitrogen, pH and dissolved oxygen. Some facilities also monitor fractions of phosphorus and nitrogen. Information concerning collection frequency and the type of samples collected is found in the excerpts from the permits of individual discharge facilities.

Analysis Methods

Each discharge facility and the contractors which they select to provide monitoring and laboratory services must adhere to all the USEPA-approved Methodology and Quality Assurance requirements specified within the permit. Delaware requires NPDES and Land Treatment permit holders to maintain records of all information resulting from any monitoring activities that are required in their permit.

Data Location

Previously data were accessible through the Delaware Environmental Navigator; however, data must now be requested directly through DNREC staff.

Management Goal

The data generated via NPDES monitoring also are used to calibrate and run the point source pollution component of the Hydrodynamic and Water Quality Model. These results determine the progress toward meeting the goals that were set by the TMDL for the Inland Bays of reducing loads of point source nitrogen by 10-15 percent and point source phosphorus by 60 percent.

C.3.2 LAND APPLICATION OF WASTEWATER

Description/Objective(s)

Land application of wastewater (spray irrigation or rapid infiltration basins) is increasingly being used within the Inland Bays watershed in lieu of point source discharges. This treatment approach uses the soil and terrestrial vegetation as a filter and storage system for wastewater constituents such as nutrients and bacteria.

Since land application has the potential to impact surface waters of the State, the Division of Water Resources, Ground Water Discharges Section, is responsible for facilitating and overseeing this activity. Land application facilities, like their NPDES counterparts, operate under a permit which establishes the parameters, limits, schedules, and conditions for each facility.

DNREC also requires compliance/monitoring reports prepared by the permittee; the surveillance, sampling and inspection of facilities; and an enforcement element. The primary objective of this monitoring is to ensure compliance with permit conditions.

Responsible Organization and Contact

The DNREC Ground Water Discharges Section, Large Systems Branch, reviews and approves spray irrigation wastewater systems, onsite wastewater treatment and disposal systems with daily flows greater than 2,500 gallons per day, Experimental/Alternative Technologies, Advanced Treatment Units, underground injection wells, and other means associated with land application wastewater treatment.

Contact: Ron Graeber, Program Manager
Ronald.Graeber@state.de.us, 302-739-9948

Data Collection Cost

The costs of monitoring and reporting of results are borne by the permitted facilities. Funding sources for DNREC operations include Federal Section 106 and 205(g) excess funds, as well as State funding and permit fees. Total cost is not available.

Data Use

Monitoring data are used to verify that the wastewater treatment process for a facility is functioning properly and that the land application activity does not adversely impact surface and groundwater quality in the area, or soils under the site.

The loading route to the Bays of constituents contributed by Land Treatment facilities is considered to be nonpoint source. Presently, nonpoint source loadings to the Bays are estimated using concentration and flow data collected at the tidal/nontidal interface. The land application monitoring data has not been used to estimate the percentage of the total nonpoint source load that is attributable to these facilities.

Record of Collection

Varies by permitted facility.

Collection Method

Following treatment at a wastewater treatment facility, the reclaimed water is tested for a variety of parameters to ensure that the reclaimed water meets appropriate treatment standards. Then, when weather conditions are suitable for irrigation, the reclaimed water is applied to the field at agronomic rates. Agronomic loading rates are determined by the nutrient levels of the reclaimed water and the nutrient needs of the crops being grown, and should be incorporated into the farm managers Nutrient Management Plan.

The effluent limitations, frequency of monitoring, parameters tested, and other special conditions vary between the individual facilities according to the requirements specified within their permits. This specific information is available for each facility within Attachment 8. Land Treatment permits also require groundwater and soil monitoring, in addition to the limitations an individual permit excerpts (Attachment 8). For one facility (the Town of Georgetown) surface water monitoring is also required in two adjacent streams. The parameters typically monitored in sprayed effluent are similar to NPDES requirements, including eutrophication indicators such as total suspended solids, biological oxygen demand, total phosphorus, total nitrogen (and fractions of nitrogen), pH and dissolved oxygen. Groundwater monitoring includes measurement of water table depth and constituents that are highly soluble and do not readily adhere to soil (under and adjacent to the spray site). Soil monitoring is done to ensure that normal soil functioning is maintained so that expected levels of effluent treatment can occur.

Analysis Methods

Each discharge facility and the contractors which they select to provide monitoring and laboratory services must adhere to all the USEPA-approved Methodology and Quality Assurance requirements specified within the permit. Delaware requires NPDES and Land Treatment permit holders to maintain records of all information resulting from any monitoring activities that are required in their permit.

Data Location

Data must be requested from the Groundwater Discharges Section at DNREC.

Management Goal

Limits are established based on minimum technology-based standards set pursuant to federal and State laws and regulations. More restrictive controls may be established, if deemed necessary, to meet Federal or State drinking water quality standards. If background conditions exceed the drinking water standards, then there shall be no concentration increase above the background levels.

C.4 Monitoring of ATMOSPHERIC DEPOSITION

C.4.1 ATMOSPHERIC DEPOSITION OF NUTRIENTS

Description/Objective(s)

Nutrients are deposited from the atmosphere directly into the Bays during both wet and dry weather. Deposition of nitrogen is the most significant.

Data from the National Atmospheric Deposition Program/Atmospheric Integrated Research Monitoring Network (NADP/AIRMoN) site at Cape Henlopen can be used to estimate wet deposition rates (fluxes associated with rainfall) of nitrate (NO_3^-), ammonium (NH_4^+), and phosphate (PO_4^{3-}) to the open waters and the tidal marshes of the Inland Bays on a daily basis. These daily measurements may be summed to determine wet deposition over other time periods.

There are, however, other species in wet deposition, and atmospheric inputs due to another set of chemical and physical processes, collectively described as dry deposition, that deliver N and P to land and water surfaces in the absence of rainfall. These species and processes may also contribute significantly to nutrient delivery to the Bays and their watersheds.

At present, the Cape Henlopen site is the only active site in the Inland Bays watershed where precipitation chemistry is determined (Site ID DE02, Latitude 38.7722, Longitude -75.0992). Therefore, it must serve as the basis for calculating current and future wet deposition rates.

Procedures to calculate the wet and dry deposition rates of N and P species directly to the Inland Bays are described in Ullman et al. (2010).

Responsible Organization and Contact

University of Delaware, College of Earth, Ocean, and Environment

Contact: Joseph Scudlark
scudlark@udel.edu, 302-645-4300

Data Collection Cost

Not available.

Data Use

Data are used to calculate the wet and dry deposition rates of nitrogen and phosphorus species to open waters and contiguous marshes in the Inland Bays, and elsewhere in Delaware for use in mass balance studies.

Record of Collection

1993 to present.

Collection Method

Samples are collected daily within 24 hours of the start of precipitation, often providing data for all or part of a single storm. The AIRMoN sites are equipped with a wet-only deposition collector and precipitation gage. Each site also has a National Weather Service standard gage for reporting storm total precipitation.

Samples are refrigerated after collection and are sent in chilled insulated shipping containers to the Central Analytical Laboratory (CAL) at the Illinois State Water Survey, where they are kept refrigerated until analysis.

Analysis Methods

The CAL measures free acidity (H^+ as pH), conductance, calcium (Ca^{2+}), magnesium (Mg^{2+}), sodium (Na^+), potassium (K^+), sulfate (SO_4^{2-}), nitrate (NO_3^-), chloride (Cl^-), and ammonium (NH_4^+). The CAL also measures orthophosphate, but only for quality assurance as an indicator of sample contamination.

The CAL reviews field and laboratory data for completeness and accuracy, and flags samples that were mishandled, compromised by precipitation collector failures, or grossly contaminated. The CAL delivers all data and information to the NADP Program Office, which applies a final set of checks and resolves remaining discrepancies. Data then are made available on the NADP Web site.

Methods to calculate N and P deposition are described in detail in Ullman et al., 2010.

Data Location

Data from the Lewes NADP/AIRMoN site (DE02; located in Cape Henlopen State Park, Lewes) is accessed through the main NADP/AIRMoN website at <http://nadp.sws.uiuc.edu/airmon/> or from the AIRMoN Data Retrieval Site at <http://nadp.sws.uiuc.edu/AIRMoN/>. Additional information about the site and data request forms, can be found there.

Management Goal

Despite significant uncertainties in the absolute values of the deposition rates calculated (Ullman et al., 2010), temporal trends may still be revealed. Based on a comparison with samples collected simultaneously on the north shore Indian River Bay, it is understood that extrapolating from Cape Henlopen rain data to the entire Inland Bays represents a conservative (minimum) estimate of wet N deposition rates for this system.

C.5 MONITORING OF WETLANDS

3.5.1. STATE WETLANDS MONITORING AND ASSESSMENT PROGRAM

Description/Objective(s)

The goal of DNREC's Wetland Monitoring and Assessment Program (WMAP) is to assess the condition, function, and services of wetlands in the state, and to integrate the latest research to understand the connection between the metrics and measures that are evaluated and the actual processes and implications on services that wetlands provide. This information is used to inform the citizens of Delaware and to improve existing education, restoration, protection, and land use planning efforts. The *Delaware Wetland Monitoring Strategy* (Delaware Department of Natural Resources and Environmental Control, 2011) guides the WMAP's efforts in the areas of protocol development, wetland monitoring and assessment activities, research, and application of information. The goals and objectives outlined in the monitoring strategy support many of the goals of the *Delaware Wetland Conservation Strategy* (Delaware Department of Natural Resources and Environmental Control, 2008).

The Program works closely with other states through the Environmental Protection Agency's Mid-Atlantic Wetlands Program to establish and conduct research methods and share information.

Objectives include:

- Develop scientifically valid wetland assessment methods.
- Assess the current condition of wetlands by watershed and identify major stressors that are impacting wetlands.
- Perform research to improve our understanding of wetland functions, the impact of stressors, and the ecosystem services provided by wetlands to humans and the environment.
- Evaluate the performance of wetland restoration and other compensatory wetland mitigation in replacing wetland acreage and function.
- Educate other state agencies, conservation partners, and the general public to improve efforts to protect and restore wetlands.
- Integrate monitoring and assessment data into watershed restoration plans and other conservation strategies.
- Meet requirements of the Clean Water Act.

The watersheds of the state were prioritized for wetlands monitoring based largely on the TMDL implementation schedule (Figure C.9). The intent of the state is to monitor these watersheds using a rotating basin approach once an initial assessment of the wetlands within each watershed has been performed. Monitoring of nontidal wetlands in the Inland Bays occurred in 2005-2006; tidal wetlands were monitored in 2008 (Jacobs et al., 2008; Rogerson et al., 2008).

Responsible Organization and Contact

DNREC Wetland Monitoring and Assessment Program

Contact: Alison Rogerson
Alison.Rogerson@state.de.us, (302) 739-9939

Data Collection Cost

Not available.

Data Use

Final reports documenting the condition of tidal and nontidal wetlands in the Inland Bays were completed in 2009 (Jacobs et al., 2009; Rogerson et al., 2009). The data obtained by DNREC from assessing wetlands is being used to design wetland restoration plans for watersheds and to better understand how certain land use decisions affect the health of our wetlands.

Record of Collection

Nontidal wetlands: 2005-2006

Tidal wetlands: 2008

Collection and Analysis Methods

Nontidal riverine sites and flats in the Inland Bays are surveyed using a combination of comprehensive and rapid assessment procedures:

DECAP - The Delaware Comprehensive Assessment Procedure is a comprehensive assessment method for collecting data that can be used to determine the condition of a wetland site relative to reference condition (closest to natural and undisturbed). DECAP can be used to assess flat, riverine and depressional nontidal wetland subclasses in the Coastal Plain of Delaware and Maryland. The comprehensive procedure can produce scores for certain wetland functions (services), including habitat, plants, hydrology, buffers, and soil cycling.

DERAP - The Delaware Rapid Assessment Procedure is a rapid field method for determining the general condition of a wetland site. The DERAP can be used in flat, riverine, and depressional wetlands in Delaware and Maryland.

Tidal wetland surveys use a rapid assessment protocol:

MidTRAM - The MidAtlantic Tidal Rapid Assessment Method is a rapid protocol for assessing the condition of estuarine emergent tidal wetlands in Delaware, Maryland, and Virginia. The MidTRAM is validated with intensive biological data based on the bird community and biomass levels.

Data Location

Data are managed and retained by the DNREC Wetland Monitoring and Assessment Program. All data are entered into an Access computer database that has been developed to consistently store wetland assessment data. The Department plans eventually to make wetlands data available through STORET.

Management Goal

DNREC's goal is to achieve an annual net gain in wetland acreage and condition. The *Delaware Wetland Conservation Strategy* highlights recommends approaches with measurable outcomes for enhancing and improving wetland protection.




FIGURE C.9 - DNREC's current schedule for statewide wetland assessments.

C.5.2 LONG-TERM SALT MARSH MONITORING

Description/Objective(s)

Three representative salt marshes in the Inland Bays were each instrumented with triplicate Sediment Elevation Tables (SETs) in order to provide high intensity baseline information on sediment elevation changes in salt marshes of the Inland Bays. Locations of the SETs are provided in Figure C.10. In addition, one marsh (Angola Neck) was outfitted with continuous water level loggers to determine water depth at marsh surface locations. These data can be used to assess whether Inland Bays marshes are keeping pace with sea level rise, and can provide information on potential causal or influential factors of sudden wetland dieback events.

Responsible Organization and Contact

The Delaware Center for the Inland Bays is responsible for this monitoring program in the watershed.

Contact: Andrew McGowan, Environmental Scientist
environment@inlandbays.org, 302-226-8105, x112

Data Collection Cost

Costs for each SET platform are roughly \$900 for initial installation, not including labor costs, and \$50 annually for maintenance. The SET arm costs \$2,000 but can be used at all sites because it is portable. Water loggers cost approximately \$2,200 each.

Labor cost is roughly \$5,000 a year to monitor the SET tables and water loggers.

Data Use

The information will be used to better understand the variation among processes that result in fringing marsh maintenance or conversion to open water, and will be included in the development of saltmarsh restoration and protection strategies for the watershed.

Record of Collection

Angola Neck: 2008 – present

Piney Point: 2010 – present

Slough's Gut: 2011 - present

Collection Method

Data are collected twice annually, within 5 days of the full moon in October and April. Each SET table has both a deep SET, which measures overall marsh height change, and a shallow SET, which measures changes in the marsh height as a result of both surface accretion/erosion, and changes in the root zone. Sampling is conducted by attaching the SET arm to a SET table, and placing nine SET pins into the corresponding holes on the SET arm. The pins are gently lowered to the marsh surface, and the height each pin extends above the SET arm is recorded for all nine pins to the nearest half millimeter. The pins are then removed, the SET are rotated 90 degrees and the process is repeated until all four directions are sampled. Both the shallow SET




FIGURE C.10 - Locations of sediment elevation tables in the Inland Bays.

and the deep SET are measured in this way. Additionally, each SET table has three feldspar clay accretion plots, which are sampled by cutting away a plug with a knife, and measuring the height of sediment above the top of the clay layer at three different locations on the plug.

Water level data are downloaded from each logger and corrected for barometric pressure using Win-Situ software.

Analysis Methods

Data are analyzed to determine if each SET platform is keeping pace with the current rate of sea level rise by first determining the slope of the change in height for each individual pin at each direction at each SET over time with a linear regression (ex. 9 slopes for direction A at Angola Neck site 1, one for each pin, then 9 slopes for direction B at Angola Neck site 1, one for each pin...etc.). Only the deep SET measurements are used as these reflect all changes in marsh height, as opposed to the shallow SET which only reflects root zone and surface layer changes. The coefficient of each slope is then averaged together for all 4 directions at each SET location (ex. 36 slope coefficients averaged at Angola Neck site 1) to determine the average rate of change at a particular SET location. The standard error is also calculated. These values are compared to the mean sea level rise rate of 3.40 mm per year with a lower tailed T-test ($\alpha = 0.05$). If the result of the test is significant, the SET table has significantly less wetland elevation change than local sea level rise. This follows the protocol recommended by the National Park Service.

To assess what specific processes are driving the changes in marsh height, each SET table is analyzed separately using the shallow SET readings, the deep SET readings, and accretion data. As described previously, a slope is calculated for each pin at each position for each SET for the shallow and deep platforms. The slopes are averaged together at each SET to determine the overall slope for a particular SET table. This procedure is performed for both the deep SET readings and the shallow SET readings. The shallow SET readings capture changes occurring only in the root zone and surface layer, or roughly the first 0.6 m of the marsh surface. These changes include root zone growth, compaction, and accretion. By subtracting the shallow SET slope from the corresponding deep SET slope, the changes occurring below the 0.6 m root zone are separated from the overall deep SET measurements. In this way, we are able to determine what is occurring below the root zone.

To determine accretion rate, the nine accretion measurements are averaged together to get the average accretion height. If the height reading is the first reading since laying down a new layer of clay, this height is divided by the number of days since the clay was laid onto the marsh. If the height value is not the first measurement since laying the clay onto the marsh, the height value is subtracted from the previous sampling event's average height value, and divided by the number of days since the last measurement. This is done to prevent previous accretion events from influencing the current measurements. Each average change in height is averaged together to determine the overall average accretion rate. The average accretion rate for each SET table is then subtracted from the shallow slope values to determine the changes occurring solely in the root zone.

Data Location

Located at the Delaware Center for the Inland Bays, available upon request.

Management Goal

Inform decision makers on how salt marshes in the Inland Bays are faring with respect to sea level rise, and what processes are responsible for maintenance of marsh elevation or conversion to open water.

Hypothesis and Test Statistics

Elevation at each SET table and each marsh as a whole is compared to current sea level rise rates, along with DNREC sea level rise planning scenarios using a lower tailed T-test ($\alpha = 0.05$). The null hypothesis is that the SET platform has equal to or greater wetland elevation change than local sea level rise.

C.6 MONITORING OF LIVING RESOURCES AND HABITATS

C.6.1 NATIONAL AQUATIC RESOURCE SURVEYS

Description/Objective(s)

The National Aquatic Resource Surveys (NARS) are statistical surveys designed to assess the status of and changes in quality of the nation's coastal waters, lakes and reservoirs, rivers and streams, and wetlands. Using sample sites selected at random, these surveys provide a snapshot of the overall condition of the nation's water. Because the surveys use standardized field and lab methods, results can be compared from different parts of the country and between years. EPA works with state, tribal and federal partners to design and implement the NARS.

The surveys are designed to answer questions such as:

- What percent of waters support healthy ecosystems and recreation?
- What are the most common water quality problems?
- Is water quality improving or getting worse?
- Are investments in improving water quality focused appropriately?

These surveys are providing nationally-consistent water quality information. Additionally, the national surveys are helping to build stronger water quality monitoring programs across the country by fostering collaboration on new methods, new indicators and new research.

The NARS are made up of four individual surveys that are implemented on a rotating basis. Stations in the Inland Bays watershed during previous field seasons are shown in Figure C.11. A summary of the indicators used in each survey is provided in Table C.7.

- National Coastal Condition Assessment (NCCA)

Goals: (1) What percent of the Nation's coastal waters are in good, fair, and poor condition for key indicators of water quality, ecological health, and recreation? (2) What is the relative importance of key stressors such as nutrients and contaminated sediments?

Design: The NCCA sampling is comprised of coastal waters extending from the shoreline to the nearshore boundary of the open water of the oceans and Great Lakes. The assessment is limited to the fringing, shallow band of coastal waters most heavily used by humans and most vulnerable to activities within adjacent coastal watersheds.

- National Lakes Assessment (NLA)

Goals: (1) What is the current biological, chemical, physical and recreational condition of lakes? (2) Is the condition of lakes getting better, worse, or staying the same over time? (3) Which environmental stressors are most associated with degraded biological condition in lakes?

Design: The NLA sampling is comprised of natural lakes, ponds, and reservoirs across the lower 48 states. Starting with the NLA2012, to be included in the survey, a water body had to be a natural or man-made freshwater lake, pond or reservoir, greater than 2.47 acres (1 hectares), at least 3.3 feet (1 meter) deep, and with a minimum quarter acre (0.1 hectare) of open water. Lakes had a minimum retention time of 1 week. The Great Lakes and the Great Salt Lake were not included in the survey, nor were commercial treatment and/or disposal ponds, brackish lakes, or ephemeral lakes. The NLA 2007 assessed only those lakes greater than 10 acres (4 hectares) in size.

- National Rivers and Streams Assessment (NRSA)

Goals: The goals of the NRSA are to determine the extent to which rivers and streams support a healthy biological condition and the extent of major stressors that affect them. The survey supports a longer-term goal: to determine whether our rivers and streams are getting cleaner and how we might best invest in protecting and restoring them. Additionally, the survey compares the condition of streams to those of an earlier study that focused on small

streams (the Wadeable Streams Assessment or WSA) conducted by the U.S. Environmental Protection Agency and its partners in 2004.

Design: The NRSA assesses the ecological condition of the full range of flowing waters in the conterminous U.S. (lower 48 states). The target population includes the Great Rivers (such as the Mississippi and the Missouri), small perennial streams, and urban and non-urban rivers. Run-of-the-river ponds and pools are included, along with tidally influenced streams and rivers up to the leading edge of dilute sea water.

- National Wetland Condition Assessment (NWCA)

Goals: The NWCA is designed to answer basic questions about the extent to which U.S. wetlands support healthy ecological conditions and the prevalence of key stressors at the national and regional scale. It is intended to complement and build upon the achievements of the U.S. Fish and Wildlife Service Wetland Status and Trends Program, which characterizes changes in wetland acreage across the conterminous United States. Paired together, these two efforts provide government agencies, wetland scientists, and the public with comparable, scientifically-defensible information documenting the current status and, ultimately, trends in both wetland quantity (i.e., area) and quality (i.e., ecological condition).

Design: The survey design is developed in partnership with the US FWS Wetlands Status and Trends Program. The NWCA sampling is comprised of all wetlands of the conterminous U.S. The survey encompasses both tidal and nontidal wetlands ranging from the expansive marshes of our coasts to the forested swamps, meadows, and waterfowl-rich prairie potholes of the interior plains.




FIGURE C.11 – Southern Delaware sites at which NARS data were collected for published reports to date.

Responsible Organization and Contact

U.S. Environmental Protection Agency, Office of Wetlands, Oceans, and Watersheds
National Aquatic Resource Surveys
1200 Pennsylvania Avenue, NW (Mailcode 4503T), Washington, DC 20460
<https://www.epa.gov/national-aquatic-resource-surveys>

Data in Delaware are collected by DNREC Environmental Laboratory Section.
Contact: Kathy Knowles, Kathy.knowles@state.de.us, 302-739-9942

Data Collection Cost

Not available.

Data Use

The U.S. EPA publishes reports for each survey. In addition, the data are publically available for use in research.

Record of Collection

NCCA: The first NCCA sampling field season was conducted in 2010. The most recent field season was conducted in 2015.

NLA: NLA field season sampling is conducted every five years. Previous field seasons were conducted in 2007 and 2012.

NRSA: NRSA sampling field seasons were conducted in 2008-2009 and 2013-2014. The next field season will be conducted in 2018-2019.

NWCA: The 2011 National Wetland Condition Assessment (NWCA) was the first sampling season. The second field sampling season was conducted in 2016.

Collection Method

Samples for Delaware sites are collected by the DNREC Division of Water, Environmental Laboratory Section.

Field Operations Manuals are available on the EPA website: <https://www.epa.gov/national-aquatic-resource-surveys/manuals-used-national-aquatic-resource-surveys>.

Analysis Methods

Field, laboratory, quality assurance and site evaluation manuals are available on the EPA website: <https://www.epa.gov/national-aquatic-resource-surveys/manuals-used-national-aquatic-resource-surveys>.

Data Location

NARS data are available for download on the EPA website: <https://www.epa.gov/national-aquatic-resource-surveys/data-national-aquatic-resource-surveys>.

Management Goals

Within each survey, the goals are:

- For each indicator of condition, estimate the proportion of the nation's waters in degraded condition within a $\pm 5\%$ margin of error and with 95% confidence.
- For each indicator of condition, estimate the proportion of waters or resources in a specific ecoregion that fall below the designated threshold for good conditions for selected measures within a $\pm 15\%$ margin of error and with 95% confidence.
- Estimate the proportion of waters ($\pm 7\%$) that have changed condition classes for selected measures with 95% confidence.

Hypothesis and Test Statistics

These are detailed in the respective Quality Assurance Project Plans, available at <https://www.epa.gov/national-aquatic-resource-surveys/manuals-used-national-aquatic-resource-surveys>.

TABLE C.7 - Indicators Evaluated for the National Aquatic Resource Surveys.

BIOLOGICAL	CHEMICAL/TOXICITY	PHYSICAL	RECREATIONAL/HUMAN HEALTH
National Coastal Condition Assessment (NCCA)			
<ul style="list-style-type: none"> • Benthic macroinvertebrates • Chlorophyll a • Ecological fish tissue contaminants 	<ul style="list-style-type: none"> • Dissolved oxygen • Nitrogen • Phosphorus • Salinity • Sediment contaminants • Sediment toxicity 	<ul style="list-style-type: none"> • Water clarity 	<ul style="list-style-type: none"> • Human health fish tissue contaminants* (Great Lakes only)
National Lakes Assessment (NLA)			
<ul style="list-style-type: none"> • Benthic macroinvertebrates • Chlorophyll a • Zooplankton 	<ul style="list-style-type: none"> • Acidification • Atrazine • Dissolved oxygen • Nitrogen • Phosphorus • Sediment mercury 	<ul style="list-style-type: none"> • Drawdown • Human disturbance • Lakeshore habitat • Physical habitat complexity • Shallow water habitat 	<ul style="list-style-type: none"> • Algal toxin (microcystin) • Cyanobacteria
National Rivers and Streams Assessment (NRSA)			
<ul style="list-style-type: none"> • Benthic macroinvertebrates • Periphyton (algae) • Fish community 	<ul style="list-style-type: none"> • Phosphorus • Nitrogen • Salinity • Acidity 	<ul style="list-style-type: none"> • Streambed sediments • In-stream fish habitat • Riparian vegetative cover • Riparian disturbance 	<ul style="list-style-type: none"> • Enterococci (fecal indicator) • Mercury in fish tissue
National Wetland Condition Assessment (NWCA)			
<ul style="list-style-type: none"> • Benthic macroinvertebrates • Chlorophyll a • Fish assemblage • Fish tissue contaminants • Macrophytes • Phytoplankton • Sediment diatoms • Wetland vegetation (introduced species) • Wetland vegetation (plant community) • Zooplankton 	<ul style="list-style-type: none"> • Acidification • Atrazine • Conductivity • Dissolved oxygen • Nitrogen • Phosphorus • Salinity • Sediment enzymes • Sediment mercury • Soil chemistry 	<ul style="list-style-type: none"> • Lakeshore habitat/riparian vegetative cover • Human disturbance • Physical habitat complexity • Shallow water habitat/in-stream fish habitat • Streambed sediments • Water clarity 	<ul style="list-style-type: none"> • Algal toxins (microcystin) • Cyanobacteria • Enterococci • Fish tissue contaminants

C.6.2. STATEWIDE VEGETATION COMMUNITY & LAND COVER MAPPING PROJECT

Description/Objective(s)

The Delaware Statewide Vegetation Community Mapping Project seeks to map all of the vegetation communities and land covers present in the state of Delaware. Delineations are drawn to the finest extent possible (no defined minimum mapping unit) using aerial imagery analysis, field observations, and data obtained from others. Approximately 10-20% of the state has been field checked. The project began with a map of the Brandywine Creek and was extended to include the entire state of Delaware.

Responsible Organization and Contact

DNREC Wildlife Species Conservation and Research Program

Contact: Joseph Rogerson
joseph.rogerson@state.de.us, 302-735-3600

Data Collection Cost

Not available.

Data Use

Currently, the maps are used for determining the rarity and status of vegetation communities in Delaware, environmental reviews, and management plans on public lands. These maps can be used as a baseline for sea level rise studies, climate change, and change over time studies from historical imagery.

Record of Collection

Historical analyses and maps of vegetation communities, land covers, and habitats have been published for the following areas in the Inland Bays:

- Thompson Island Nature Preserve (Coxe, 2011)
- Assawoman Wildlife Area (Coxe, 2012a)
- Delaware Seashore State Park (Coxe, 2012b)
- Fenwick Island State Park (Coxe, 2012c)
- Cape Henlopen State Park (Coxe, 2012d)
- James Farm Ecological Preserve (Coxe, 2013)

Collection Method

Field data is obtained primarily from environmental reviews and surveys of public lands in Delaware. Land covers are obtained from the same methods and impervious surfaces are from 2007 impervious surface layer, except for the Brandywine and Red Clay Creek watersheds which were digitized from 2002 aerial imagery. Vegetation communities are determined using the *Guide to Delaware Vegetation Communities* which is derived and linked to the National Vegetation Classification System (NVCS). The NVCS is a national effort by The Nature Conservancy and NatureServe to standardize the names and classification of vegetation communities in North America. Common names of the vegetation communities in Delaware are the same as those used in the NVCS.

Analysis Methods

All shapefiles for the Delaware Statewide Vegetation Community Map are organized by watershed. Each watershed file has a year after (i.e. 1997, 2002, or 2007) that signifies the imagery that the map is based on. About once a month a complete map will be produced which will be called Delaware Statewide Vegetation Community Map with the date after it.

Data can be used to map a particular vegetation community for a watershed, or to query how much acreage of a vegetation community is present in a watershed.

Data Location

DNREC Wildlife Species Conservation and Research Program, and the University of Delaware Water Resources Agency.

C.6.3 SEAWEED MONITORING

Description/Objective(s)

Small to medium amounts of macroalgae are healthy for estuarine systems. They provide habitat for blue crabs, as well as numerous species of fish, especially in the absence of seagrass beds. However, excess macroalgae can have the opposite effect by smothering benthic organisms and creating hypoxic zones particularly during the early summer mornings before photosynthetic activity resumes. In order to assess levels of macroalgae over time, particularly in response to efforts to reduce nutrient pollution to the Inland Bays, 12 locations in Rehoboth and Indian River Bay are monitored for macroalgae.

The objectives of this study are to see if macroalgae types, distribution, and density appear remarkably different than previously observed levels, and to assess the abundance and distribution of macroalgae over a full growing season in Indian River Bay and Rehoboth Bay. Furthermore, the rapid macroalgae sampling approach employed in this study can be investigated for its feasibility as a means for more regular monitoring of macroalgae abundance and distribution using citizen volunteers.

Responsible Organization and Contact

The Delaware Center for the Inland Bays is responsible for this monitoring program beginning in the spring of 2017. Previous to this, DNREC Division of Water was in charge of the monitoring program.

Contact: Andrew McGowan, Environmental Scientist
environment@inlandbays.org, 302-226-8105, x112

Data Collection Cost

Roughly \$2,000 for CIB staff time per year
Equipment: \$100 in gas and supplies

Data Use

Data is used to track macroalgae abundance over time, and to assess seasonal abundance and distribution.

Record of Collection

Data has been collected irregularly, with sampling occurring in 1999, 2009, 2011, and 2012.

Collection Method

Macroalgal density is determined using a stainless steel grappling hook tossed off the windward side of a boat. Sampling occurs once a month at 12 sites from May through September (Figure C.12). The hook is 25 cm long, has a width of 24.3 cm with six tines spaced about 9 cm apart and is attached to a 10 m length of 0.95 cm diameter nylon line. The hook is tossed off the windward side of the boat and allowed to settle to the bottom, at which time the line is given five steady tugs and then the hook is hauled into the boat. It was determined that the five tugs result in the hook covering a distance across the bottom of about 3 to 4 m. Three tosses will constitute a single sample that will be placed into a sieve bucket graduated in liters. The bucket will then be shaken from side-to-side a few times to settle and uniformly distribute the algae.

Analysis Methods

The approximate amount of area covered by the grappling hook during the three tosses is 2.5 m. (3.5 meters per toss x width of hook, 0.24m x 3 tosses). The sample of algae collected is reported as liters of algae. Density is categorized as light (0 to 3 liters), moderate (4 to 7 liters) and heavy (> 8 liters). Dominant groups of macroalgae (e.g. Ulva, Gracilaria, and Agardhiella) are sorted while remaining non-dominant groups are lumped together. Relative percentages of the groups are determined visually.

Data Location

Located at the Delaware Center for the Inland Bays, available upon request.

Management Goal

Document the abundance of macroalgae in the Inland Bays over time, including seasonal patterns of abundance and distribution. Data generated through this program can be used to gauge the potential success of sea grass restoration in areas near sampling locations, and can be used in conjunction with water quality measurements to gauge the success of nutrient pollution reduction efforts.

Hypothesis and Test Statistics

Abundance between sites as well as between years for macroalgae in general, along with each dominant group, can be compared using Kruskal Wallis tests ($\alpha = 0.05$).




FIGURE C.12 - Map showing location of seaweed monitoring stations in Rehoboth and Indian River Bays.

C.6.4 COASTAL FINFISH ASSESSMENT SURVEY

Description/Objective(s)

The relative abundance and distribution of a number of recreationally important finfish species are assessed using trawl sampling in the Delaware Estuary and Delaware's Inland Bays (Indian River and Rehoboth Bays). The 16-foot trawl survey used in the Inland Bays is primarily intended to monitor juvenile fish abundance.

Responsible Organization and Contact

DNREC Division of Fish and Wildlife, Fisheries Section

Contact: Michael Greco
Michael.Greco@state.de.us, 302-735-6780

Data Collection Cost

Not available.

Data Use

The indices generated from these surveys are used in the development of interstate fishery management plans and stock assessments. In particular, the surveys are used in the Weakfish (*Cynoscion regalis*), Striped Bass (*Morone saxatilis*), Bluefish (*Pomatomus saltatrix*), Atlantic Menhaden (*Brevoortia tyrannus*), Black Drum (*Pogonias cromis*) and Summer Flounder (*Paralichthys dentatus*) fishery management plans and stock assessments. In addition, data from the surveys are used in establishing time of year restrictions for beach replenishment and dredging. The CIB uses the data to develop fish abundance indicators.

The surveys also serve as platforms for providing specimens to researchers studying life history and biology of fishes. These surveys have provided samples for use in genetics, tissue contaminants, age and growth, food habits, reproduction, and many other studies.

Record of Collection

The Delaware Bay trawl surveys were expanded in 1986 to include monthly sampling (April to October) in the Indian River and Rehoboth Bays.

Collection Method

Sampling with a 16-foot semi-balloon otter trawl is conducted monthly from April through October at 12 fixed stations in the Inland Bays (Figure C.13).

Sampling at each station consists of a ten-minute trawl tow, typically made against the prevailing tide. Occasionally, tows less than ten minutes are made in cases of unforeseen gear conflicts, draft considerations, etc. In such cases, tows are required to be at least five minutes in duration to be considered valid. Catches from short tows are standardized to ten minutes.

The trawl is hauled over the stern and the catch emptied on a sorting table upon completion of each tow. Fish were sorted by species and enumerated. A representative subsample of 30 specimens per species is measured for fork length to the nearest half centimeter; the remainder are enumerated. Surface temperature, salinity, and dissolved oxygen, tidal stage, weather conditions, water depth, and engine speed are recorded at the beginning of each tow.

Analysis Methods

Data analysis includes monthly and annual summaries of the catch including a listing of species collected, total number of each species taken, mean catch per tow, and standard deviations. Mean surface salinities and temperatures are calculated similarly by month, station or subarea. Annual young-of-the-year index values are calculated, as geometric mean catch per tow, for target species.




FIGURE C.13 - Map showing DNREC otter trawl survey locations in the Inland Bays.

Data Location

Data are maintained by the Division of Fish and Wildlife Fisheries Section.

Management Goal/Hypothesis and Test Statistics

Species-specific Stock Assessment Subcommittees test geometric means for individual assessments. First, the relative prevalence for that particular species is determined to see if the survey interacts with it enough to warrant inclusion in the assessment. Next, the survey may be standardized using a generalized linear model. Further, trend analyses are conducted on the survey itself (using ARIMA, Mann-Kendall tests), and combined with age, growth (Von Bertalanffy), maturity F, M (Lorenzen) data for analyses.

C.6.5 VOLUNTEER INSHORE FISH AND BLUE CRAB SURVEY

Description/Objective(s)

The shorelines of the Inland Bays provide critical habitat to many juvenile and young-of-the-year fish species along with blue crabs. In an effort to obtain fish population and diversity data from these inshore areas, an Inland Bays volunteer fish monitoring program was implemented collect data that can be used to create fish indices that complement the data being collected by the DNREC Division of Fish and Wildlife Trawl Survey. The objectives of this program are to conduct seining surveys to determine the abundance, species diversity, and body lengths of the near-shore fish communities in the Delaware Inland Bays and to measure physical and biological parameters to determine which conditions are favorable for nearshore fish and blue crabs in the Inland Bays.

Responsible Organization and Contact

The Delaware Center for the Inland Bays is responsible for this monitoring program.

Contact: Andrew McGowan, Environmental Scientist
environment@inlandbays.org, 302-226-8105, x112

Data Collection Cost

Equipment: \$700 for nets, buckets, first aid kits, tongs, miscellaneous gear
Staff time: \$5,400

Data Use

The data collected from this project will be used primarily by the Delaware Division of Fish and Wildlife and the Center for the Inland Bays. These data will be used to help document trends, seasonality, and annual variability in fish populations over time. Additionally, this data will be used to assess the importance of the shorezone waters for commercially and recreationally important species.

Record of Collection

2011 to present.

Collection Method

16 shoreline sites are monitored once in April and twice a month from May through October using a 30-foot-long 4-foot-tall seine net (Figure 3.10). To sample a beach, one volunteer holds one end of the seine net along the beach while another volunteer wades out with the opposite end until the net is fully extended (30 ft). Both volunteers drag the net for 70 feet along the shoreline, with the volunteer in the deeper water walking slightly ahead of the shallow water volunteer. At the 70-foot mark, the inshore volunteer stops and the deep water volunteer swings inshore with the net, at which point both volunteers drag the net ashore having seined 100 feet of shoreline. All fish are identified to species and counted, and the first 25 fish of each species are also measured to the nearest millimeter. The number of blue crabs is also counted. In addition, physical parameters such as dissolved oxygen, water temperature,

salinity, wave height, amount of rain in the last 24 hours, and wind speed are also recorded prior to each seine.

Analysis Methods

To assess species abundance over time, geometric mean catch for each species is calculated each year. Differences in catch between the bays are assessed with pairwise Wilcoxon rank sum tests ($\alpha = 0.05$). To assess which variables are important to inshore fish species, correlations to physical variables are assessed for recreationally or commercially important species using Kendall's Tau correlation tests ($\alpha = 0.05$).

Data Location

Data are stored at the Center for the Inland Bays and are made available online at www.inlandbays.org.

Management Goal

Data generated through this program can be used to document trends in fish species abundance over time. These data can also be used to inform decision makers on the relative importance of shorelines on various species of fish.

Hypothesis and Test Statistics

Differences between bays are assessed using pairwise Wilcoxon rank sum tests ($\alpha = 0.05$). The null hypothesis is that there are no differences in fish abundance between the Inland Bays. Correlations between physical parameters and recreationally or commercially important species' abundance are assessed with a Kendall's Tau correlation tests ($\alpha = 0.05$). Geometric mean catch per year are calculated for each fish species caught, and these data will be used to assess species trends over time.




FIGURE C.14 - Map of Inshore Fish and Blue Crab Survey seining sites.

C.6.6 RECREATIONAL FISHING SURVEYS

Description/Objective(s)

The Marine Recreational Information Program (MRIP) survey is used by NOAA Fisheries to quantify and report marine recreational fishing catch and effort. These surveys document pounds of fish caught per trip, number of fishing trips, and pounds of individual fish species caught per year. Data for the Inland Bays are based on surveys conducted by NOAA at the Indian River Inlet.

Responsible Organization and Contact

NOAA Fisheries Service is responsible for this program. DNREC's Division of Fish and Wildlife requests and analyzes local data each year.

Contact: Gordon Colvin, NOAA
301-427-8118, Gordon.colvin@noaa.gov

Data Collection Cost

Not available.

Data Use

Locally, the data are used primarily by DNREC Division of Fish and Wildlife Fisheries Section to assess the impact of recreational fishing on Delaware's fish species, and adjust management decisions accordingly.

Record of Collection

2004 to present. Recreational fishing survey data exist prior to 2004, but collection used a method differing from MRIP.

Collection Method

Data are collected through field, harvester-intercept interview surveys that record catch rates for species from anglers, and a telephone (or mail survey) that is designed to estimate effort. Average catch rates are applied to the effort estimates to generate landings per year. The MRIP survey also takes into account potential bias due to differences in catch rate at high-activity or low-activity sites, or the amount of fishing occurring at different parts of the day.

Analysis Methods

Each estimate is a combination of catch rates and effort in a particular waterbody. Previously, all fishing locations were treated equally, and in some cases high activity sites were sampled much more than low activity sites in an effort to maximize the amount of data being collected. However, the MRIP survey takes into account site activity and other bias associated with the time the survey was conducted to more accurately arrive at a catch estimate.

Data Location

Data are available from DNREC Division of Fish and Wildlife Fisheries Section upon request.

Management Goal

To incorporate the data into stock assessments, thereby accounting for the impact recreational fishing has upon each species.

Hypothesis and Test Statistics

N/A

C.6.7 HARD CLAM SURVEYS

Description/Objective(s)

The hard clam is the most important commercial resource in the Inland Bays, and along with the blue crab, is the most important recreational shellfishery. To enhance the wellbeing of this shellfishery and maintain optimum levels of sustainable harvest an adequate knowledge of the hard clam stock and its variability are necessary. A vacuum suction dredge survey of Rehoboth Bay and Indian River Bay over two years were performed to support these goals and provide information with which to evaluate overall bay health and this important living resource. Objectives of this survey were to determine density and distribution of hard clams and bay scallops within Delaware's Inland Bays, evaluate clam recruitment and

survival since previous surveys were conducted, and evaluate study results and effort in light of management and monitoring objectives.

Responsible Organization and Contact

DNREC Division of Watershed Stewardship

Contact: Michael Bott, Environmental Scientist
Michael.Bott@state.de.us, 302-739-9939

Data Collection Cost

\$10,000 per survey.

Data Use

Data is used primarily by DNREC Division of Fish and Wildlife and the Center for the Inland Bays. Data generated by this survey can be used to infer the health of the Inland Bays hard clam shellfishery, and inform the shellfish aquaculture program of potential conflicts in aquaculture sites due to high hard clam densities.

Record of Collection

1967, 1976, 2011

Collection Method

The Venturi Suction Dredge survey method was used for this survey. The dredge was fabricated from aluminum with a mesh net attached capable of capturing clams greater than 8mm. The dredge was powered by a 4", water pump with a 4" reinforced intake hose and 3" effluent hose attached to the Venturi dredge. The sample quadrat was a 1m² frame constructed from 1" PVC piping with holes drilled into the material to allow the pipe to fill with water and stay weighted on the sediment. The sample stations duplicated the survey points used for the 1967 and 1976 surveys. Sample stations which were located near the Indian River Inlet or in navigational channels were either eliminated or moved to adjacent locations due to safety hazards. In the event a sample location was too near to a shore line to allow operation of the dredge, the quadrat was located as near to the original point as possible and the new coordinates were recorded. Sample stations were located using a hand held GPS to get within 3m of the location, and a weighted buoy was thrown randomly to mark the specific site. Two divers would place the quadrat on the northern side of the buoy weight and remove and record any macro-algae present. One diver would operate the Venturi dredge, excavating all substrate to a minimum depth of 12". The second diver stabilized the quadrat to prevent movement and verified the station was completely sampled before surfacing. If the substrate could not be excavated to a depth of 12", the actual depth and limiting factors were recorded. Any clams which were partly located within the inside portion of the quadrat were included in the sample. Clams were included if the diver could run a finger along the inside edge of the PVC pipe and feel the shell within the quadrat. To avoid sampling additional substrate which caved into the sample area, the diver did not re-dredge any portions which were already sampled. Once samples were brought to the surface, all live bi-valve molluscan shellfish were sorted from the material, and measurements of the shell width and length, along with species were recorded. Qualitative sediment type was also recorded for each site, along with water depth. All calipers used were frequently calibrated on a known standard.

Analysis Methods

The data was analyzed using the Kolmogorov-Smirnov test to measure differences in clam distributions between the 1976 survey and the 2011 survey in the Rehoboth Bay, and Indian River Bay. Differences were also assessed between Rehoboth Bay and Indian River Bay for the 2011 survey.

Data Location

Upon request to DNREC Division of Fish and Wildlife.

Management Goal

This survey can be used to inform decision makers on the current status of hard clams in the Inland Bays in reference to historical levels. Data generated through this survey details the current health of the hard clam population, if restoration efforts are needed, how current management efforts are impacting hard clams, and where the highest densities of hard clams reside in the Inland Bays. This data can also be used to prevent conflicts between aquaculture sites and high density hard clam beds.

Hypothesis and Test Statistics

Kolmogorov-Smirnov tests were used to assess differences between the 1976 survey and the 2011 survey, along with differences in hard clam abundance between bays in the 2011 survey.

C.6.8 VOLUNTEER HORSESHOE CRAB SURVEY AND TAGGING PROGRAM

Description/Objective(s)

Due to their importance to both the medical industry and migratory birds, horseshoe crab spawning activity is monitored through a volunteer spawning survey, at five sandy beaches in the Inland Bays from May through June (Figure C.15). The objectives of the survey are to assess the importance of the Inland Bays' horseshoe crab population in regards to the regional population, track the number of spawning crabs over time, and monitor horseshoe crab movement through a long-term tagging program.

Responsible Organization and Contact

The Delaware Center for the Inland Bays is responsible for this monitoring.

Contact: Andrew McGowan, Environmental Scientist
environment@inlandbays.org, 302-226-8105, x112

Data Collection Cost

Equipment costs: \$150 annually to support five teams

Staff time: \$5,400 annually

Data Use

Data from this survey is used primarily by the Center for the Inland Bays and DNREC Division of Fish and Wildlife. The data generated through this survey can help determine the size of the Inland Bays' horseshoe crab population, relative to the regional population, measure if their numbers are increasing or decreasing over time, and identify important spawning areas that need to be managed and protected.

Record of Collection

2015 - present (Surveys began in 2012, but the protocol was switched in 2015 to facilitate comparisons with the Delaware Bay Survey).

Collection Method

Sampling occurs on five beaches in the Inland Bays two nights prior to, the night of, and two nights following the new and full moons from May through June. Sampling begins during the



FIGURE C.15 - Map of Inland Bays horseshoe crab survey and tagging sites. The Ellis Point beach was dropped from the survey in 2015.

nighttime high tide as the tide begins to recede. Teams begin by randomly selecting one end of the beach to start from using a coin flip. Once the end of the beach is determined, the team will walk to that end and extend a pull rope with markings every 1m, at the high tide line towards the opposite end of the beach. The length of the pull rope is dependent on the length of the beach, and is designed to systematically allow the placement of 100 1m² quadrats along the beach. The length of the rope is determined by dividing the overall length of the beach by 50. James Farm, Coastal Kayak, and Tower Road all use a 4 meter pull rope. Bay Colony uses a 6 meter pull rope. Peninsula does not use a pull rope because the length of the beach is only 100m; and therefore all quadrats along the beach are counted. In addition to randomizing the direction of travel, the placement of the quadrats within each rope pull is randomized for a single night. Two quadrats are sampled per rope pull, for a total of 100 quadrats. The same two randomized locations along the pull rope are used for the duration of the night. Once the pull rope has been extended, the 1m² quadrat is placed at the first random quadrat location for that given night. The quadrat is positioned so that one end is even with the line of crabs, and the other end is towards the bay. All crabs which have at least half their body inside the quadrat are sexed and counted. Upon completion of the first quadrat, the team moves the quadrat to the second randomly selected location and repeats the counting process. Once the two quadrats have been counted for the first rope pull, the rope is extended along the next portion of the beach, and the same two random quadrat locations are sampled. This is repeated until 100 quadrats have been sampled. The 'Horseshoe Crab line' that is followed is not a straight line, and it may be above or below the water line, however, it is never more than 1m² away from the high tide line.

At each of the sites, salinity samples are also taken for each night. These samples are later tested for salinity. Air and water temperature are also taken with a thermometer.

In addition to counting crabs, 1,000 crabs are also tagged with US Fish and Wildlife Service tags on the left posterior portion of the prosoma. The tagged crabs are sexed and measured for carapace width.

Analysis Methods

Average spawning densities per 1m² are calculated for each beach by dividing the total number of crabs per night by 100 (the number of quadrats), and averaging each night to obtain one spawning density per beach. The cumulative spawning density for the Inland Bays represents the average of the five beaches' averages. Spawning index is calculated in the same manner but includes only females.

Total crab abundance is compared between sites using a pairwise Wilcoxon rank sum test ($\alpha = 0.05$).

Data Location

Located at the Delaware Center for the Inland Bays, available upon request.

Management Goal

The data generated through this survey can be used to identify important areas for horseshoe crab spawning activity, assess overall trends in horseshoe crab spawning activity over time, and inform decision makers on the movements of horseshoe crabs over the course of a single spawning season or multiple years.

Hypothesis and Test Statistics

Correlations between physical parameters and crab abundance are assessed with Kendall's tau correlation tests ($\alpha = 0.05$). Differences in crab abundance between sites is assessed with pairwise Wilcoxon rank sum tests ($\alpha = 0.05$).

C.6.9 DELAWARE BREEDING BIRD ATLAS

Description/Objective(s)

The Delaware Breeding Bird Atlas is primarily a volunteer citizen science project. From 1983 through 1987, over 100 volunteers participated in compiling data for Delaware's first breeding bird atlas, Delaware's largest and most comprehensive ornithological project. This effort, incorporated in *The Birds*

of Delaware (Hess et al., 2000), summarized the distribution of the breeding birds during those years along with species accounts and data about all birds that occurred, or thought to have occurred, in the State. A second Atlas project was conducted between 2008 and 2012.

Responsible Organization and Contact

DNREC Division of Fish and Wildlife, Wildlife Species Conservation & Research Program

Contact: Anthony Gonzon, Project Coordinator

Phone: (302)-653-2880, ext. 123, Email: Anthony.Gonzon@state.de.us

Data Collection Cost

The second Atlas project was funded through the Delaware Division of Fish and Wildlife with funding from the State Wildlife Grants Program, Division of Federal Assistance, United States Fish & Wildlife Service.

Data Use

The data are useful for tracking trends in populations and diversity that occur with changes in land use, habitats, and climate.

Since being published, the first atlas has become an important resource, providing much of the distribution data about Delaware's breeding avifauna. It is often used by researchers, scientists, government officials, and birders. The first Delaware breeding bird atlas established the baseline data that will be used to compare and examine changes in the distributions of breeding species upon completion of future atlas projects.

Record of Collection

Surveys for the first Breeding Bird Atlas project were conducted from 1983-1987. The second set of surveys were conducted from 2008-2012.

Collection Method

Volunteer atlasers are assigned to a geographic "block" to survey. Within that block, the atlaser attempts to confirm breeding for as many species as possible. A block is approximately 10 sq. miles and is created using a standardized grid that is overlaid onto a map of Delaware. Delaware has over 265 blocks divided into six regions in the State. Each of the six regions contains more than 40 blocks.

Each atlaser spends time in their block surveying during the breeding season (for most birds, March - July), and observing and recording data about the birds they encounter on field cards. Volunteer atlasers report this data using an online, interactive data entry application or by submitting their field cards for entry. Once the data are entered the volunteers "save" it to the database for verification. A regional compiler serves as the primary contact for each region and verifies data. The project coordinator is responsible for maintaining communications between volunteers, compilers, project supporters, landowners, and others, and also handles all of the financial and logistical issues of the atlas, including reports and data analyses.

Analysis Methods

Maps are produced for each breeding species along with other important information such as breeding safe dates, nesting dates, and arrival and departure estimates.

Data Location

Data are kept by the DNREC Wildlife Species Conservation & Research Program. Data may be accessed online through the Breeding Bird Atlas Explorer, maintained by the USGS Patuxent Wildlife Research Center (<http://www.pwrc.usgs.gov/bba>).

C.6.10 MID-WINTER WATERFOWL SURVEYS

Description/Objective(s)

For almost 40 years, the Division of Fish and Wildlife has conducted four aerial waterfowl surveys annually to measure long-term trends in duck and goose populations. Flights are usually made in mid-October, mid-November, mid-December and the second week in January, subject to weather and mechanical delays. The January flight is part of a coast-wide effort to survey waterfowl throughout the Atlantic Flyway at approximately the same time. The state surveys cover the primary waterfowl habitat in Delaware, approximately the eastern half of the state, and are divided into 11 zones.

Not all ducks and geese can be seen equally well from a plane. The surveys give fairly accurate information about geese, but duck populations such as wood ducks and sea ducks are almost impossible to count.

The important feature of these counts is that they augur for long-term trends that are useful to measure changes in waterfowl management strategies and the environment. In most cases no single survey count is especially important in itself, but cumulative counts have revealed important changes over the years.

Responsible Organization and Contact

DNREC Wildlife Species Conservation and Research Program

Contact: Joseph Rogerson
joseph.rogerson@state.de.us, 302-735-3600

Data Collection Cost

Not available.

Data Use

Data are used by the US Fish and Wildlife Service and the Delaware Division of Fish and Wildlife to adjust hunting regulations in response to population trends.

Data from both the state and Atlantic Flyway counts are used to produce a winter waterfowl indicator for the *State of the Delaware Inland Bays* reports.

Record of Collection

1974 to present.

Collection Method

These surveys are carried out via a small plane with a pilot and biologist aboard, taking similar routes and using the same techniques each time.

Analysis Methods

Waterfowl counts are summarized within the following zones:

- Zone 1** - (Width of the State) Pennsylvania Line to the Delaware Memorial Bridge.
- Zone 2** - (Width of the State) Delaware Memorial Bridge to the C&D Canal.
- Zone 3** - (Width of the State) C&D Canal to a Line from Liston Point to Kenton.
- Zone 4** - (Width of the State) Liston Point to Route 6 (Smyrna).
- Zone 5** - (Width of the State) Route 6 to Route 8 Port Mahon.
- Zone 6** - (East of 113) Route 8 to Big Stone Beach.
- Zone 7** - (East of Route 1) Big Stone Beach to Broadkill River.
- Zone 8** - (East of Route 1) Broadkill River to Bend in Lewis Rehoboth Canal South of Gordon's Pond.
- Zone 9** - South of Gordon's Pond and All of Rehoboth Bay.
- Zone 10** - All Indian River Bay West to Millsboro and South to Salt Pond.
- Zone 11** - Salt Pond South to Maryland Line and West to Route 17.

Zones 9, 10, and 11 fall within the Inland Bays watershed.

Data Location

DNREC Division of Fish and Wildlife. Data also are posted to the Delaware Open Data Site (<https://data.delaware.gov/Energy-and-Environment/Aerial-Waterfowl-Survey-Data/bxyv-7mgn>).

C.6.11 BALD EAGLE & OSPREY NESTING SURVEYS

Description/Objective(s)

The Delaware Division of Fish and Wildlife's Species Conservation and Research Program (SCRP) monitors bald eagle and osprey populations in the state.

State monitoring efforts are supplemented by data collected through the Citizen Osprey Monitoring Program, managed by the SCRCP, in which volunteers monitor osprey platforms and nests. As of 2016, the volunteer monitoring data are submitted to the OspreyWatch program (www.ospreywatch.org), but state data are still analyzed by SCRCP. The mission of OspreyWatch is to collect information on a large enough spatial scale to be useful in addressing global climate change, depletion of fish stocks, and environmental contaminants.

The U.S. Geological Survey biologists tested eggs and chicks for contaminants in 2001. Historically, osprey surveys were concentrated in the Inland Bays and Nanticoke River system, but the entire state of Delaware was surveyed in 2003, 2007 and 2014.

Responsible Organization and Contact

DNREC Division of Fish and Wildlife, Wildlife Species Conservation & Research Program

Contacts: Anthony Gonzon (Bald Eagles)
Anthony.Gonzon@state.de.us, 302-653-2880, ext. 123

Kate Fleming (Ospreys)
Kate.Fleming@state.de.us, 302-735-8658

Data Collection Cost

Not available.

Data Use

The data for Delaware are analyzed by SCRCP staff to improve understanding of arrival dates, nest success and other aspects of the species' biology in the state. Information supports conservation decisions and assists the state in identifying any new or emerging issues.

Bald Eagle and Osprey nesting are used as indicators in the *State of the Delaware Inland Bays* reports. Both species are high on the food chain and eat foods - primarily fish - that accumulate toxins. Bald eagle nesting is influenced by habitat disturbance. Ospreys also rely mostly on coastal habitats and may be affected by climate change and sea level rise.

Record of Collection

Data used for indicator reports: Bald Eagles: 1987 to present.
Ospreys: 1991 to present.

The Division has conducted surveys in some form since 1970 to document Osprey nest success. The SCRCP decided to end state-wide Osprey surveys on an annual basis in 2007, with the intention of returning to it every five years or so.

Collection Method

Eagle nests are counted by aerial survey. The only state-wide Osprey nest survey conducted since 2007 was in 2014, and it was done entirely from helicopter. Each site was visited only once between June 20th and July 10th. In prior years, nest sites were visited at least twice, so the number of nests documented in 2014 likely is low, as any early season failed nesting attempts would have missed.

Analysis Methods

Starting in 2016, volunteers provide osprey nest observations to the OspreyWatch online reporting application, hosted by the Center for Conservation Biology at <http://www.osprey-watch.org/>. The data for Delaware are analyzed by SCRCP staff to improve understanding of osprey arrival dates, nest success and other aspects of osprey biology in the state.

Data Location

Data are managed by the SCRCP, and are available upon request to the Program.

Starting in 2016, volunteers provide osprey nest observations to the OspreyWatch online reporting application, hosted by the Center for Conservation Biology at <http://www.osprey-watch.org/>. Delaware and Inland Bays data may be viewed on this site.

APPENDIX D

ASSESSMENT, LISTING, AND REPORTING METHODOLOGIES FOR 305(B) REPORT AND 303(D) LIST DEVELOPMENT, 2016

State of Delaware Ambient Surface Water Quality Monitoring Program - FY 2016

Department of Natural Resources and Environmental Control Watershed Assessment and
Management Branch

Table 1 Station Locations, Descriptions Parameters and Sampling Frequency

Executive Summary

Delaware's Surface Water Quality Monitoring Program for Fiscal Year 2016 is described in this report. Delaware maintains a General Assessment Monitoring Network (GAMN) of 134 stations. GAMN stations are considered long term stations whose data is used to do long term status and trend assessments of water quality conditions of the State's surface waters and support compilation of Watershed Assessment Reports as mandated by the Clean Water Act under section 305(b). This plan implements an updated monitoring strategy that monitors 22 stations monthly, and the remaining stations either 6 or 12 times a year on a rotating basin basis.

Table 1 Station Locations, Descriptions Parameters and Sampling Frequency

Ambient Surface Water Quality Monitoring Program - FY 2015

The purpose of the Ambient Surface Water Quality Monitoring Program is to collect data on the chemical, physical and biological characteristics of Delaware's surface waters. The information that is collected under this Program is used to:

- Describe general surface water quality conditions in the State;
- Identify long term trends in surface water quality;
- Determine the suitability of Delaware surface waters for water supply, recreation, fish and aquatic life, and other uses;
- Monitor achievement of Surface Water Quality Standards;
- Identify and prioritize high quality and degraded surface waters;
- Calculate annual nutrient loads and track progress toward achieving Total Maximum Daily Loads (TMDLs) targets; and
- Evaluate the overall success of Delaware's water quality management efforts.

There are five major components to Delaware's Surface Water Quality Monitoring Program:

- General Assessment Monitoring
- Biological Assessment Monitoring
- Toxics in Biota Monitoring
- Toxics in Sediment Monitoring
- Monitoring under the Watershed Approach to Toxics Assessment and Restoration (WATAR) Plan.

This report discusses the General Assessment Monitoring network and plan. The current Toxics in Biota Monitoring Plan, Toxics in Sediment Monitoring Plan and the WATAR plan are all available on request.

Table 1 Station Locations, Descriptions Parameters and Sampling Frequency

Figure 1. State of Delaware Basins




Table 1 Station Locations, Descriptions Parameters and Sampling Frequency

Part I The General Assessment Monitoring Network (GAMN)

The General Assessment Monitoring Network (GAMN) provides for routine water quality monitoring of surface waters throughout Delaware. Each station is monitored for conventional parameters such as nutrients, bacteria, dissolved oxygen, pH, alkalinity, and hardness. Some stations are monitored for dissolved metals. See tables 2, 3 and 4 for parameters and methods. The data from this monitoring is entered into the STORET database, is reviewed and then analyzed in assessing the water quality of each basin for the Watershed Assessment Report (CWA Section 305 (b) Report).

The plan provides for monitoring at stations within each watershed in the state. The network was recently reviewed and updated. The review is discussed in section I.1. See also Table 1: FY 2016 Monitoring Plan and Schedule.

I.1 Changes for Surface Water Quality Monitoring Plan

Over the past several years, a main objective of the Watershed Assessment and Management Section's Ambient Surface Water Quality Monitoring Program was to collect water quality data that could be used for developing and calibrating hydrodynamic and water quality models. These models were used to establish Total Maximum Daily Loads (TMDLs) for nutrients and bacteria in impaired waters of the State.

Now, with the establishment of nutrient and bacteria TMDLs for most impaired waters of the State, a major objective of the Ambient Surface Water Quality Monitoring Program is to collect appropriate data that can be used to track water quality changes and to determine if TMDL requirements are being met.

Considering this (and other emerging) needs, and since the Department's monitoring budget is limited, the surface water quality monitoring plan has been prepared with the following changes: Monitoring stations in earlier monitoring plans were reviewed to determine which stations were critical to meet data needs and which could be dropped. The retained stations fall into 2 categories;

- Stations were assigned to one of the following categories:
- a. C1 – Category 1 stations are high priority stations that will be used for calculating annual loads and/or long-term trends. These stations are generally co-located with a USGS stream gaging station, or are located at the mouth of a tidal river. Because of importance of these stations, monitoring at these stations will be conducted monthly, regardless of priority basin schedule (22 stations)
 - b. C2 – The remaining stations are part of Category 2 stations and monitoring frequency at these stations follow Priority Basin schedule.
2. A Rotating Basin Monitoring Plan has been implemented. In this scheme of monitoring, the State is divided into 5 Monitoring Basins. Every year, two of the Basins are considered "Priority Basins" and all stations in a Priority Basin are monitored monthly.

Table 1 Station Locations, Descriptions Parameters and Sampling Frequency

Monitoring for stations in other basins will be conducted bimonthly. Priority Basin monthly monitoring will be conducted according to the following schedule:

- a. FY 2014 – Lower Delaware River/Bay, Piedmont
- b. FY 2015 – Piedmont, Chesapeake
- c. FY 2016 – Chesapeake, Inland Bays
- d. FY 2017 – Inland Bays, Upper Delaware River/Bay
- e. FY 2018 – Upper Delaware, Lower Delaware River/Bay

I.2 Objectives

The objective of this monitoring is to collect water quality data for status and trends assessment on all basins within Delaware. The data will also be compared to water quality standards to assess designated use support, as mandated by Section 305(b) of the Clean Water Act. In addition, the data will be used to calculate annual nutrient loads and to track progress toward achieving TMDL targets.

I.3 Scope of Monitoring

Table 1 provides a listing of all stations to be monitored during FY 2015.

Table 2 provides a listing of parameters that will be monitored at all stations in the network. Stations shown for metals testing in Table 1 shall be sampled according to the specifications in Table 3.

The Department is in the process of bringing the Environmental Laboratory's recently purchased ICP/MS instrument online and expects to update metals sampling protocols in the near future. The monitoring plan will be adjusted as those details are known.

Table 4 shows additional parameters needed for use in the Biotic Ligand Model that is part of the Surface Water Quality Standards for copper in freshwaters.

Part II Special Project Monitoring

Special project monitoring is needed from time to time in specific watersheds to address specific concerns. These projects are generally short term in nature. The Department is not conducting any special projects during the FY 2016 monitoring year.

II.1 Special Surveys

The purpose of special survey monitoring is to collect data that are not obtained using other monitoring activities and are needed for modeling purposes as described above. Special surveys include deployment of continuous monitors (YSI Data Sondes) and sediment sampling. No special survey sediment sampling is called for in this monitoring year.

II.2 Continuous Monitoring

The Department is implementing a network of continuous water quality monitoring stations to collect data for dissolved oxygen and other parameters several times each day using YSI (or similar) datasondes. The Department is cooperating with Delaware Geological Survey (DGS) and the United States Geological Survey (USGS) in operating a number of continuous monitors in the

Table 1 Station Locations, Descriptions Parameters and Sampling Frequency

State. The information from these continuous monitoring sites are available on real-time basis via the USGS website and via the Delaware Environmental Observing System (DEOS) website.

Part III Field and Laboratory Procedures

Field procedures for sample collection activities are detailed in the Quality Assurance Management Plan, Environmental Laboratory Section. Method references, STORET codes and reporting levels for parameters listed in Table 2 are from an Access database maintained by the Environmental Laboratory Section. Any deviation from standard field, laboratory procedures, or this sampling plan shall be documented with a complete description of the alteration.

Boat run surveys

Boat run surveys should be conducted within one day of tributary sampling in the watershed.

Part IV Quality Assurance, Documentation, Data Usage and Reporting

The quality assurance objectives and quality control procedures for these surveys are documented in the Quality Assurance Management Plan, Environmental Laboratory Section. A duplicate water column sample will be collected and analyzed on 10% of the samples from this project. All analytical results from the duplicate analyses shall be reported with the other data.

All analytical results shall be reported to the Watershed Assessment and Management Section digitally and on paper (using standard Environmental Laboratory Section data report forms).

Table 1 Station Locations, Descriptions Parameters and Sampling Frequency

STATION INFORMATION - FY 2016	STORET #	Type	GAMN	Cu, Pb & Zn	As	Fe	DIN & DIP	Storm Sampling	No. of Samples for FY 2016	Boat Run
PIEDMONT DRAINAGE										
Brandywine Creek										
Brandywine Creek @ Foot Bridge in Brandywine Park	104011	C2	✓	✓					6	
Brandywine Creek @ New Bridge Rd. (Rd. 279)(USGS gage 01481500)	104021	C1	✓	✓				4 storms	12	
Brandywine Creek @ Smith Bridge Rd. (Rd. 221)	104051	C2	✓	✓					6	
Christina River										
Christina River beneath Rt. 141 in Newport off Water St.	106021	C2	✓	✓					6	
Little Mill Creek @ DuPont Rd.	106281	C2	✓	✓					6	
Christina River @ Conrail Bridge (USGS tide gage 01481602)	106291	C1	✓	✓					12	
Christina River @ Nottingham Rd. (Rt. 273) above Newark	106191	C2	✓	✓					6	
Christina River @ Sunset Lake Rd. (Rt. 72) (USGS 01478000 at Cooches bridge)	106141	C1	✓	✓				4 storms	12	
Smalleys Dam Spillway @ Smalleys Dam Rd.	106031	C2	✓	✓					6	
Red Clay Creek										
Red Clay Creek @ W. Newport Pike (Rt. 4) Stanton (USGS gage 01480015)	103011	C2	✓	✓					6	
Burrough's Run @ Creek Rd. (Rt 82)	103061	C2	✓	✓					6	
Red Clay Creek @ Barley Mill Rd. (Rd. 258A) Ashland	103041	C2	✓	✓					6	
Red Clay Creek @ Lancaster Pike (Rt. 48) Wooddale (USGS gage 01480000)	103031	C1	✓	✓				4 storms	12	
White Clay Creek										
White Clay Creek @ Delaware Park Blvd. (Race Track) (USGS gage 014790000)	105151	C1	✓	✓				4 storms	12	

Table 1 Station Locations, Descriptions Parameters and Sampling Frequency

STATION INFORMATION - FY 2016	STORET #	Type	GAMN	Cu, Pb & Zn	As	Fe	DIN & DIP	Storm Sampling	No. of Samples for FY 2016	Boat Run
White Clay Creek @ McKees Lane	105171	C2	✓	✓					6	
White Clay Creek @ Chambers Rock Rd. (Rd. 329)	105031	C2	✓	✓					6	
Naamans Creek										
Naaman Creek @ State Line near Hickman Rd.	101021	C2	✓						6	
S. Branch Naaman Creek @ Darley Rd. (Rd. 207)	101031	C2	✓						6	
Naamans Creek at Rt 3 (Marsh Road)	101061	C2	✓						6	
Shellpot Creek										
Shellpot Creek @ Hay Rd. (Rd. 501)	102041	C2	✓			✓			6	
Rt. 13 Bus (Market Street) Bridge, USGS station is located about 700 ft downstream.	102051	C1	✓					4 storms	12	
Shellpot Crk at Carr Road Bridge	102081	C2	✓						6	
CHESAPEAKE BAY DRAINAGE										
Chester River										
Sewell Branch @ Sewell Branch Rd. (Rd. 95)	112021	C2	✓						12	
Choptank River										
Cow Marsh Creek @ Mahan Corner Rd. (Rd. 208)	207021	C2	✓						12	
Tappahanna Ditch @ Sandy Bend Rd. (Rd. 222)	207081	C2	✓						12	
Culbreth Marsh Ditch @ Shady Bridge Rd. (Rd. 210)	207091	C2	✓						12	
White Marsh Branch @ Cedar Grove Church Rd. (Rd. 268)	207111	C2	✓						12	
Marshyhope Creek										
Marshyhope Creek @ Fishers Bridge Rd. (Rd. 308)	302031	C1	✓					8 storms	12	
Nanticoke River										
Nanticoke River @ Sharptown	304011	C2	✓	✓					12	

Table 1 Station Locations, Descriptions Parameters and Sampling Frequency

STATION INFORMATION - FY 2016	STORET #	Type	GAMN	Cu, Pb & Zn	As	Fe	DIN & DIP	Storm Sampling	No. of Samples for FY 2016	Boat Run
Nanticoke River @ buoy 66 (confluence with DuPont Gut)	304151	C2	✓	✓					12	✓
<i>Nanticoke River Tributaries</i>										
Racoon Prong @ Pepperbox Rd. (Rd. 66)	304671	C2	✓	✓					12	
Nanticoke River @ Rifle Range Rd. (Rd. 545)	304191	C1	✓	✓				8 storms	12	
Concord Pond @ German Rd. (Rd. 516)	304311	C2	✓	✓					12	
Williams Pond @ East Poplar St. (across from Hospital)	304321	C2	✓	✓					12	
Bucks Branch @ Conrail Rd. (Rd. 546)	304381	C2	✓	✓					12	
Nanticoke River @ Rt. 13	304471	C2	✓	✓					12	
Records Pond @ Willow St.	307011	C2	✓	✓					12	
Horseys Pond @ Sharptown Rd. (Rt. 24)	307171	C2	✓	✓					12	
Gravelly Branch @ Coverdale Rd. (Rd. 525)	316011	C2	✓	✓					12	
Trap Pond on Hitch Pond Branch @ Co. Rd. 449 or Trap Pond Rd	307081	C2	✓	✓					12	
Deep Creek above Concord Pond, near Old Furnace at Rd. 46	304591	C2	✓	✓					12	
Gravelly Branch at Deer Forest Road (Rd 565) on west edge of Redden State Forest Jester Tract	316031	C2	✓	✓					12	
Broad Creek at Main Street in Bethel (Rd 493)	307031	C2	✓	✓					12	
Nanticoke River at Beach HWY (Ellendale Greenwood HWY) on east edge of Greenwood	304681	C2	✓	✓					12	
Clear Brook at Rt 18 (Bowdens Garage Rd)	304491	C2	✓	✓					12	
<i>Pocomoke River</i>										
Pocomoke River @ Bethel Rd. (Rd. 419)	313011	C2	✓						12	
<i>DELAWARE BAY DRAINAGE</i>										
<i>Appoquinimink River</i>										

Table 1 Station Locations, Descriptions Parameters and Sampling Frequency

STATION INFORMATION - FY 2016	STORET #	Type	GAMN	Cu, Pb & Zn	As	Fe	DIN & DIP	Storm Sampling	No. of Samples for FY 2016	Boat Run
Drawyer Creek off DuPont Parkway. (Rt. 13) at parking area	109071	C2	✓	✓					6	
Shallcross Lake @ Shallcross Lake Rd. (Rd. 428)	109191	C2	✓	✓					6	
Noxontown Pond @ Noxontown Rd. (Rd. 38)	109131	C2	✓	✓					6	
Appoquinimink River @ DuPont Prkwy. (Rt. 13)	109041	C2	✓	✓					6	
Appoquinimink River @ MOT Gut (west bank)	109171	C2	✓	✓					6	
Deep Creek Br of Appoquinimink River at Rt. 71 Bridge (Middletown Natural Area), duplicate with 109081	109251	C1	✓	✓				4 storms	12	
Appoquinimink River @ Silver Run Rd. (Rt. 9) NE side	109121	C2	✓	✓					6	
Appoquinimink River @ confluence with Delaware River	109091	C1	✓	✓					12	✓
Army Creek										
Army Creek @ River Rd. (Rt. 9)	114011	C2	✓						6	
Chesapeake & Delaware Canal										
C & D Canal @ DuPont Pky. (Rt. 13) St. Georges Bridge	108021	C2	✓						6	✗
Lums Pond @ Boat ramp	108111	C2	✓						6	
Dragon Run										
Dragon Creek @ Wrangle Hill Rd. (Rt. 9)	111011	C2	✓						6	
Dragon Creek @ S. DuPont Hgwy. (Rt. 13)	111031	C2	✓						6	
Red Lion Creek										
Red Lion Creek @ Bear Corbitt Rd. (Rt. 7)	107011	C2	✓						6	
Red Lion Creek @ Rt. 9	107031	C2	✓						6	
Blackbird Creek										
Blackbird Creek, Road 463 East of RR Tracks. USGS gage	110011	C1	✓					4 storms	12	
Blackbird Landing Rd 455	110031	C2	✓						6	

Table 1 Station Locations, Descriptions Parameters and Sampling Frequency

STATION INFORMATION - FY 2016	STORET #	Type	GAMN	Cu, Pb & Zn	As	Fe	DIN & DIP	Storm Sampling	No. of Samples for FY 2016	Boat Run
Blackbird Creek @ Taylors Bridge Rd. (Rt. 9)	110041	C2	✓						6	
Leipsic River										
Garrisons Lake @ DuPont Highway (Rt. 13)	202021	C2	✓						6	
Leipsic River @ Denny St. (Rt. 9)	202031	C2	✓						6	
Upstream of Masseys Millpond at Rt. 15	202191	C2							6	
Little River										
Little River @ Bayside Dr. (Rt.9)	204031	C2	✓						6	
Little River @ N. Little Creek Rd. (Rt. 8)	204041	C2	✓						6	
Smyrna River										
Mill Creek @ Carter Rd. (Rd. 137)	201021	C2	✓						6	
Smyrna River @ Rt. 9 (Flemings Landing)	201041	C2	✓						6	
Duck Creek @ Smyrna Landing Rd. (Rd. 485)	201051	C2	✓						6	
201011 Mill Creek at Rt. 13	201011	C2	✓						6	
Providence Creek @ Duck Creek Rd. (Rt.15)	201161	C2	✓						6	
Broadkill River										
Ingram Branch, Savanah Ditch @ Rd. 246	303011	C2	✓						6	
Ingram Branch @ Rd. 248	303021	C2	✓						6	
Rt. 5 Bridge	303031	C1	✓					4 storms	12	
Rt. 1 Bridge (Mainstem)	303041	C2	✓						6	
Broadkill River 0.10 Miles From Mouth of Broadkill	303061	C1	✓						6	✓
Red Mill Pond at Rt. 1	303051	C2	✓						6	
Beaverdam Creek at Rd. 88	303171	C2	✓						6	

Table 1 Station Locations, Descriptions Parameters and Sampling Frequency

STATION INFORMATION - FY 2016	STORET #	Type	GAMN	Cu, Pb & Zn	As	Fe	DIN & DIP	Storm Sampling	No. of Samples for FY 2016	Boat Run
Beaverdam Creek above Rd. 259, Hunters Mill Pond	303181	C2	✓						6	
Round Pole Branch at Rd. 88	303311	C2	✓						6	
Waples Pond at Rt. 1	303331	C2	✓						6	
Pemberton Branch at Rt. 30 above Wagamons Pond	303341	C2	✓						6	
<i>Cedar Creek</i>										
Swiggetts Pond @ Cedar Creek Rd. (Rt. 30)	301021	C2	✓						6	
Cedar Creek @ Coastal Hgwy. (Rt. 1)	301031	C2	✓						6	
Cedar Creek @ Cedar Beach Rd. (Rt. 36)	301091	C2	✓						6	
<i>Mispillion River</i>										
Mispillion River @ Rt. 1	208021	C2	✓						6	
Mispillion River/Cedar Creek confluence @ Lighthouse	208061	C1	✓						6	
Abbotts Pond @ Abbotts Pond Rd. (Rd. 620)	208181	C2	✓						6	
Silver Lake @ Maple Ave.	208211	C2	✓						6	
Beaverdam Branch @ Deep Grass Ln. (Rd. 384)	208231	C2	✓						6	
<i>Delaware Bay</i>										
Roosevelt Inlet, Mouth	401011	C2	✓						6	✓
<i>Murderkill River</i>										
Murderkill River @ confluence of Black Swamp Creek at Rt. 13	206011	C1	✓	✓				4 storms	12	
Browns Branch @ Milford - Harrington Hwy. (Rt. 14)	206041	C2	✓	✓					6	
Murderkill River @ Bay Rd. (Rt. 1/113)	206091	C2	✓	✓					6	✓
Murderkill River @ Bowers Beach Wharf (mouth)	206101	C1	✓	✓					6	✓
Murderkill River near levee @ Milford Neck Wildlife Area (3.25 miles from mouth)	206141	C2	✓	✓					6	✓

Table 1 Station Locations, Descriptions Parameters and Sampling Frequency

STATION INFORMATION - FY 2016	STORET #	Type	GAMN	Cu, Pb & Zn	As	Fe	DIN & DIP	Storm Sampling	No. of Samples for FY 2016	Boat Run
Murderkill River @ confluence of Kent County WWTF discharge ditch	206231	C2	✓	✓					6	✓
McColley Pond @ Canterbury Rd. (Rt. 15)	206361	C2	✓	✓					6	
Coursey Pond @ Canterbury Rd. (Rt. 15)	206451	C2	✓	✓					6	
Double Run @ Barretts Chapel Rd. (rd. 371)	206561	C2	✓	✓					6	
St. Jones River										
St. Jones River @ Barkers Landing	205041	C2	✓						6	
St. Jones River @ Rt. 10	205091	C2	✓						6	
Fork Branch @ State College Rd. (Rd. 69)	205151	C2	✓						6	
Moores Lake @ S. State St.	205181	C2	✓						6	
Silver Lake @ Spillway (Dover City Park)	205191	C1	✓					4 storms	12	
St. Jones at Bowers Beach, mouth to Del.Bay.	205011	C1	✓						6	
Derby Pond @ Rt. 13A	205211	C2	✓						6	
INLAND BAYS DRAINAGE										
Tributary Stations										
Burton Pond @ Rt. 24	308031	C2	✓	✓			✓		12	
Millsboro Pond @ Rt. 24	308071	C1	✓	✓			✓	4 storms	12	
Pepper Creek @ Rt. 26 (Main St.)	308091	C2	✓	✓			✓		12	
Blackwater Creek @ Omar Rd. (Rd. 54)	308361	C2	✓	✓			✓		12	
Dirickson Creek @ Old Mill Bridge Rd. (Rd. 381)	310031	C2	✓	✓			✓		12	
Bunting Branch										
Buntings Branch @ Rt. 54 (Polly Branch Rd.)	311041	C2	✓	✓			✓		12	
Guinea Creek										

Table 1 Station Locations, Descriptions Parameters and Sampling Frequency

STATION INFORMATION - FY 2016	STORET #	Type	GAMN	Cu, Pb & Zn	As	Fe	DIN & DIP	Storm Sampling	No. of Samples for FY 2016	Boat Run
Guinea Creek @ Banks Rd. (Rd. 298)	308051	C2	✓	✓			✓		12	
Iron Branch										
Whartons Branch @ Rt. 20 (Dagsboro Rd.)	309041	C2	✓	✓			✓		12	
Lewes & Rehoboth Canal										
Lewes & Rehoboth Canal @ Rt. 9	305041	C2	✓	✓			✓		12	
Little Assawoman Canal										
Little Assawoman Bay @ Rt. 54 (The Ditch)	310011	C2	✓	✓			✓		12	
White Creek @ mouth of Assawoman Canal	312011	C2	✓	✓			✓		12	
Love Creek										
Bundicks Branch @ Rt. 23	308371	C2	✓	✓			✓		12	
Miller Creek										
Beaver Dam Ditch @ Beaver Dam Rd. (Rd. 368)	310121	C1	✓	✓			✓		12	
Stockley Branch/Cow Bridge										
Cow Bridge Branch @ Zoar Rd. (Rd. 48)	308281	C2	✓	✓			✓		12	
Swan Creek										
Swan Creek @ Mount Joy Rd. (Rd. 297)	308341	C2	✓	✓			✓		12	
Vines Creek										
Ocean Boundary Stations										
Lewes & Rehoboth Canal @ Rt. 1	305011	C2	✓	✓			✓		12	
Indian River Inlet @ Coast Guard Station	306321	C1	✓	✓	✓		✓		12	
Boat Run Stations										
Rehoboth Bay @ Buoy 7	306091	C2	✓	✓	✓		✓		12	✓

Table 1 Station Locations, Descriptions Parameters and Sampling Frequency

<i>STATION INFORMATION - FY 2016</i>	<i>STORET #</i>	<i>Type</i>	<i>GAMN</i>	<i>Cu, Pb & Zn</i>	<i>As</i>	<i>Fe</i>	<i>DIN & DIP</i>	<i>Storm Sampling</i>	<i>No. of Samples for FY 2016</i>	<i>Boat Run</i>
Masseys Ditch @ Buoy 17	306111	C2	✓	✓	✓		✓		12	✓
Indian River Bay @ Buoy 20	306121	C1	✓	✓	✓		✓		12	✓
Indian River @ Buoy 49 (Swan Creek)	306181	C2	✓	✓	✓		✓		12	✓
Indian River @ Island Creek	306331	C2	✓	✓	✓		✓		12	✓
Island Creek upper third	306341	C2	✓	✓	✓		✓		12	✓
Little Assawoman Bay Mid-bay (Ocean Park Lane)	310071	C2	✓	✓			✓		12	✓

Table 2 Water Quality Parameters to be analyzed at all Stations in the Monitoring Network, FY 2015

<i>Parameter</i>	<i>Method Reference (EPA)</i>	<i>Reporting Level¹</i>
<i>Water Column Nutrients</i>		
Total Phosphorus	EPA365.1 M	0.005 mg/l P
Soluble Ortho-phosphorus	EPA365.1	0.005 mg/l P
Ammonia Nitrogen	EPA350.1	0.005 mg/l N
Nitrite+Nitrate N	EPA353.2	0.005 mg/l N
Total N	SM 4500 NC	0.08 mg/l N
<i>Carbon and Organics</i>		
Total Organic Carbon	EPA415.1	1 mg/l
Dissolved Organic Carbon	EPA415.1	1 mg/l
Chlorophyll-a (Corr)	EPA 445.0	1 µg/l
<i>Biochemical Oxygen Demand</i>		
BOD ₅ , N-Inhib (CBOD)	SM20 th ed-5210B	2.4 mg/l
BOD ₂₀ , N-Inhib (CBOD)	SM20 th ed-5210B	2.4 mg/l
<i>General</i>		
Dissolved oxygen – Winkler ²	EPA360.2	0.25 mg/l
Dissolved oxygen – Field	EPA360.1	0.1 mg/l
Total Suspended Solids	EPA160.2	2 mg/l
Alkalinity	EPA310.1	1 mg/l
Hardness	EPA130.2	5 mg/l
Field pH	EPA150.1	0.2 pH units
Conductivity – Field	EPA120.1	1 µS/cm
Salinity	SM20 th ed-2520B	1 ppt
Temperature	EPA170.1	°C
Secchi Depth ³	EPA/620/R-01/003	meters
Light Attenuation ⁴	EPA/620/R-01/003	%
Turbidity	EPA180.1	1 NTU

Chloride	EPA325.2	1 mg/l
Bacteria		
Enterococcus	SM20 th ed-9230C	1 cfu/100 ml

- ¹ As documented in the ELS Quality Assurance Management Plan, the ELS defines the Limit of Quantitation (LOQ) as the lowest standard in the calibration curve or, in instances where a standard curve is not specified by the procedure, LOQ represents the limitations of the method. For those tests where reference spiking material exists, the ELS measures Method Detection Limit (MDL), as defined in the Federal Register 40 CFR Part 136 Appendix B. MDL values are generated or verified once per year. Results less than the MDL are considered to be not detected and “< MDL” is reported. Results greater than the MDL but less than the LOQ are qualified with a J to indicate a result that is extrapolated or estimated. For tests where MDL is not applicable, results less than the LOQ are reported as “< LOQ”, ELS MDLs meet or exceed (i.e. are lower than) the reporting level requirements listed in Table 3.
- ² Secchi Depth to be measured at designated stations.
- ³ Light attenuation to be conducted as practical to obtain correlation with Secchi disk readings

Table 3 Metals Parameters

<i>Dissolved Metals (dissolved and total)</i>	<i>Method Reference (EPA)</i>	<i>Reporting Level</i>
Copper	EPA 200.7 M	5.0 ug/l
Lead	EPA 200.7 M	3.0 ug/l
Zinc	EPA 200.7 M	10 ug/l
Iron	EPA 200.7 M	100 ug/l

Table 4 Additional Parameters Needed for Freshwater Stations with Biotic Ligand Model Sampling for Copper

<i>Dissolved Parameters</i>	<i>Method Reference (EPA)</i>	<i>Reporting Level</i>
Alkalinity	APHA 2320	1mg/l
Chloride	APHA 4500-Cl ⁻ E	3mg/l
Calcium	EPA 200.7 M	1000 ug/l
Magnesium	EPA 200.7 M	1000 ug/l
Potassium	EPA 200.7 M	1000 ug/l
Sodium	EPA 200.7 M	1000 ug/l
Sulfate	USEPA 300.0	0.75 mg/l